

**Natural History Museum Library**



**000163746**








S. 3. C. 65.







PHILOSOPHICAL  
TRANSACTIONS,

OF THE

ROYAL SOCIETY

OF

L O N D O N.

V O L. LXXIX. For the Year 1789.

P A R T I.



L O N D O N,

SOLD BY LOCKYER DAVIS, AND PETER ELMSLY,  
PRINTERS TO THE ROYAL SOCIETY.

MDCCLXXXIX.



## A D V E R T I S E M E N T.

**T**HE Committee appointed by the *Royal Society* to direct the publication of the *Philosophical Transactions*, take this opportunity to acquaint the Public, that it fully appears, as well from the council-books and journals of the Society, as from repeated declarations which have been made in several former *Transactions*, that the printing of them was always, from time to time, the single act of the respective Secretaries, till the Forty-seventh Volume: the Society, as a Body, never interesting themselves any further in their publication, than by occasionally recommending the revival of them to some of their Secretaries, when, from the particular circumstances of their affairs, the *Transactions* had happened for any length of time to be intermitted. And this seems principally to have been done with a view to satisfy the Public, that their usual meetings were then continued, for the improvement of knowledge, and benefit of mankind, the great ends of their first institution by the Royal Charters, and which they have ever since steadily pursued.

But the Society being of late years greatly enlarged, and their communications more numerous, it was thought advisable, that a Committee of their members should be appointed to reconsider the papers read before them, and select out of them such as they should judge most proper for publication in the future *Transactions*; which was accordingly done upon the 26th of March 1752. And the grounds of their choice are, and will continue to be, the importance and singularity of the subjects, or the advantageous manner of treating them; without pretending to answer for the certainty of the facts, or propriety of the reasonings, contained in the several papers so published, which must still rest on the credit or judgment of their respective authors.

It is likewise necessary on this occasion to remark, that it is an established rule of the Society, to which they will always adhere, never to give their opinion, as a Body, upon any subject, either of Nature or Art, that comes before them. And therefore the thanks, which are frequently proposed from the Chair, to be given to the authors of such papers as are read at their accustomed meetings, or to the persons through whose hands they receive them, are to be considered in no other light than as a matter of civility, in return for the respect shewn to the Society by those communications. The like also is to be said with regard to the several projects, inventions, and curiosities of various kinds, which are often exhibited to the Society; the authors whereof, or those who exhibit them, frequently take the liberty to report, and even to certify in the public news-papers, that they have met with the highest applause and approbation. And therefore it is hoped, that no regard will hereafter be paid to such reports, and public notices; which in some instances have been too lightly credited, to the dishonour of the Society.



---

---


# C O N T E N T S

O F

## V O L. LXXIX. P A R T I.

- I. **D**ESCRIPTION of an Improvement in the Application of the Quadrant of Altitude to a celestial Globe, for the Resolution of Problems dependant on Azimuth and Altitude. By Mr. John Smeaton, F. R. S.; communicated by Mr. William Wales, F. R. S. Page 1
- II. *Objections to the Experiments and Observations relating to the Principle of Acidity, the Composition of Water, and Phlogiston, considered; with farther Experiments and Observations on the same Subject.* By the Rev. Joseph Priestley, LL.D. F. R. S. P. 7
- III. *Observations on the Class of Animals called, by Linnæus, Amphibia; particularly on the Means of distinguishing those Serpents which are venomous from those which are not so.* By Edward Whitaker Gray, M. D. F. R. S. p. 21
- IV. *Observations on the Dryness of the Year 1788. In a Letter from the Rev. Mr. B. Hutchinson to Sir Joseph Banks,* Bart. P. R. S. P. 37

- V. *On the Method of determining, from the real Probabilities of Life, the Value of a contingent Reversion in which Three Lives are involved in the Survivorship.* By Mr. William Morgan; communicated by the Rev. Richard Price, D. D. F. R. S. p. 40
- VI. *Result of Calculations of the Observations made at various Places of the Eclipse of the Sun, which happened on June 3, 1788.* By the Rev. Joseph Piazzi, C. R. Professor of Astronomy in the University of Palermo; communicated by Nevil Maskelyne, D. D. F. R. S. and Astronomer Royal. p. 55
- VII. *An Account of a bituminous Lake or Plain in the Island of Trinidad.* By Mr. Alexander Anderson; communicated by Sir Joseph Banks, Bart. P. R. S. p. 65
- VIII. *An Account of a particular Change of Structure in the human Ovarium.* By Matthew Baillie, M. D.; communicated by John Hunter, Esq. F. R. S. p. 71
- IX. *Some Account of the Vegetable and Mineral Productions of Boutan and Thibet.* By Mr. Robert Saunders, Surgeon at Boglepoor in Bengal; communicated by Sir Joseph Banks, Bart. P. R. S. p. 79
- X. *A Meteorological Journal kept at the Apartments of the Royal Society, by Order of the President and Council.* p. 113



THE LAWYERS' ASSOCIATION OF THE DISTRICT OF COLUMBIA

MEMORANDUM FOR THE BOARD OF DIRECTORS

RE: M. D. B. B.

Printed in the

OFFICE OF THE SECRETARY

APPROVED AND FORWARDED:

---


THE PRESIDENT and COUNCIL of the ROYAL SOCIETY  
adjudged, for the Year 1788, the Medal on Sir GODFREY  
COPLEY'S Donation, to CHARLES BLAGDEN, M. D. Sec.  
R. S. for his Two Papers on Congelation, printed in the  
last Volume of the Philosophical Transactions.

---

E R R A T U M.

Page 37. line 15. for  $\frac{175}{5}$  read  $\frac{175}{7}$





P H I L O S O P H I C A L

T R A N S A C T I O N S.

---

- I. *Description of an Improvement in the Application of the Quadrant of Altitude to a celestial Globe, for the Resolution of Problems dependant on Azimuth and Altitude. By Mr. John Smeaton, F. R. S.; communicated by Mr. William Wales, F. R. S.*

Read November 20, 1788.

PERHAPS there are few instruments that better fulfil their design in general, or more naturally represent the movements they are intended to explain and illustrate, than the terrestrial and celestial globe, which are also applied to resolve some of the problems of the sphere, which they most readily do. I believe, however, that whoever applies to

them for the last mentioned purpose, will find them more defective in some respects than they are in others.

The difficulty that has occurred in fixing a semicircle, so as to have a center in the *zenith* and *nadir* points of the globe, at the same time that the meridian is left at liberty to raise the pole to its desired elevation, I suppose, has induced the globe-makers to be contented with the *strip* of thin flexible brass, called the *quadrant of altitude*; and it is well known how imperfectly it performs its office.

The improvement I have attempted, is in the application of a *quadrant of altitude*, of a more solid construction; which being affixed to a brass socket of some length, and this ground, and made to turn upon an upright steel spindle, fixed in the zenith, steadily directs the *quadrant*, or rather *arc*, of *altitude* to its true *azimuth*, without being at liberty to deviate from a vertical circle to the right hand or left: by which means the azimuth and altitude are given with the same exactness as the measure of any other of the great circles.

With respect to the horary circle, as the common application seems very convenient on account of the ready adjustment of its index to answer the culmination of any of the heavenly bodies; and as I find that a circle of four inches diameter is capable of an actual and very distinguishable division into 720 parts, answerable to two minutes of time each, which may serve a globe of the largest size; it seems that it should rather be *improved* than omitted; and, if instead of a *pointer*, an index *stroke* is used in the same plane with that of the divisions, the single minutes, and even half minutes, may be readily distinguished.

This globe, though mounted merely as a model for experiment, and only nine inches in diameter, appears capable of bringing out the solution to a quarter of a degree; which, I  
 appre-

apprehend, may be esteemed sufficient not only as a check upon numerical computation, but to come near enough to find stars in the day-time in the field of telescopes, which, having no equatorial motion, are only capable of direction in altitude and azimuth; but from globes of a larger size, we may expect to come proportionably nearer.

*Explanation of the figures, Plate I.*

The figures 1. and 2. being different views of the same things, AB represents a line, in common to both, in the surface of the horizon, which here is of brass.

CD, CD, are vertical lines, supposed to pass through the center of the globe in each figure; and

EFG, EFG, are portions of great circles of the globe.

Fig. 1. supposes the spectator looking at the apparatus of the globe from the south point of the horizon; therefore the circular arch EFG, in this position, will be a part of the *prime vertical*, and the small parallelogram HI is supposed to be a *section* of the brass meridian, according to that vertical plane.

Fig. 2. is a view of the same parts, the spectator being supposed to look at them from the west point of the horizon; and in this position HI is supposed to be a *portion* of the *brass meridian*. This being fixed in mind, in what follows the same letters denote the same parts in both figures.—KLM denotes a piece of brass, or brass carriage, made to fit upon the vertical part of the meridian, and capable of sliding  $5^{\circ}$  on each side of that point, so as to adjust to it, and to fix fast there, by means of the finger screw N\*. This piece of brass carries

\* The holes represented in the portion of the brass meridian (HI, fig. 2.) are screw holes at five degrees distance, in this quarter of the circle, into any of which the finger screw N is to be put as occasion may require; the *fl* allowing sufficiently for adjustment.

the *steel spindle* PQ, which is firmly socketed into it at K, according to the dotted lines *o r*, *o r*. The axis of this spindle is therefore capable of being set upright upon the *zenith point*, and to maintain that position with a sufficient degree of firmness.—Rq, Rq, represents the section of a brass socket made to fit the spindle, and turn round freely upon it; and when home to the shoulder at *oo*, to turn without shake; the socket and spindle being a small matter taper, and *ground* together. On one side of the socket is firmly fixed the arm ST, by screws or folder.—UW is an arch of 80 degrees, serving instead of the quadrant of altitude, and of the same *substance* as the meridian. This is firmly screwed to the arm, and adjusted by construction, so that when the *spindle* is vertical, the face of this arch shall make part of a vertical circle.—This arch being a portion of a circle, of the same diameter as the brass meridian, when its point *zero* at W rests upon the brass horizon, its inside surface is made to agree with that of the horizon by means of a small thin *nib* of brass; that being attached to the inside of the bottom of the quadrant of altitude at W, and projecting a little below it, gently bears against the inside of the horizon, in substance occupying about half the *clearance* between the body of the globe and its surrounding horizon: this *nib*, seen edgewise, is shewn at the letter X. By this means the altitude of the object is shewn upon the working face of the quadrant, and the quadrant's bottom shews the azimuth upon the horizon; at the same time the globe is free to revolve upon its axis, clear of all the circles.

The quadrant might be made complete to 90°; but as in these middle latitudes there is very little business for azimuths when the altitudes are above 80°, and as I judged it eligible, that the quadrant should be made capable of working on both sides the meridian; *that* would be prevented by the necessary thickness

thickness that the circles require to give them solidity, in contradistinction to *mathematical planes*; unless a part of a quadrant was cut out next the vertex to give them clearance: by this means the arch being lifted up from the spindle, and put on the other side of the brass meridian for the afternoon, it will then come within  $10^{\circ}$  or  $15^{\circ}$  of the meridian; and if the use of this space should be wanted, it can be supplied by reversing the similar operation for the morning; and the back side of the upper end of the quadrant at U being champered, or *bevelled* off, this will admit it to come as near to the meridian as I have mentioned.

The steel spindle is easily adjusted to the *zenith*; for the globe being rectified to its *latitude*, set the brass carriage at liberty, bring the quadrant and meridian together, face to face, and slide the carriage, till the lower extremity of the quadrant *buts* upon the horizon, and there screw it fast.

It is, however, to be noted, that I have found something necessary by way of *holdfast*, to prevent the brass meridian from shifting its latitude, and that without confining it in any other respect.—What I have found to answer this purpose is represented, fig. 3. The crutch-like piece of wood ABC is shewn as seen looking right down upon it. The circle DE is the horizontal section of the south pillar of the globe. The strong wire pin FG, that goes through the two arms of the crutch and pillar, serves as an axis upon which its other extremity at B is at free liberty to lift up and down, but without shake upon the pin; and the whole being split with a fine saw, from B to H, the notch BK lays hold of the under side of the brass meridian, and by tightening the finger screw LM, it firmly clips it, and retains it in any given position. And that it may be under no confinement *cross-ways*, the hole in the pillar is opened on both sides,

as shewn in the section, to give it liberty of accommodation; the pin being fast in the two ends of the crutch, and turning gently in the pillar; the whole being slender and compliant, except in point of length.

N. B. Those that would use the globe to the best advantage to solve problems, should be careful to get a *just* declination, as also a *distinct* point to mark it; and as the circles and divisions upon the surface of the globe itself, are not always sufficiently to be depended on for this purpose, I have found the following expedient fully to answer. Chuse any plain white part of the globe's surface, answerable to the declination given, and with the point of a needle or protracting pin, by the help of the divisions of the brass meridian, mark a fine point upon the blank surface of the globe, and upon this point make a dot with ink, with the small point of a pen, which rub off with the finger, and it will leave a fine black speck behind. This dot being brought to the meridian, rectify the horary index to it, and it will accurately represent the center of the celestial body whose investigation is wanted.






Fig. 2.




Fig. 1.




Fig. 3.





II. *Objections to the Experiments and Observations relating to the Principle of Acidity, the Composition of Water, and Phlogiston, considered; with farther Experiments and Observations on the same Subject. By the Rev. Joseph Priestley, LL.D. F. R. S.*

Read November 27, 1788.

HAVING never failed, when the experiments were conducted with due attention, to procure some *acid* whenever I decomposed dephlogisticated and inflammable air in close vessels, I concluded that an acid was the necessary result of the union of those two kinds of air, and not water only; which is an hypothesis that has been maintained by Mr. LAVOISIER and others, and which has been made the basis of an intirely new system of chemistry, to which a new system of terms and characters has been adapted. The *facts* that I alleged were not disputed; but to my *conclusion* it was objected, that the acid I procured might come from the phlogisticated air, which in one of my processes could not be excluded; and that it was reasonable to conclude that this was the case, because Mr. CAVENDISH had procured the same acid, *viz.* the nitrous, by decomposing dephlogisticated and phlogisticated air with the electric spark. In other cases it has been said, that the *fixed air* I procured came from the *plumbago* in the iron from which my inflammable air had been extracted.

With respect to the former of these objections I would observe, that my process is very different from that of Mr.

CAVENDISH;

CAVENDISH; his decomposition being a very slow one by electricity, and mine a very rapid one by *simple ignition*, a process by which phlogisticated air, as I found by actual trial, was not at all affected; the dephlogisticated and inflammable airs uniting, and leaving the phlogisticated air (as they probably would any other kind of air with which they might have been mixed) just as it was.

I would also observe, that there is no contradiction whatever between Mr. CAVENDISH's experiment and mine, since phlogisticated air may contain phlogiston, and by means of electricity this principle may be evolved, and unite with the dephlogisticated air (or with the acid principle contained in it) as in the process of simple ignition the same principle is evolved from inflammable air, in order to form the same union; in consequence of which, the water, which was a necessary ingredient in the composition of both the kinds of air, is precipitated. That in other circumstances than those in which I made the experiments, the acid wholly escaped, and nothing but water was found, may be easily accounted for, from the small quantity of the acid principle in proportion to the water, and the extreme volatility of it, owing, I presume, to its high phlogistication when formed in this manner.

In order to ascertain the effect of the presence of phlogisticated air in this process, I now not only repeated the experiment of mixing a given quantity of phlogisticated air with the two other kinds of air, and found, as before, that it was not affected by the operation; but I made the experiment with atmospheric air, instead of dephlogisticated. Since the air of the atmosphere contains a greater proportion of phlogisticated air, it might be expected that, if the acid I got before came from the small quantity of phlogisticated air which I could

not

not possibly exclude, I should certainly get more acid, when, instead of endeavouring to exclude it, I purposely introduced a greater quantity. But the consequence was the production of much less acid than before, the liquor I procured being sometimes not to be distinguished from pure water, except by the greatest attention possible: for though the decomposition was made in the same copper vessel which I used in the former experiments, there was now no sensible tinge of green colour in it.

When I repeated this experiment in a glass vessel, I perceived, as I imagined, the reason of the small produce of acid in these new circumstances: for the vessel was filled with a vapour which was not soon condensed, and being diffused through the phlogificated air (which is not affected by the process) is drawn away along with it, when the exhausting of the tube is repeated; whereas, when there is little or no air in the vessel besides the two kinds which unite with each other, and are decomposed, the acid vapour, having nothing to attach itself to and support it (by being entangled with it) much sooner attacks the copper, making the deep green liquor which I have described. Sometimes, however, I have procured a liquor which was sensibly green by the decomposition of atmospheric and inflammable air, but by no means of so deep a colour, or so sensibly acid, as when the dephlogificated air is used.

The extreme volatility of the acid thus formed (and which accounts for the escape of some part of it in all these processes) is apparent from this circumstance, that if the explosions be made in quick succession (the tube being exhausted immediately after each of them, and filled again as soon as possible) no liquor at all will be collected, the whole of the

acid vapour, together with the water with which it was combined, being drawn off uncondensed in every process. I once made twenty successive explosions of this kind, in a copper tube, out of which I found that I drew 37 ounce measures of air by the action of the pump, and found not a single drop of liquid, though near an hour was employed in the whole process, and the vessel was never made more than a little warmer than my hand. This was a degree of heat by no means sufficient to keep the whole of any quantity of water in a state of vapour; and is a circumstance that of itself sufficiently proves, that the vapour did not consist of water only.

Indeed, I think it impossible for any one to *see* this vapour in a tall glass vessel, and especially to observe how it falls from one end of it to the other, and the time that is required to its wholly disappearing, without being satisfied that it consists of something else than mere water, the vapour of which would be more equally diffused. If the appearance to the eye should fail to convince any person of this, the sense of *smell* would do it: for even in a glass vessel it is very offensive, though it might not be pronounced to be *acid*. I conjecture, however, that this, and every other species of *smell*, is produced by some modification of the acid or alkaline principle. Some may be disposed to ascribe this smell to the *iron* from which the inflammable air was produced; but the smell is the same, or nearly so, when the air is from tin, and would probably be the same if it were from any other substance.

Besides using atmospheric air, which contains a greater proportion of phlogisticated air, I have sometimes used dephlogisticated air which was not very pure; and in this case I have always observed, that the liquor I procured had less colour, and was less sensibly acid.

These observations might, I should think, satisfy any reasonable person, that the acid liquor which I procured by the explosion of dephlogisticated and inflammable air in close vessels did not come from the phlogisticated air which could not be excluded, whether it was that which remained in the vessel after exhausting it by the air pump, or that with which the dephlogisticated air was more or less contaminated.

But besides these experiments, in which I procured the green acid liquor by the explosion of dephlogisticated and inflammable air in close vessels, I made another, to which I thought the same objection could not have been made, because no air pump was used in it, and nothing but the purest dephlogisticated air was employed, being separated in the process from *precipitate per se* in contact with the purest inflammable air in a glass vessel which had been previously filled with mercury. Accordingly, the only objection made to *this* experiment was, that the preparation I made use of might be impure, containing something which might yield phlogisticated air. This appeared to me highly improbable, as the precipitate had been made by M. CADET, and for the purpose of philosophical experiments. Besides, if the heat of a burning lens should dislodge phlogisticated air from any unperceived impurity in this preparation, mere *heat* will not decompose this air. Let any person try the effect of a lens on such air, or any substance containing it, and produce an acid if he can.

M. BERTHOLLET, however, thinking that this might be the case, desired that I would send him a specimen of my precipitate *per se*. Accordingly, I sent him all that remained of it; and, in return, he sent me a quantity on the goodness of which I might depend. With this preparation I repeated my former experiment; and, by giving more attention to the

process, found it to be far more decisively conclusive in favour of my opinion than I had imagined. In the former experiment I had attended only to the drop of *water* which was found in the vessel in which the process was made; and finding that it turned the juice of turnsole red, I concluded, that it contained nitrous acid: but I now examined the *air* that remained in the vessel, and found that a considerable proportion of it was fixed air; so that I am now satisfied *this* was the acid with which it was impregnated, and not the *nitrous*. Still, however, some acid is the constant result of the union of the two kinds of air, and not water only. A quantity of the same precipitate *per se* yielded no fixed air by heat.

Comparing this experiment with that in which iron is ignited in dephlogisticated air, this general conclusion may be drawn, *viz.* that when either inflammable or dephlogisticated air is extracted from any substance in contact with the other kind of air, so that one of them is made to unite with the other in what may be called its *nascent state*, the result will be *fixed air*; but that if both of them be completely formed before their union, the result will be *nitrous acid*.

It has been said, that the fixed air produced in both these experiments may come from the *plumbago* in the iron from which the inflammable air is obtained. But since we ascertain the quantity of plumbago contained in iron by what remains after its solution in acids, it is in the highest degree improbable, that whatever plumbago there may be in iron, any part of it should enter into the inflammable air procured from it. Besides, according to the antiphlogistic hypothesis, all inflammable air comes from water only.

As it cannot be said, that any real fixed air is found in inflammable air from iron (since it is not discoverable by lime-water)

water) it must be supposed, that the elements, or component parts of fixed air are in it; but one of these elements is pure air, and the mixture of nitrous air shews, that it contains no such thing, though, according to M. LAVOISIER, fixed air contains 72 parts in 100 of pure air.

However, being apprized of this objection to inflammable air from iron, I made use of inflammable air from *tin*, and I had the same result as with that from iron. I also calculated the weight of the fixed air which I got in the process, and comparing it with the plumbago which the iron necessary to make the inflammable could have contained, I found, that, in all the cases, it far exceeded the weight of the plumbago; so that it was absolutely impossible, that the fixed air which I found should have had this origin. For the greater satisfaction, I shall recite the particulars of a few experiments of this kind.

In ten ounce measures of inflammable air from malleable iron I revived *red precipitate* till there remained only 1.1 oz. measure of air, and of this 0.07 oz. m. was fixed air, being completely absorbed by water. The weight of this air would be 0.063 gr. But, since 960 grains of iron will yield 1054 oz. measures of inflammable air, the iron employed in procuring all the inflammable air that was used in this experiment, *viz.* 8.9 oz. measures (without allowing for any that went to the revivification of the mercury) would be 8.1 grains; and since M. BERGMAN supposes, that 100 grains of iron contains 0.12 gr. of plumbago, the quantity of it in this iron would only be 0.01008 gr. which is not quite a sixth part of the weight of the fixed air.

With the *precipitate per se*, sent me by M. BERTHOLLET, I revived mercury till  $8\frac{1}{2}$  oz. m. of inflammable air was reduced to  $2\frac{1}{2}$  oz. m., and of this 0.04 oz. m. at least was fixed air.

This

This is not quite so much in proportion as in the preceding experiment, but abundantly more than the weight of the plumbago.

In 8 oz. m. of inflammable air I revived *minium* (which I found to have exactly the same effect in this process as red precipitate, or precipitate *per se*), till it was reduced to 1.2 oz. m.; and of this 0.028 oz. m. was fixed air, which would exceed the weight of the plumbago more than three times. In reviving lead from massicot (which I prepared by expelling the pure air from minium) I had no fixed air in the residuum.

In 7 oz. m. of inflammable air from tin by spirit of salt, I revived red precipitate till it was reduced to 1.1 oz. m.; and in this the fixed air was something more than in proportion to that in the last experiment.

In my last volume of *Experiments*, p. 30. I mentioned some instances of the revival of red precipitate in inflammable air, without finding any fixed air, though in one I perceived a slight appearance of it. To this I can only say, that I now always find it, and have, in the preceding cases, measured the quantity of it; so that, though I did not find any before, I must presume that I did not use the same precautions that I did at this time: and it is possible, that I might not attend to the effect of admitting a large quantity of water to a small quantity of fixed air, which would presently absorb the greatest part of it. I also think I recollect, that I then continued the process as far as I possibly could, and consequently left very little air in the vessel; whereas I now purposely left a good deal, that the admission of water might have less effect on the fixed air diffused through it.

This also may be said in favour of the greater accuracy of my present experiments, that they intirely remove a very great difficulty,



difficulty, which I acknowledged, p. 128. in finding different results from seemingly similar circumstances; whereas I now find that both the circumstances and the results are different. Besides, the *positive* evidence of actually finding a substance is always more conclusive than the *negative* one, of not finding it.

I do not know that any objection can be made to the inflammable air from *tin*, as this metal has not been proved to contain plumbago. I wished, however, to repeat this experiment with inflammable air from *sulphur*. But though, when steam is sent over melted sulphur, a small quantity of inflammable air is procured, as I observed in my last volume of experiments; yet, as sulphur cannot part with much phlogiston, except in proportion as it imbibes pure air, to form oil of vitriol, I could not in this manner easily procure enough for my purpose.

In order to supply the sulphur with pure air, I mixed with it a quantity of *turbith mineral*; but this made it yield vitriolic acid air, though in great abundance, there not being, I imagine, *water* enough to form inflammable air: for when iron is dissolved in concentrated acid of vitriol, vitriolic acid air is produced; but in diluted vitriolic acid, the produce is inflammable air. With a view to supply these materials with water, I sent steam over them; but it did not combine with the air, which was still only vitriolic acid air.

Since, however, vitriolic acid air unquestionably contains the same principle which forms the inflammability of inflammable air, this experiment proves, that sulphur is not that simple substance which the antiphlogistians suppose it to be; but that it contains phlogiston. Had it been nothing more than a substance which had a strong affinity to pure air, it would have

have united with the pure air from the turbith mineral, and have made vitriolic acid; but no vitriolic acid air would have been produced.

That vitriolic acid air contains the same inflammable principle with inflammable air is evident from the quantity of vitriolic acid air which I produced by reviving copper from blue vitriol in inflammable air. See my *Experiments*, vol. VI. p. 15. Mr. KIRWAN also produced this air from sulphur and red precipitate. See his *Treatise on Phlogiston*, p. 29.

When I used a small quantity of sulphur in proportion to the turbith mineral, the first produce was vitriolic acid air, and afterwards dephlogisticated air, from the turbith mineral alone, the effect of the sulphur having been exhausted.

According to the antiphlogistic theory, *phosphorus*, as well as sulphur, is a simple substance; and when it is ignited imbibes pure air, and thereby becomes the phosphoric acid, without parting with any thing. But I find, that after the accension of it in dephlogisticated air, there is a considerable quantity of fixed air in the residuum; and this fixed air could only be formed by the union of the dephlogisticated air in the vessel with the phlogiston contained in the phosphorus. Mr. KIRWAN had a similar result from phosphorus confined in atmospheric air. As it is not pretended, that there is any plumbago in phosphorus, this experiment is not liable to the objection that has been made to those in which inflammable air from iron was made use of.

It will be expected, that in this reply to the objections that have been made to my experiments establishing the doctrine of phlogiston, I should consider what has been alledged by Mess. LAVOISIER, BERTHOLLET, and DE FOURCROY, in favour of their new system, in their *Report* on the subject of the new  
chemical

chemical characters invented by Mess. HASSENFRAZ and ADET, subjoined to the new *Nomenclature Chymique*. I shall therefore notice what appears to me to be most important in that publication.

“ One of the articles of the modern doctrine” (of which they say, p. 311. “ that it cost more than twenty years labour, which “ the force of reasoning has obliged many celebrated chemists to “ adopt, and in favour of which much greater numbers are ready “ to decide;” and the evidence for which they say, p. 301. “ is the most complete chemical proof), which seems the “ most solidly established,” p. 298, “ is the formation, the “ decomposition, and recomposition of water; and how is “ it possible,” they add, “ to doubt of it, when we see that, “ in burning together 15 grains of inflammable air and 85 of “ pure air, we get exactly 100 grains of water; and when we “ can, by decomposition, find again these same two principles, “ in the same proportions?”

To this I must say, as I have done, Experiments, vol. VI. p. 139. (and when I wrote that, I was myself a believer in the decomposition of water), that I have never been able to find the full weight of the air decomposed in the water produced by the decomposition; and that now I apprehend it will not be denied, that the produce of this decomposition is not mere water, but always some acid.

As to the supposed decomposition of water by means of iron, I have shewn that it is a fallacy; since the iron imbibes nothing but water when it parts with its phlogiston. And I have observed (Experiments, vol. VI. p. 83.), that when this finery cinder is reconverted into iron by inflammable air, nothing but water is expelled from it; and that the residuum of the air is purely inflammable, without containing any fixed

air. It is evident, therefore, that the iron had imbibed pure water only. Had the iron imbibed dephlogisticated air from the water, and not water itself, there seems to be no reason why fixed air should not be found in this, as well as in the exactly similar process with minium and precipitate *per se*. Also, it can never be supposed, that the addition which iron gains, of one-third of its weight, is from air contained in steam, if it could be proved to contain any; because, if there be a sufficient quantity of iron, the whole of the water will be imbibed; so that, on this hypothesis, water must be nothing but dephlogisticated air condensed.

There is, I acknowledge, a great difficulty in explaining the experiment of iron first imbibing water, and parting with phlogiston, and again parting with its water, and imbibing phlogiston, in circumstances of heat so nearly similar as those which I have described. It seems as if the affinity of iron to water and to phlogiston was each, in their turns, stronger than the other. To this I can only say, that the whole doctrine of affinities, as far as it is true, is founded on facts; and these are clearly such as I have represented; and that a difference of circumstances, which is not apparent at present, may become so when we shall have given sufficient attention to them.

In order to satisfy myself whether any thing besides *water* was expelled from finery cinder by heat, I went through similar processes with this substance and *massicot*, from which all air had been previously expelled; and after reviving both of them in inflammable air, I found the results, in all respects, the very same. The residuums of the inflammable air were equally free from fixed air; and when they were fired with equal quantities of dephlogisticated air, the diminutions of  
bulk

bulk were very nearly the same, less than when the original inflammable air was used, because all the impurities in the whole quantity were retained in a small residuum, the metals having imbibed nothing but pure phlogiston. Also the inflammable air had been long confined by water, in consequence of which it is always altered more or less. The particulars of the processes were as follows :

The finery cinder was revived in 7 oz. m. of inflammable air, which was thereby reduced to  $1\frac{1}{4}$  oz. m.; and an oz. m. of this residuum being fired together with an equal quantity of dephlogisticated air, not very pure, the diminution of both was to 28 divisions of a tube, of which 30 was one oz. m. when with equal quantities of the same dephlogisticated and the original inflammable air the diminution was to 18.

The massicot was reduced in 8 oz. m. of inflammable air till it was reduced to  $1\frac{1}{4}$  oz. m.; and after the process with the dephlogisticated air, the diminution was to 29, when with the original inflammable air it was to  $17\frac{1}{2}$ .

In both the residuums, after the explosion, there was a slight appearance of *fixed air*, though none could be perceived before the explosion; but in both cases it was so slight that it could not have been perceived by the diminution of its bulk. But since both fixed air and nitrous acid are produced from the same materials in different circumstances, it cannot be thought extraordinary if, in some cases, both should be produced at the same time.

M. LAVOISIER and his associates farther observe, p. 300. with respect to my experiments, that “ when a calx is revived  
“ in inflammable air, more water is found in the vessel than the  
“ weight of inflammable air that disappears, so that it could  
“ not have been contained in that air.” They only refer to

my experiments in general; but as they speak of the water produced as appearing both on the inside of the vessel, and on the surface of the mercury, it can be no other than the experiment of the revival of iron from fiery cinder; and the water that is found in this process was never supposed to come from the little that is contained in the inflammable air, but the much greater quantity contained in the cinder.

Before I conclude this Paper, I shall just mention a few circumstances attending the many explosions I have made of inflammable and dephlogisticated air in the long metallic and glass vessels I have made use of, as they were pretty remarkable. The explosions were made by a small electric spark at one end of the vessel, and the greatest force of the explosion was always at the other end. No tinned iron vessel could bear many of them before they swelled out at that end, and at length burst; and even the flat end of the copper vessel, which was not less than one-tenth of an inch thick, was in time made quite convex, and the cylindrical part next to it was made very sensibly wider than any other part of the tube. This must have been effected by mere *force*, and not by *heat*; for the hottest part of the tube, after every explosion, was never there, but always about the middle, though something nearer to that end than the other, and in the glass vessel the dense cloud was always formed at that end.

The probability is, that the air where the electric spark is made taking fire first, the inflammation does not extend itself so rapidly but that the air at the opposite end is first condensed, in consequence of the inflammation and expansion of the air at the other end, so that the air is there fired in a condensed state; and hence its greater force.

III. *Observations on the Class of Animals called, by Linnæus, Amphibia; particularly on the Means of distinguishing those Serpents which are venomous, from those which are not so.*  
By Edward Whitaker Gray, M. D. F. R. S.

Read December 18, 1788.

OF the various classes of the animal kingdom, no one has been so little attended to as the class, called by LINNÆUS, Amphibia. What he himself did in that class (though far superior to what any other person has done) was evidently done in a hurry; false references are, at least, as common in that, as in any other part of his works, and many of his descriptions are given in a very careless manner; others there are, however, which are truly worthy of their author, and in which the specific characters are pointed out with that clearness and precision, which so eminently distinguish the descriptions of LINNÆUS from those of all his predecessors.

In the construction of the class, LINNÆUS has been particularly unfortunate; as he has erred, not only in making an unilocular heart one of the characters of it, but also in making the cartilaginous fishes a part of it. I think it needless to mention the causes which led him to this latter error; every anatomist now agrees that the Amphibia Nantes are not furnished with lungs; and every naturalist is convinced of the propriety of removing them, from the class of Amphibia, to that of Fishes. I shall only observe that, by the removal, the name of the class

class (which some naturalists have cavilled at) becomes much less objectionable; there being few genera, in the two orders of which it is now presumed to consist, which do not contain animals to which the term amphibious may, with some propriety, be given; whereas, in the order of Nantes, not one species occurs which has the smallest claim to that title. With respect to the other error I noticed (*viz.* that of supposing the hearts of the Amphibia to be single) it would be easy to shew that it was not an uncommon one, at the time LINNÆUS formed his system. And indeed he appears to have been led into it, by following an author whom he probably supposed of too great fame not to be safely relied on. At least, in defence of his opinion, he quotes the following words of BOERHAAVE. “*In omnibus animalibus in quibus sanguis non calet, ventriculus cordis est unicus.*” Whether the hearts of all the different genera, of which the class is composed, have yet been accurately examined; and whether an exact similarity of structure is found throughout the class; are questions I do not mean, at present, to examine. It is sufficient for my purpose to observe, that the hearts of most of the Amphibia are now well known to be double, with an immediate communication between the two cavities; which structure seems peculiarly adapted to that change of element, which (as I before observed) many of them can, for a time, support; and thereby furnishes another argument in favour of the name LINNÆUS has given to the class.

To consider the structure of the heart, however, is not absolutely necessary in forming the characters of the class: the animals of which it consists being sufficiently distinguished from all others, by having cold red blood, and breathing by means of lungs. These two characters render the class perfectly distinct from the rest; the two superior ones, *viz.*



Mammalia and Birds, having warm blood; and the three inferior ones, *viz.* Fishes, Insects, and Worms, not being furnished with lungs.

In his generic characters, LINNÆUS has been more successful than in those of the class; insomuch that they may, I think, be considered as the best hitherto given. Whoever will be at the pains of comparing LINNÆUS's genera of Amphibia with those of GRONOVIVS, will find, that the generic characters of the former, though few in number, are precise and distinct; while those of the latter, though more numerous, are vague, indistinct, and sometimes inaccurate. As a glaring instance of inaccuracy, I need only refer to the Chamæleon, which by GRONOVIVS is made a distinct genus, of which one of his characters is, *Pedes unguibus destituti*; whereas, in fact, the feet of that animal are furnished with very distinct, and pretty large, claws.

But though LINNÆUS's genera of Amphibia are, upon the whole, well formed, it must be allowed to be a great imperfection in them, that the venomous serpents are not separated from the others.

From some expressions of his, in the Preface to the *Museum Regis*, and in the Introduction to the Class Amphibia, in the *Systema Naturæ*, it seems, that he thought it not easy to distinguish them, by any external characters; and his ideas respecting the venomous fangs themselves were (as we shall see hereafter) so vague and confused, that it was hardly possible for him to attempt to found a generic distinction upon them\*.

\* As a sort of comparative excuse for LINNÆUS, it may be observed, that GRONOVIVS (though he made two more genera of Serpents than LINNÆUS) did not separate the venomous ones; neither has he distinguished them by a mark (as LINNÆUS has) or by any other means.

Whether

Whether venomous Serpents can be, with certainty, distinguished from others, and if so, how they are to be known, is what I mean to consider in this Paper; in doing which I shall examine, first, how far they may be distinguished by any external characters; secondly, supposing the venomous fangs to be the only certain criterion, how those fangs are to be distinguished from common teeth.

Though Serpents, by their internal organization, naturally belong to the third class of the animal kingdom, they are, in their external form, more simple than most of the animals belonging to the three inferior classes; their external characters must consequently be very few. I shall first examine those of the head; and, as all venomous Serpents (so far as our present experience extends) are contained in the three first of LINNÆUS's genera, I shall, at present, consider only those three.

In the first genus, *Crotalus*, the head is broader than the neck, depressed or flat at top, and covered with small scales. These three characters are particularly observable in the three intermediate species *horridus*, *Dryinas*, and *Durissus*. In the *miliarius* the scales of the head are rather larger than in the others. The *mutus* I have never seen; but it certainly should not be placed among the *Crotali* \*.

As all the species of this genus are venomous, one is naturally led, by the examination of it, to consider the fore-mentioned characters as being, in some measure, proper to venomous serpents. In order to see how far they are so, I shall, for the present, pass over the next genus, *Boa*, and consider

\* LINNÆUS's reason for not placing it among the *Boæ* seems to have been, that he supposed none of them were venomous. He appears, however, to have had his doubts about the *contortrix*. I have examined it, and am convinced it is venomous.

that which follows it, Coluber. In that genus are many venomous species, and it is very certain that, in general, they have the fore-mentioned characters; examples of which may be seen in the Atropos, Ceraftes \*, atrox, Berus, and others. It is, however, equally certain, that there are some in which they are not to be found. As an example of this, I need only mention the Naja, a species well known to be very venomous; the head of which is neither depressed nor broad, is covered with large scales, and is, in every respect, a complete exception to what has been said, respecting the heads of venomous Serpents.

Since then, there are venomous Serpents in which the fore-mentioned characters, *viz.* a broad and depressed head, covered with small scales, are not to be found; I shall next examine whether those characters are to be found in any of those Serpents which are not venomous. In the genus Coluber there are very few (except venomous ones) which have the head much broader than the neck; and of those few, I believe, every one has the head covered with large scales. But in the genus Boa, though no species is venomous, except the contortrix, almost every one has the head broad, depressed, and covered with small scales. The canina, Constrictor, hortulana, besides some others not described by LINNÆUS, furnish

\* The Ceraftes is not marked by LINNÆUS as a venomous species. He probably depended upon HASSELQUIST's description, which I suspect to have been made from a mutilated specimen. Mr. ELLIS's description in the Philosophical Transactions, Vol. LVI. p. 287. is only a Translation of HASSELQUIST's. But he observes, that Dr. TURNBULL told him it was venomous. That it is so, I have not the smallest doubt, though in the only specimen I have seen of it the fangs were wanting. IMPERATO, who has given a figure of it (Hist. Nat. p. 784. Ed. Nap.), says it is very venomous.

examples of this. It must, however, be confessed, that the general character of the head of the Boa, though differing very widely from that of those Colubri which are not venomous, is not quite that of the Crotalus; but the difference, though very obvious to a person accustomed to the examination of Serpents, is perhaps not easy to be fully expressed in words. It seems, however, to consist principally in a lateral compression, and elongation, of the anterior part of the head, so as to form a kind of snout. Hence the trivial name of *canina* is given by LINNÆUS to one of the species.

From the characters of the head (as the trunk affords none deserving consideration) I shall proceed to those of the other extremity.

In the Crotali I have never found the tail (exclusive of the Rattle) to exceed one-ninth part of the whole length; sometimes I have found it much shorter. In some of the venomous Colubri, the proportion is still less. In the Atropos I found it only one-thirteenth. In the English Viper (*Coluber Berus*) it is commonly about one-seventh or eighth. In some venomous species, however, the proportion is something greater. In the *Naja* I have found it as much as one-sixth; which proportion is, I believe, as great as I have ever observed: but that I may be sure to keep within the truth, I will only say, that I have never met with a venomous Serpent, the tail of which was equal to one-fifth of the whole length\*.

\* The tail of the *Boa contortrix* is said by LINNÆUS to be one-third; but his own enumeration of the Scuta sufficiently shews that this must be an error. The *Coluber Leberis*, *Dipsas*, and *mycterizans* appear, by the number of scales under their tail, to furnish exceptions to what I have said. The two first I have never seen, but suspect they are not venomous; that the last is not so I am very certain, having examined many specimens of it.

With respect to those Colubri which are not venomous, it must be confessed, that there are many whose tails are within the limits assigned to the venomous ones. In the Coluber *Æsculapii*, *doliatus*, *gétulus*, and some others, the tail is not, in general, more than one-seventh of the whole length. In the *lemniscatus* I have found it not exceeding one-twelfth or thirteenth; but I know no other Linnæan species in which it is so short. In the greater number, however, the proportion of tail is more considerable; in many, it is full one-third. In the *Ahætulla*, and in some species not described by LINNÆUS, I have seen it more than two-fifths; but have never met with a species in which it was quite so long as the trunk, or half of the whole length.

I have not considered the *Boæ*, because none of the Linnæan species, of that genus, have their tails either remarkably long, or short; but, in two species, not described by LINNÆUS, I found the tail very little exceeding the proportion I have assigned to the Coluber *lemniscatus*.

In the thickness of the tail, or in the acuteness of its termination, I have observed no difference worth remarking. In every species of the three first genera, the tail is thinner than the trunk; and in most of them it is more or less acute. The few exceptions I have observed were, I believe, none of them venomous; but they are too few to deserve any particular consideration.

A character of great use in distinguishing the species of Serpents, and which was not overlooked by LINNÆUS, is, that elevated line, or carina, with which the scales of many species are furnished. In order to shew how far this is to be considered as serving to distinguish venomous Serpents from others, I need only observe, that I have examined one hundred and twelve species of Serpents, not venomous, belonging to the three first

genera; and find that eighty of them have smooth scales, and thirty-two only have carinated ones. Of venomous Serpents I have examined twenty-six; of which number, twenty have carinated scales, and only six have smooth ones. Upon the whole, therefore, carinated scales must be considered as being, in some measure, a character of venomous Serpents.

In what I have hitherto said, I have considered only the three first genera of Serpents; I shall now make some remarks upon the three last.

These three (*viz.* Anguis, Amphibæna, and Cæcilia), besides the characters assigned them by LINNÆUS, have some others which are common to all, and which render them very different, in their external appearance, from any of the three first genera. These are, a very thick and obtuse tail, and a head which is very indistinct\*, and furnished with very small eyes. This last character (*viz.* very small eyes) is sometimes, though very rarely, met with among the Colubri, for instance, in the lemniscatus; in the three last genera, however, it takes place, I believe, without exception. The thickness of the tail is also common to every species; and though in the Anguis bipes, and in another species, not described by LINNÆUS, but figured in BROWNE'S History of Jamaica (Tab. XLIV. fig. 1. †), the tail has an acute termination, yet in both those species, especially in the last, it continues thick to the end, and becomes suddenly sharp, being what in botanical language would be called, *obtusa cum acumine*. With respect to the proportionate length of tail, however, it is very remarkable, that the genus

\* This indistinctness of the head, which is more or less common to each genus, is in the Amphibæna so considerable, as to have given rise to the supposition of that Serpent's having a head at each end.

† This figure is, by LINNÆUS, erroneously quoted as his Anguis lumbicalis.

Anguis affords examples of much less proportion, and also of much greater, than is to be found in any of the three first genera. In the Anguis Scytale the tail is not above one-twentieth of the whole length; in the maculata it is not above one-fortieth; yet in the Anguis fragilis, and in the ventralis, the tail is always longer than the trunk, or, in other words, is more than half the whole length. Indeed, in one specimen of the last mentioned species, I found the tail nearly two-thirds of the whole length. It may, however, be questioned whether that species is really an Anguis, or a Lacerta\*.

I shall make no further remarks on the external characters of Serpents; the principal inferences to be deduced from those I have already made, are the following.

1st, That a broad head, covered with small scales, though it be not a certain criterion of venomous Serpents, is, with some few exceptions, a general character of them.

2dly, That a tail under one-fifth of the whole length, is also a general character of venomous Serpents; but, since many of those which are not venomous have tails as short, little dependence can be placed upon that circumstance alone. On the other hand, a tail exceeding that proportion, is a pretty certain mark that the species, to which it belongs, is not venomous.

3dly, That a thin and acute tail is by no means to be considered as peculiar to venomous Serpents; though a thick and obtuse one is only to be found among those which are not venomous.

\* The Anguis ventralis of LINNÆUS, is so very like the Lacerta apoda, described by PALLAS, in Vol. XIX. of the Novi Comment. Petrop. as to render it doubtful whether it may not be the same. When I first examined it, I considered it as a Lacerta, on account of the projecting future along the body, and the open ears; but I have since met with a specimen, which had two large echinated *Penes* (as they are called) a character which is, I believe, peculiar to Serpents.

4thly, That carinated scales are, in some measure, characteristic of venomous Serpents, since in them they are more common than smooth ones, in the proportion of nearly 4 to 1; whereas, smooth scales are, in those Serpents which are not venomous, more common, in the proportion of nearly 3 to 1.

Upon the whole therefore it appears, that though a pretty certain conjecture may, in many instances, be made, from the external characters; yet, in order to determine, with certainty, whether a Serpent be venomous or not, it becomes necessary to have recourse to some more certain diagnostic. This can only be sought for in the mouth; I shall therefore next consider, how the fangs, with which the mouths of venomous Serpents are furnished, are to be distinguished from common teeth.

To those who form their ideas of the fangs of venomous Serpents, from those of the Rattle-snake, or even from those of the English Viper, it will appear strange, that there should be any difficulty in distinguishing those weapons from common teeth; and indeed the distinction would really be very easy, were all venomous Serpents furnished with fangs as large as those of the fore-mentioned species. But the fact is, that in many species the fangs are full as small as common teeth, and consequently cannot, by their size, be known from them; this is the case with the *Coluber laticaudatus* \*, *lacteus*, and several others. I cannot, however, better demonstrate that the distinction, between the venomous fangs and common teeth, is not very obvious, than by shewing how very vague and erroneous

\* This species is by LINNÆUS reckoned venomous, in the *Museum Regis*, though the mark is not affixed to it in the *Systema Naturæ*. To me it appears to be certainly venomous, and is the only water Serpent I have met with that is so.



LINNÆUS's ideas about them were; nor can I better prove the want of information on this subject, than by observing that, erroneous as the ideas of LINNÆUS were, no one, that I know of, has yet attempted to furnish more correct ones.

LINNÆUS thought the fangs might be distinguished by their mobility; this, at least, may be fairly inferred, from his never mentioning them in the *Museum Regis*, without adding the epithet *mobilia*, except in one instance (the *Coluber aulicus*); and, in that very instance, the want of mobility in the supposed fangs appears evidently to raise doubts in his mind, whether they are really fangs or not. His words are, “*Dentes, sive tela, duo, rigida, parva, non mobilia.*” These doubts, respecting the above-mentioned species, I am not able to remove, as I am not sure that I have ever seen it\*. But with regard to mobility, considered in general as a character of venomous fangs, I must assert, not only that I have never found it so, but also, that I have never been able to discover in them any thing which I thought could properly be called mobility. I have, indeed, sometimes found some of them loose in their sockets; but then I have found others, in the same specimen, quite fixed. The same thing was observed both by Dr. NICHOLLS †, and by the Abbé FONTANA ‡, in the common Viper, even during life. The loose fangs may be such as have not yet been firmly fixed in their socket, or they may have been loosened by some accident: for I suspect that the fangs may be at any time loosened, and even displaced, by a small degree of violence; and that, perhaps, may be one

\* I have seen one, which agreed pretty well with LINNÆUS's description; if that was really his species, it is not venomous.

† Appendix to Dr. MEAD's Account of the Viper.

‡ FONTANA, *Traité sur le Venin de la Vipere*, chap. 1st and 2d.

reason why there is always a certain number of small fangs, near the base of the full grown ones, ready to enlarge and take their place, if they should be, by any accident, torn out.

LINNÆUS seems also to have thought that the fangs might be known by their situation. In the Introduction to the class Amphibia in the *Systema Naturæ*, he says they are, “*Dentibus simillima sed extra maxillam superiorem collocata;*” and in the description of the *Crotalus Dryinas*, in the *Amœnitates Academicæ*, he says, “*Dentes ejus duo canini uti in reliquis venenatis Serpentibus non in maxillis hærent, iis enim vulnerando, non autem ictus infligendo utitur.*”

These two quotations shew, that LINNÆUS thought the situation of the fangs different from that of the common teeth; the last also shews that he thought their mode of action influenced by it. What difference in situation may be found by accurate dissection, it is foreign from my present purpose to enquire; I am, however, very certain that common examination \* will not discover any difference, in that respect, between the fangs of venomous Serpents, and the teeth of others.

But the most singular opinion of LINNÆUS, respecting the venomous fangs, was, that they were sometimes fixed in the base of the jaw. Of this he has given two instances in the *Museum Regis*. One in the description of the *Coluber severus*, of which he says, “*Hastæ mobiles solitarie versus basin maxillarum interius adhærent.*” The other in that of the *Coluber stolatus*. His words there are, “*Tela mobilia ad basin maxil-*

\* By common examination I mean such as may be made without dissecting, or otherwise damaging, the specimen to be examined; and such only do I suppose allowable in the distinction I am seeking to establish.

“*larum affixa, ut vix vulnerare valeat hostes, solum cibos veneno*  
“*inficere.*”

LINNÆUS's opinion respecting the use of the fangs, in the last mentioned species, appears to me not very clearly expressed \*. But I have quoted both descriptions, merely to shew that LINNÆUS thought the fangs were sometimes placed in the base of the jaw; an idea for which I have never been able to discover any foundation. The first of the two species in question I have never seen; of the *stolatus* I have examined several specimens, and am convinced it is not venomous.

I shall not dwell any longer on the false notions which have been entertained, respecting the fangs of venomous Serpents, but shall proceed to shew how, in my opinion, they may be most easily, and most certainly, distinguished from common teeth.

With respect to their size, I have already observed that it is very various, consequently no certain judgement can, in all cases, be made from that circumstance. In some species they are so large, that their size alone sufficiently distinguishes them from common teeth; but in others they are so small, that it is very difficult to discover them.

The size of the common teeth also varies very much, in different species. In the *Coluber mycterizans* they are remarkably large, especially those which are situated near the apex of the upper jaw; which circumstance probably helped to lead LINNÆUS into the erroneous opinion he entertained, that this Ser-

\* LINNÆUS's opinion seems not unlike that of the Abbé FONTANA, who (in the work already mentioned, chap. 12.) supposes the poison of the Viper may be of use, to the animal, in digestion. To me the venomous fangs have ever appeared to be merely offensive weapons; nor can I see greater difficulty in supposing such a weapon, with the power of injecting poison, placed in the head of a Viper or Rattle-snake, than in supposing such an one, with a similar power, placed in the tail of a Wasp or Hornet.

pent was venomous. But in many species the teeth are so small, that it is impossible to discover, merely by looking into the mouth, that the animal has any. Yet in that case they may be very easily detected, by drawing a pin (or any other hard substance) with a moderate degree of pressure, along the edge of the jaw, from the apex to the angle of the mouth, when they will be felt to grate against the pin, like the teeth of a saw.

Although the size of the venomous fangs is very various, their situation is, I believe, always the same; namely, in the anterior and exterior part of the upper jaw, which situation I consider as the only one in which venomous fangs are ever found. But as, in those Serpents which are not venomous, common teeth are found in that part of the jaw, it is plain that we cannot, by situation alone, distinguish one from the other. They may, however, be distinguished with great ease, and I believe also with great certainty, by the following simple operation. When it is discovered that there is something like teeth in the fore-mentioned part of the upper jaw, let a pin be drawn, in the manner already described, from that part of the jaw to the angle of the mouth (which operation may, for greater certainty, be tried on each side). If no more teeth are felt in that line, it may I believe be certainly concluded, that those first discovered are what I have distinguished by the name of fangs, and consequently, that the Serpent is a venomous one \*. If, on the contrary, the teeth first discovered are found not to stand alone, but to be only a part of a complete row, it may as certainly be concluded, that the Serpent is not venomous.

\* If a specimen should be met with, in which no teeth, of any kind, can be discovered in the margin of the upper jaw, the presumption is, that it is a venomous Serpent, which has lost its fangs; but I have never met with such an one, except the *Coluber Ceraastes* already mentioned.

In the upper jaw, both of venomous Serpents and others, besides the teeth already spoken of, there are two interior rows; consequently, the distinction I have endeavoured to establish might be expressed in other words, by saying, that all venomous Serpents have only two rows of teeth, in the upper jaw, and all others have four\*. I think it better, however, to leave the interior rows out of the question, as, in many species, the teeth of which they are composed are so small, as to make it very difficult to discover them. Indeed, in two species of Anguis, I can hardly be sure that I have discovered them; but as, in every other species, I have never failed to do so, I presume I may, with very little risk of error, assert, that all Serpents whatever are furnished with them; and that those only, which are not venomous, have the exterior rows.

What I have said sufficiently shews that LINNÆUS's ideas, respecting venomous serpents, were such as did not permit him to separate them from the others; if the method I have proposed shall be found to render the distinction of them sufficiently clear and easy, it naturally follows, that they should be made generically distinct. Some other reforms might also be made in LINNÆUS's class of Amphibia, the consideration of which I do not mean, at present, to enter further into. But, before I conclude, I think it necessary to notice an inaccuracy of LINNÆUS, of a different kind from those I have already pointed out.

\* GRONOVIVS, of whose inaccuracy I have already given one instance, in describing the *Crotalus Durissus*, in his *Museum Ichthyologicum*, says it has no teeth, except the venomous fangs. KLEIN, in his *Tentamen Herpetologiæ*, has gone still further, having actually made a genus of Serpents without teeth, which he calls *Anodon*. He appears not to have examined the mouth of a single species; but to have depended intirely upon the descriptions of SEBA.

In the Preface to the *Museum Regis*, and in the Introduction to the class Amphibia, in the *Systema Naturæ*, LINNÆUS says, that the proportion of venomous Serpents to others, is 1 in 10; yet in the *Systema Naturæ*, in which the sum total of species is one hundred and thirty-one, he has marked twenty-three as venomous, which is somewhat more than 1 in 6. How he came to be so much at variance with himself, I know not; but the last mentioned proportion seems to me to be not far from the truth; as I find that I have examined one hundred and fifty-four species of Serpents, of which number twenty-six appear to be venomous.

I have already mentioned, that the *Coluber stolatus* and the *mycterizans*, though marked by LINNÆUS as venomous Serpents, certainly are not so; and that I suspect the same may be said of the *Leberis*, and *Dipsas*. I have also observed, that the *Boa contortrix*, *Coluber Cerastes*, and *laticaudatus*, none of which are marked in the *Systema Naturæ*, are all of them venomous; to these last may be added the *Coluber fulvus*.

If LINNÆUS's species were all accurately examined, I have no doubt but more errors, of both kinds, would be found; for it must be observed, that though I have examined a greater number of species than LINNÆUS, not above half that number are of those described by him; consequently there remains more than one-third of his species which I have never seen. The number I have examined, however, seems to me sufficiently great to warrant the inferences I have drawn from that examination. That some exceptions to them might be found, by the examination of a greater number, is very possible; but, if these observations shall tend to rectify the false notions which have been entertained respecting venomous Serpents, and to render the distinction between them and others more clear, I trust they will be thought not totally useless.

IV. *Observations on the Dryness of the Year 1788. In a Letter from the Rev. Mr. B. Hutchinson to Sir Joseph Banks, Bart. P. R. S.*

Read January 15, 1789.

S I R,

Kimbolton, January 8, 1789.

**A**S the defect of rain has been very considerable in 1788; and in consequence a great want of water on the close of the year universally felt; perhaps the quantity fallen here, compared with that of the seven preceding years, may not be unacceptable to yourself and the Royal Society.

	Inches.	
Rain 1781	21,6	}
1782	32,3	
1783	23,6	
1784	28,0	
1785	21,0	
1786	24,7	
1787	23,8	
	<hr style="width: 50%; margin: 0 auto;"/>	
1788	14,5	

$\frac{175}{5} = 25$  inches, the mean of seven years.

By estimation it therefore appears, that the average quantity of rain of the seven preceding years is 25 inches, and the rain which fell last year is only 14,5, that is, not much more than half that quantity, if we deduct 1,3 now lying in snow, which fell in December, and not in solution. On the supposition

sition which, I believe, is not far from truth, that the whole island has had the same defect; a greater failure of the produce of the earth might have been expected than what the country has experienced; for, except in hay, and a little failure in turneps, the crops have in general been as plentiful as in most of the former years, and in fruits of the orchard much more so.

It has always been said of England, that drought never occasions want; this year verifies the assertion. But to account for crops that, taken on the whole, are rather abundant, we may consult the following monthly state of rain for 1788.

	Inches.
In January	0,3
February	1,7
March	0,7
April	0,0
May	0,6
June	1,8
July	0,8
August	3,4
September	3,4
October	0,3
November	0,2
December	1,3
	<hr/>
	14,5

Having premised, Sir, that there were no extremes of cold and heat throughout the year; the thermometer in a northern exposure never falling below the freezing point during the day-time, except on the 14th and 15th of January, the 6th, 7th,



7th, 8th, 10th, 11th, 12th, 13th, and 17th of March, and on none of those days at noon, so that there never were twenty-four hours together successive frost; therefore vegetation was never entirely at a stand. In summer it did not rise to 80 degrees, except on

		Deg.	
May	26	80	
	27	81	
	28	81	
June	18	83	with thunder and rain: then cool for a
	27	80	week.
July	11	80	
	12	82	
August	4	81	: the rest of the time exceedingly tem-
			perate.

Now, the rain that fell on February was towards the end of the month; which, together with that which fell in March, brought up the spring corn, gave an early first crop of hay to the large towns, and covered the meadows and pastures in the country; that they were not so entirely dried up through the defect of April, as to prevent the rain, which fell plentifully on the 29th of May, succeeded by more in June, giving a second crop to the former situations, and a first, though late one, to the latter: and as fructification chiefly depends on rain falling at the latter end of the season of flowering, this rain set the blossoms of wheat, and of the useful fruit-trees; as the great rains in August swelled the kernel, filled, as they term it, the bushel, and gave an opportunity for a second crop of turneps that proved more vigorous than the first.



V. *On the Method of determining, from the real Probabilities of Life, the Value of a contingent Reversion in which Three Lives are involved in the Survivorship. By Mr. William Morgan; communicated by the Rev. Richard Price, D. D. F. R. S.*

Read January 29, 1789.

**I**N a Paper which I had lately the honour of communicating to the Royal Society, respecting the method of determining the values of reversions depending on survivorships between two persons from the real probabilities of life, I observed, that the investigation of those cases in which three lives were involved in the survivorship (though attended with much more difficulty) might, however, be effected in a similar manner. The further pursuit of this subject has now convinced me that, as it is never safe, so likewise it can never be necessary to have recourse to the *expectations* of life in any case; and that the solution even of those problems which include three lives is far from being so formidable as at first sight it appears to be. I am sensible of the impropriety of entering minutely in this place into the vast variety of propositions which refer to the different orders of survivorship between three lives; but as the following problem seems to be of considerable importance on account of its being applied to the solution of many other problems, the demonstration of it, perhaps, may not be thought an improper addition to my former Paper.

PROBLEM.

## P R O B L E M.

Supposing the ages of A, B, and C, to be given; to determine, from any table of observations, the value of the sum S payable on the contingency of C's surviving B, provided the life of A shall be then extinct.

## S O L U T I O N.

Let  $a$  represent the number of persons living in the table at the age of A. Let  $a'$ ,  $a''$ ,  $a'''$ ,  $a''''$ , &c. represent the decrements of life at the end of the 1st, 2d, 3d, 4th, &c. years from the age of A. Let  $b$  represent the number of persons living at the age of B, and  $m$ ,  $n$ ,  $o$ ,  $p$ , &c. the number of persons living at the end of the 1st, 2d, 3d, 4th, &c. years from the age of B. In like manner let  $c$  represent the number of persons living at the age of C, and  $d$ ,  $e$ ,  $f$ ,  $g$ , &c. the number of persons living at the end of the 1st, 2d, 3d, 4th, &c. years from the age of C. Let  $r$  also denote the value of  $\text{£. 1}$  increased by its interest for a year. In order to receive the sum S in the first year, it is necessary either that all the three lives shall have died in that year, A having died first, B next, and C last; or that only the two lives A and B shall have died (A having died first), and that C shall have lived to the end of that year. The probability that the three lives shall die in the first year is

$\frac{a' \cdot \overline{b-m} \cdot \overline{c-d}}{abc}$ . The probability that they shall die in the order

above mentioned is  $\frac{a' \cdot \overline{b-m} \cdot \overline{c-d}}{6 \cdot abc}$ . The probability that both

A and B shall die in the first year is  $\frac{a' \cdot \overline{b-m}}{ab}$ . Half this frac-

tion, or  $\frac{a' \cdot \overline{b-m}}{2ab}$ , is the probability that the death of A shall happen before the death of B in this year. The probability that C shall survive A and B, restrained to the contingency of A's having died first, is  $\frac{a' \cdot \overline{b-m} \cdot d}{2abc}$ . The value therefore of the

sum S for the first year is  $S \times \frac{a' \cdot \overline{b-m} \cdot \overline{c-d}}{6 \cdot abc} + \frac{a' \cdot \overline{b-m} \cdot d}{2abc} = \frac{S}{abc} \times$

$\frac{a'bc}{6} - \frac{a'mc}{6} + \frac{a'ab}{3} - \frac{a'md}{3} \dots \dots \dots$  In the second year the pay-

ment of the given sum will depend on either of four events happening. First, on the contingency of all the three lives dying in that year, A having died first, B next, and C last. 2dly, On the contingency of B's dying in that year, C's living to the end of it, and A's dying in the first year. 3dly, On the contingency of B's dying after A in the second year (both of them having survived the first year) and of C's living to the end of that year. 4thly, On the contingency of A's dying in the first year, and of B and C's both dying in the second year, B having died first. The probability of the first contingency is expressed by the fraction  $\frac{a'' \cdot \overline{m-n} \cdot \overline{d-e}}{6 \cdot abc}$ . The probability of the second by the fraction  $\frac{a' \cdot \overline{m-n} \cdot e}{abc}$ . The probability of the third by the fraction  $\frac{a'' \cdot \overline{m-n} \cdot e}{2 \cdot abc}$ . And the probability of the fourth contingency by the fraction  $\frac{a' \cdot \overline{m-n} \cdot \overline{d-e}}{2abc}$ . These several fractions, therefore, multiplied into  $\frac{S}{r^2}$  will be the value of the given sum for the second year, and may be easily found =

$\frac{S}{abc r^2} \times \frac{a'' dm}{6} - \frac{a'' dn}{6} + \frac{a'' em}{3} - \frac{a'' en}{3} + \frac{a' em}{2} - \frac{a' en}{2} + \frac{a' dm}{2} - \frac{a' dn}{2}$ . In like

manner

manner the payment of the given sum in the third year will depend on the contingency of the same number of events as in the second year; that is, it will, first, depend on the contingency of all the three lives dying in that year, A having died first, B next, and C last; 2dly, on the contingency of B's dying in that particular year, C's living to the end of it, and A's dying in the first or second years; 3dly, on the contingency of B's dying after A in the third year (both of them having survived the two preceding years), and of C's living to the end of that year; and, 4thly, on the contingency of A's dying in the first or second year, and of B and C's both dying in the third year, C having died last. These several contingencies are expressed by the respective fractions

$$\frac{a''' \cdot \overline{n-o} \cdot \overline{e-f}}{6 \cdot abc} \dots \frac{a' + a'' \cdot \overline{n-o} \cdot f}{abc} \dots \frac{a' \cdot \overline{n-o} \cdot f}{2abc} \dots \text{and}$$

$$\frac{a' + a'' \cdot \overline{n-o} \cdot \overline{e-f}}{2abc}.$$

Consequently the value of the sum S for

the third year will be =  $\frac{S}{abc r^3} \times \frac{a''' \cdot en}{6} - \frac{a''' \cdot eo}{6} + \frac{a''' \cdot fn}{3} - \frac{a''' \cdot fo}{3} +$

$$\frac{a' + a'' \cdot fn}{2} - \frac{a' + a'' \cdot fo}{2} + \frac{a' + a'' \cdot en}{2} - \frac{a' + a'' \cdot eo}{2}.$$

And by reasoning in the same manner the value of the sum S for the fourth

year may be found =  $\frac{S}{abc r^4} + \frac{a'''' \cdot fo}{6} - \frac{a'''' \cdot fp}{6} + \frac{a'''' \cdot go}{3} - \frac{a'''' \cdot gp}{3} +$

$$\frac{a' + a'' + a''' \cdot go}{2} - \frac{a' + a'' + a''' \cdot gp}{2} + \frac{a' + a'' + a''' \cdot of}{2} - \frac{a' + a'' + a''' \cdot fp}{2}.$$

If either B or C be the oldest of the three lives, these series continued to the extremity of that life will express the whole value of the reversion, which will be =  $\frac{S}{6} \times$

$$\frac{a'bc}{abc r} + \frac{a'dm}{abc r^2} + \frac{a'''en}{abc r^3} + \frac{a''''fo}{abc r^4} + \&c. + \frac{S}{2r} \times \frac{a'dm}{abc r} + \frac{a' + a'' \cdot en}{abc r^2} +$$

$$\frac{a' + a'' + a''' \cdot fo}{abcr^3} + \&c. - \frac{S}{6} \times \frac{a'mc}{abcr} + \frac{a''na}{abcr^2} + \frac{a'''oe}{abcr^3} + \frac{a'''' \cdot pf}{abcr^4} + \&c. -$$

$$\frac{S}{2r} \times \frac{a'dn}{abcr} + \frac{a' + a'' \cdot eo}{abcr^2} + \frac{a' + a'' + a''' \cdot fp}{abcr^3} + \&c. + \frac{S}{3} \times \frac{a'db}{abcr} + \frac{a'' \cdot en}{abcr^2} +$$

$$\frac{a''' \cdot fn}{abcr^3} + \frac{a'''' \cdot go}{abcr^4} + \&c. + \frac{S}{2r} \times \frac{a'em}{abcr} + \frac{a' + a'' \cdot fn}{abcr^2} + \frac{a' + a'' + a''' \cdot go}{abcr^3} + \&c.$$

$$- \frac{S}{3} \times \frac{a'md}{abcr} + \frac{a''en}{abcr^2} + \frac{a''' \cdot fo}{abcr^3} + \frac{a'''' \cdot gp}{abcr^4} + \&c. - \frac{S}{2r} \times \frac{a'en}{abcr} + \frac{a' + a'' \cdot fo}{abcr^2} +$$

$$\frac{a' + a'' + a''' \cdot gp}{abcr^3} + \&c.$$

In order to sum up the first and second of these series let  $\beta$  represent the number of persons living at the age of F, a person one year younger than B, and  $x$  the number of persons living at the age of K, a person one year younger than C. Let FK, BC, AFK, and ABC, represent the value of an annuity on the two and three joint lives of F and K, of B and C, of A, F and K, and of A, B and C respectively; then

will the series  $\frac{S}{6} \times \frac{a'bc}{abcr} + \frac{a''dm}{abcr^2} + \frac{a'''en}{abcr^3} + \&c.$  be  $= \frac{S \cdot \beta \cdot x}{6 \cdot bc} \times$

$$\frac{bc}{\beta x r} - \frac{a - a' \cdot bc}{a\beta x r} + \frac{dm}{\beta x r^2} - \frac{a - a' - a'' \cdot dm}{a\beta x r^2} - \frac{a'dm}{a\beta x r^2} + \frac{en}{\beta x r^3} - \frac{a - a' - a'' - a''' \cdot en}{a\beta x r^3}$$

$$- \frac{a' + a'' \cdot en}{a\beta x r^3}, \&c. = \frac{S \cdot \beta x}{bc} \times \frac{FK - AFK}{6} \left( - \frac{S}{6r} \times \frac{a'dm}{abcr} + \frac{a' + a'' \cdot en}{abcr^2}, \&c. \right.$$

$$\left. = - \frac{S}{6r} \times \frac{dm}{bcr} - \frac{a - a' \cdot dm}{abcr} + \frac{en}{bcr^2} - \frac{a - a' - a'' \cdot en}{abcr^2}, \&c. = \right) - \frac{S}{6r} \times$$

BC - ABC. The sum, therefore, of the two first series, or of

$$\frac{S}{6} \times \frac{a'bc}{abcr} + \frac{a'' \cdot dm}{abcr^2} + \&c. + \frac{S}{2r} \times \frac{a'dm}{abcr} + \frac{a' + a'' \cdot en}{abcr^2} + \&c. \text{ is } = \frac{S}{6} \times$$

$\frac{\beta x \cdot FK - AFK}{bc} + \frac{S}{3r} \times BC - ABC.$  Again, let P represent a life one year older than B, and let BK, PC, ABK, and APC, represent the values of annuities on the two and three joint lives of B and

B and K, P and C, A, B and K and of A, P and C: then the sum of the third and fourth series, or of  $-\frac{S}{6} \times$

$$\frac{a'mc}{abcr} + \frac{a''nd}{abcr^2} + \&c. - \frac{S}{2r} \times \frac{a'dn}{abcr} + \frac{a'+a'' \cdot eo}{abcr^2} + \&c. \text{ being } = -\frac{S \cdot x}{6c} \times$$

$$\frac{mc}{abr} - \frac{a-a' \cdot mc}{abxr} + \frac{dn}{bxr^2} - \frac{a-a'-a'' \cdot dn}{abxr^2} + \frac{eo}{bxr^3} - \frac{a-a'-a''-a''' \cdot eo}{abxr^3}, \&c.$$

$$\left( -\frac{S}{3r} \times \frac{a'dn}{abcr} + \frac{a'+a'' \cdot eo}{abcr^2}, \&c. = \right) - \frac{S \cdot m}{3br} \times \frac{a'dn}{acmr} + \frac{a'+a'' \cdot eo}{acmr^2} + \&c.,$$

will be  $= -\frac{S}{6} \times \frac{x \cdot BK - ABK}{c} - \frac{S}{3r} \times \frac{m \cdot PC - APC}{b}$ . The fifth se-

$$\text{ries, } \frac{S}{3} \times \frac{a'db}{abcr} + \frac{a''em}{abcr^2} + \&c., \text{ is } = \frac{S}{3} \times \frac{\beta}{b} \times \frac{db}{\beta cr} - \frac{a-a' \cdot db}{a\beta cr} + \frac{em}{\beta cr^2} -$$

$$\frac{a-a'-a'' \cdot em}{a\beta cr^2} + \frac{fn}{\beta cr^3} - \frac{a-a'-a''-a''' \cdot fn}{a\beta cr^3}, \&c. - \frac{S}{3r} \times \frac{a'em}{abcr} + \frac{a'+a'' \cdot fn}{abcr^2}, \&c.$$

= (putting FC and AFC for the values of the two and three joint lives of F and C and of A, F, and C)  $\frac{S}{3} \times \frac{\beta \cdot FC - AFC}{b} -$

$$\frac{S}{3r} \times \frac{a'em}{abcr} + \frac{a'+a'' \cdot fn}{abcr^2}, \&c. \text{ The sum, therefore, of the fifth and}$$

sixth series, or of  $\frac{S}{3} \times \frac{a'db}{abcr} + \frac{a''em}{abcr^2}, \&c. + \frac{S}{2r} \times \frac{a'em}{abcr} + \frac{a'+a'' \cdot fn}{abcr^2} + \&c.$

$$\text{is } = \frac{S}{3} \times \frac{\beta \cdot FC - AFC}{b} + \left( \frac{S}{6r} \times \frac{a'em}{abcr} + \frac{a'+a'' \cdot fn}{abcr^2}, \&c. = \right) \frac{S \times d}{6cr} \times$$

$$\frac{em}{bdr} - \frac{a-a' \cdot em}{abdr} + \frac{fn}{bdr^2} - \frac{a-a'-a'' \cdot fn}{abdr^2} + \&c. \text{ If T denote a life one}$$

year older than C, and BT, and ABT denote the values of the two and three joint lives of B and T and of A, B, and T, this

last series will be  $= \frac{S}{6r} \times \frac{d \cdot BT - ABT}{c}$ , and consequently the sum

of the fifth and sixth series will be  $= \frac{S}{3} \times \frac{\beta \cdot FC - AFC}{b} + \frac{S}{6r} \times$

$\frac{d \cdot BT - ABT}{c}$ . Lastly, the seventh and eighth series, or -

$$\frac{S}{3} \times \frac{d'md}{abc} + \frac{a''en}{abc^2} + \&c. - \frac{S}{2r} \times \frac{d'en}{abc} + \frac{a' + a'' \cdot fo}{abc^2} + \&c. = - \frac{S}{3} \times \frac{dm}{bc} - \frac{a - a' \cdot dm}{abc} + \frac{en}{bc^2} - \frac{a - a' - a'' \cdot en}{abc^2} + \&c. - \frac{S}{6r} \times \frac{a'en}{abc} + \frac{a' + a'' \cdot fo}{abc^2} + \&c.$$

$$= - \frac{S}{3} \times \overline{BC - ABC} - \left( \frac{S}{6r} \times \frac{md}{bc} \times \frac{en}{mdr} - \frac{a - a' \cdot en}{amdr} + \frac{of}{mar^2} - \frac{a - a' - a'' \cdot of}{amdr^2} + \&c. \right) \frac{S}{6r} \times \frac{md \cdot \overline{PT - APT}}{bc}$$

represent the values of the two and three joint lives of P and T, and of A, P, and T. If these several expressions be added together, &c. we shall at last have

$$\frac{S \cdot x}{6c} \times \frac{\beta \cdot \overline{FK - AFK}}{b} - \overline{BK - ABK} + \frac{S \cdot \beta}{3b} \times \overline{FC - AFC} - \frac{S \cdot r - 1}{3r} \times \overline{BC - ABC} - \frac{S \cdot m}{3br} \times \overline{PC - APC} + \frac{S \cdot d}{6cr} \times \overline{BT - ABT} - \frac{m \cdot \overline{PT - APT}}{b}$$

for the value of the sum S, when either B or C are the oldest of the three lives.

In order to determine the value of the reversion when the life of A is the oldest of the three lives, let *s, t, u, w, &c.* be the number of persons living at the end of the 1st, 2d, 3d, 4th, &c. years from the age of A, and let *b', b'', b''', b''''*, &c. be the decrements of life at the end of 1, 2, 3, 4, &c. years from the age of B; then, by reasoning as above, the value of the sum S for the

first year will be expressed by the series  $\frac{S \times b' \cdot \overline{a-s} \cdot \overline{c-d}}{6abc} + \frac{S \times b' \cdot \overline{a-s} \cdot d}{2abc}$ , for the second year by the series  $\frac{S \cdot b'' \cdot \overline{s-t} \cdot \overline{d-e}}{6abc^2} + \frac{S \cdot b'' \cdot \overline{s-t} \cdot e}{2abc^2} + \frac{S \cdot b'' \cdot a-s \cdot e}{abc^2} + \frac{S \cdot b'' \cdot \overline{a-s} \cdot \overline{d-e}}{2abc^2}$ , for the third year by the series  $\frac{S \cdot b''' \cdot \overline{t-u} \cdot \overline{e-f}}{6abc^3} + \frac{S \cdot b''' \cdot \overline{t-u} \cdot f}{2abc^3} + \frac{S \cdot b''' \cdot \overline{a-t} \cdot f}{abc^3} + \frac{S \cdot b''' \cdot \overline{a-t} \cdot \overline{e-f}}{2abc^3}$ , and so on for the remaining years of A's life.

These several series may be found =  $\frac{S}{abc} \times \frac{ab'c}{3} - \frac{b'cs}{6} - \frac{ab'd}{6} - \frac{b'ds}{3} + \frac{ab'd}{ab'd}$



$$\frac{ab'd}{2} + \frac{ab'c}{2} + \frac{s}{abc r^2} \times \left[ -\frac{sb''d}{3} - \frac{b'dt}{6} - \frac{sb''e}{6} - \frac{be''t}{3} + \frac{ab''e}{2} + \frac{ab''d}{2} + \frac{s}{abc r^3} \times \right. \\ \left. - \frac{tb'''e}{3} - \frac{b'''eu}{6} - \frac{tb'''f}{6} - \frac{b'''fu}{3} + \frac{ab'''f}{2} + \frac{ab'''e}{2} \right], \&c. \&c.$$

Let  $\alpha$  represent the number of persons living at the age of H, a person one year younger than A; let N denote a person one year older than A, and let the several combinations BN, BNC, AB, &c. denote, as in the former case, the values of annuities on the joint lives of B and N, of B, N and C, of A and B, &c.; then by proceeding in the same manner as in the fore-

going demonstration the series  $\frac{ab'c}{3abc r} + \frac{sb''d}{3abc r^2} + \frac{tb'''e}{3abc r^3} + \&c.$  may

be found =  $\frac{\alpha x}{3ac} \times \overline{HK - HBK} - \frac{AC - ABC}{3r}$ ; the series  $\frac{b'cs}{6abc r} + \frac{b'dt}{6abc r^2}$

+  $\frac{b'''eu}{6abc r^3} + \&c. = \frac{x}{6c} \times \overline{AK - ABK} - \frac{s}{6ar} \times \overline{NC - NBC}$ ; the series

$\frac{ab'd}{6abc r} + \frac{sb''e}{6abc r^2} + \frac{tb'''f}{6abc r^3} + \&c. = \frac{\alpha}{6a} \times \overline{HC - HBC} - \frac{d}{6cr} \times \overline{AT - ABT}$ ;

and the series  $\frac{b'ds}{3abc r} + \frac{b''et}{3abc r^2} + \frac{b'''fu}{3abc r^3} + \&c. = \frac{AC - ABC}{3} - \frac{sd}{3acr} \times$

$\overline{NT - NTB}$ . These four series, therefore, supposing them all

to be positive quantities are =  $\frac{x}{3c} \times \frac{\alpha \cdot \overline{HK - HBK}}{a} + \frac{\overline{AK - ABK}}{2} +$

$\frac{\alpha}{6a} \times \overline{HC - HBC} + \frac{r-1}{3r} \times \overline{AC - ABC} - \frac{s}{6ar} \times \overline{NC - NBC} - \frac{d}{3cr} \times$

$\frac{\overline{AT - ABT}}{2} + \frac{s \cdot \overline{NT - NTB}}{a}$ . With respect to the two remaining

series  $\frac{b'd}{2bcr} + \frac{b''e}{2bcr^2} + \frac{b'''f}{2bcr^2} + \&c. \dots$  and  $\frac{b'c}{2bcr} + \frac{b'd}{2bcr^2} + \frac{b'''e}{2bcr^3} + \&c.$ ,

these, it is evident, are to be continued after the decease of A till the extinction of the joint lives of B and C, and have been already proved in the solution of the second problem in my former Paper, to denote the value of the given sum payable if C should survive B. Let this value be represented

presented by R and the sum of the foregoing expressions (or  $\frac{x}{3c} \times \frac{a \cdot \overline{HK} - \overline{HBK}}{a} + \frac{\overline{AK} - \overline{ABK}}{2} + \frac{a}{6a} \times \overline{HC} - \overline{HBC}$ , &c.) by M, then will the value of the sum S (when A is the oldest of the three lives) be =  $S \times \overline{R} - M$ . Q. E. D.

If the three lives be equal, the value of the given sum for the first year will be =  $\frac{S \cdot \overline{c-d}^3}{6 \cdot c^3 \cdot r} + \frac{S \cdot \overline{c-d}^2 \cdot d}{2c^3 \cdot r} = S \times \frac{1}{6r} + \frac{2a^3}{6c^3r} - \frac{3dd}{6c^2r}$ ;

the value of the same sum for the second year will be =  $\frac{S \cdot \overline{d-e}^3}{6c^3r^2} + \frac{S \cdot \overline{d-e}^2 \cdot e}{2c^3r^2} + \frac{S \cdot \overline{d-e} \cdot \overline{c-d} \cdot e}{c^3r^2} + \frac{\overline{d-e}^2 \cdot \overline{c-d}}{2c^3r^2} = S \times$

$\frac{2e^3}{6c^3r^2} - \frac{3ee}{6c^2r^2} - \frac{2d^3}{6c^3r^2} + \frac{3dd}{6c^2r^2}$ ; the value for the third year will be =

$S \times \frac{2f^3}{6c^3r^3} - \frac{3ff}{6c^2r^3} - \frac{2e^3}{6c^3r^3} + \frac{3ee}{6c^2r^3}$ , and so on for the other years to

the extremity of life. Let CC and CCC denote the values of the two equal and three equal joint lives, the sum of these

series may then be found =  $\frac{S}{6} \times \frac{1}{r} + \frac{2CCC - 3CC}{r}$

= (supposing the perpetuity, or  $\frac{1}{r-1}$ , to be denoted by V)

$\frac{S}{6} \times \frac{r-1}{r} \times V - \frac{3 \cdot CC - 2CCC}{r}$ .

It must be here remembered, that from other principles it is well known, that the number of years purchase expressing the value of an *estate* or *perpetual annuity* to be entered upon at the failure of two out of any three equal lives is, “the difference  
“between three times the values of two equal joint lives, and  
“twice the values of three equal joint lives subtracted from  
“the perpetuity,” or  $V - \frac{3CC - 2CCC}{r}$ . The value, therefore, of such a reversion, supposing it to depend on the failure of the three equal lives in any one particular order, is (since

there are six such orders equally probable)  $\frac{1}{6} \times \overline{V - 3CC - 2CCC}$ . But it appears, from the correction explained in Dr. PRICE's Treatise on Reversionary Payments, Vol. I. p. 34. that the value of a reversionary *sum* is always less than the value of an equivalent reversionary *estate* in the proportion of 1 to  $r$ . The *sum* being  $S$  the equivalent estate or perpetual annuity is always  $S \times \overline{r - 1}$ ; and consequently the value of the *sum*  $S$  depending on the ceasing of three equal lives in any one particular order and thus determined, is the same with that determined by the foregoing investigation, that is,  $\frac{S}{6} \times \frac{\overline{r - 1}}{r} \times \overline{V - 3CC - 2CC}$ . The investigation, therefore, is right, and the correction and investigation demonstrate one another.

But the foregoing expression for determining the value of the reversion in this particular case is not only obtained immediately from the series, but also from the two different rules which have been given for determining the value when the lives are unequal; and hence a proof arises of the truth of these rules, as well as of the reasoning upon which they are founded. Thus the first rule, supposing the lives all equal,

becomes  $\frac{x^2}{c^2} \times \frac{KK - CKK}{6} - \frac{dd}{cc \cdot r} \times \frac{TT - CTT}{6} - \frac{\overline{r - 1}}{r} \times \frac{CC - CCC}{3} + \frac{x}{c} \times \frac{CK - CCK}{6} - \frac{d}{cr} \times \frac{CT - CCT}{6}$ , and the second rule becomes  $\frac{\overline{V - CC} \cdot \overline{r - 1}}{2r} - \frac{x^2}{c^2} \times \frac{KK - CKK}{3} + \frac{dd}{cc \cdot r} \times \frac{TT - CTT}{3} - \frac{\overline{r - 1}}{r} \times \frac{CC - CCC}{3} - \frac{x}{c} \times \frac{CK - CCK}{3} + \frac{d}{cr} \times \frac{CT - CCT}{3}$ . Let the value according to the first rule be denoted by  $L$ , and the second rule will be =  $\frac{\overline{r - 1} \cdot \overline{V - CC}}{2r} - 2L - \frac{\overline{r - 1} \cdot \overline{CC - CCC}}{r} (= L)$ . Hence  $3L =$

$$\frac{r-1 \cdot \overline{V} \quad \overline{CC} - 2 \cdot r \quad \overline{1 \cdot CC} - \overline{CCC}}{2r} \text{ and } L = \frac{r-1}{6r} \times \overline{V} - \overline{3CC} - \overline{2CCC}.$$

Q. E. D.

Were we possessed of complete tables of the values of annuities on two and three joint lives, the preceding rules would give an easy and exact solution of this problem in all cases. But as such tables, computed for every age, would be a work of immense difficulty, especially in regard to the values of three joint lives, Mr. SIMPSON'S rule for approximating to these from the given values of the two joint lives, has hitherto been adopted, and it seems upon the whole to answer the purpose very well. In the present problem it is attended with no other inconvenience than increasing the labour of the computations; for the values of the reversions derived from it appear in general to be perfectly correct. This is more fully ascertained by a table which Dr. PRICE has given in his Treatise on Reversionary Payments (Vol. II. Table 37.), of the values of three equal joint lives computed at 4 *per cent.* from the probabilities of life at NORTHAMPTON. By the assistance of this table, when the lives are of the same age, it is evident, from what has been already observed, that the exact value of the reversion may be easily obtained. The few following specimens computed from it, and compared with the values of the reversions deduced from the first and second of the preceding rules, demonstrate the accuracy of those rules: for, notwithstanding the approximated values of the three joint lives have been used in every instance in which the rules have been employed, yet the results approach so near the truth, even in the last stages of life, when the decrements are most irregular, that, though derived from these approximations, there can be little doubt of their correctness in almost every other period of life.

Common age.	Exact value * of £. 100. computed from Dr. PRICE'S Tables of the values of two and three equal joint lives.	Value of £. 100. computed from the first of the foregoing rules, and from Mr. SIMPSON'S approximation to the values of three joint lives.	Value of £. 100. computed from the second of the foregoing rules, and from Mr. SIMPSON'S approximation to the values of three joint lives.
70	- 12.000	- 12.005	- 12.000
75	- 12.944	- 12.943	- 12.943
80	- 13.840	- 13.810	- 13.880
85	- 14.450 †	- 14.670	- 14.340

Mr. DODSON †, and Mr. SIMPSON §, are the only writers who have solved, or rather who have approximated to the solution of this problem. But the former, by deducing his rules immediately from a wrong hypothesis, having rendered

\* That is, *one-sixth* part of the whole reversion.

† The several reversions in this column, when computed from SIMPSON'S approximation to the values of the three joint lives, are 12.012, 12.933, 13.847, and 14.803 respectively; which upon the whole differing nearly as much from the real values as those in the two other columns afford a convincing proof, that the very small deviation from the truth in these latter values proceeds not from any inaccuracy in the rules themselves, but solely from having used the *approximated* instead of the *real* values of the three joint lives. And this also will account for the difference in the values by the first and second rules. Were those values computed from tables which give the correct values of two and three joint lives at all ages, they would come out exactly the same. In the two first examples, where the values by one rule are true, it appears, that the values by the other rule are equally so. In the two last examples, where the values are not quite so accurate, it may be observed, that they differ as much in excess by one rule as they do in defect by the other; which must in general be the case from the very nature of those rules; for if  $L$  (or the value by the first rule) be greater than the truth, the difference between  $\frac{r-1 \cdot V-CC}{2r}$  and  $2L$  (or the value by the second rule) must be less than the truth; and, on the contrary, if  $L$  be less, this difference will be greater than the truth.

‡ See DODSON'S Mathematical Repository, Vol. III. Questions 42, 43, &c.

§ See SIMPSON'S Select Exercises, Prob. 38.

most of them (especially those in which three lives are concerned) of no use, it will be unnecessary to take notice of what he has done on the subject. With regard to the latter, whose rule is not only the sole guide for determining the value of this reversion, but also the source from which a great variety of other problems are solved, perhaps it may not be improper to examine how far his solution is to be depended upon; and the following examples have therefore been computed for this purpose.

T A B L E I.

Ages of			Value, by SIMPSON'S rule, of £. 100. payable on the contingent reversion specified in this pro- blem, when either C or B are el- dest, according to the <i>Northamp-</i> <i>ton Table</i> and at 4 per cent.			True value of the same reversion computed from the first rule in the fore- going solution.		
C.	B.	A.						
80	70	40	-	1.926	-	-		1.179
75	65	25	-	1.873	-	-		1.032
65	50	15	-	2.090	-	-		1.690
70	80	40	-	6.615	-	-		6.117
50	65	15	-	5.580	-	-		3.879
78	78	20	-	2.583	-	-		1.982
45	60	12	-	5.571	-	-		4.133
60	45	12	-	2.292	-	-		1.686

T A B L E II.

Ages.			Value of the same reversion by SIMPSON'S rule, when A is the oldest of the three lives.			True value of the same reversion by the second rule in the foregoing solution.		
C.	B.	A.						
24	65	75	-	34.636	-	-		31.792
65	24	75	-	6.305	-	-		7.895
49	9	69	-	7.351	-	-		5.960
18	78	78	-	37.554	-	-		33.019

T A B L E III.

Common Age.	Value of the same reversion by SIMPSON'S rule, when the ages of the three lives are equal.				True value of the same reversion.	
70	-	-	13.20	-	-	12.00
75	-	-	14.98	-	-	12.94
80	-	-	16.58	-	-	13.84
85	-	-	17.86	-	-	14.45

By comparing the values in the preceding tables, Mr. SIMPSON'S rule appears in almost every instance to be exceedingly incorrect. Even when the lives are equal (in which case it might have been expected to be sufficiently accurate) it seems to deviate, in old age at least, so widely from the truth as to be unfit for use. When C or B are eldest (which, however, is a case that does not often occur), the results sometimes exceed the truth *one-half*, and generally by more than *one-third* of the real value. When A is the oldest of the three lives (which is the most common case) these results are erroneous in nearly an equal degree. Nay, in some cases, Mr. SIMPSON'S rule is not only wrong but absurd. Thus, in the last example in the second table, the value of £. 100. payable on the contingency of C aged 18 surviving B aged 78 after A aged 78, is by this rule = £. 37.554. The value, therefore, of the same sum on the contingency of C's surviving A after B is also £. 37.554. Hence the value of £. 100. on the contingency of C's surviving A and B (without the restriction of one dying before the other) is  $2 \times 37.554 =$  £. 75.108\*. By another rule of Mr. SIMPSON †, the value

\* See SIMPSON'S Select Exercises, Prob. 39. † Ibid. Prob. 32.

of £. 100, on the contingency of C's surviving B only, is no more than £. 74\*. Now it is self-evident, that this latter value, instead of being *less*, ought to have been *greater* than the former, inasmuch as the probability of C's surviving only one life must be greater than that of his surviving two lives.

Many additional instances might be produced in which this rule, being made the basis upon which the solutions of other problems are founded, leads to conclusions equally erroneous. But these enquiries would be improper here; and I shall only observe, that had the foregoing examples been computed from the SWEDEN or LONDON, instead of the NORTHAMPTON Table, this rule would have appeared to be still more incorrect than it does from those computations.

When Mr. SIMPSON wrote his Select Exercises, he was in a great measure obliged to have recourse to DE MOIVRE's hypothesis, for want of those excellent tables of the real probabilities of life, and also of the values of single and joint lives which have been since published. Had he been possessed of these, it is most likely that his superior abilities would have directed him to a more accurate method of investigation. At present there can be no just reason for ever recurring to this wretched hypothesis. The solutions of all cases of two and even of three lives may be effected without much difficulty from principles strictly true. But I must here take my leave of this subject, hoping that its importance may engage other mathematicians to the further prosecution of it.

\* The true values are £. 66.038. and £. 74.884. respectively.





VI. *Result of Calculations of the Observations made at various Places of the Eclipse of the Sun, which happened on June 3, 1788. By the Rev. Joseph Piazzi, C. R. Professor of Astronomy in the University of Palermo; communicated by Nevil Maskelyne, D. D. F. R. S. and Astronomer Royal.*

Read January 15, 1789.

TO DR. MASKELYNE.

S I R,

THE satisfaction I had in observing the eclipse of the sun on the third of June last, with you and M. D'ARQUIER, at Greenwich, induces me to give you an account of the use made of the observations, and the consequences I have drawn from them. The observations which I have collected concerning the same eclipse, and which were made in other places, contribute to the extensiveness of my calculations, and to determine the position of certain places, which had not been before accurately determined, as that of Dublin, that of Mitau in Courland, and Perinaldo in Italy. The longitudes of all the other places must be referred to that of the Royal Observatory at Greenwich, as being the first in Europe, and because the observations which you have made in it, are by far more accurate than any others made elsewhere.

The

The result of the observation at Greenwich is confirmed in the most satisfactory manner by the observations of Oxford and Loampit-Hill, which Dr. HORNSBY and Mr. AUBERT have done me the honour to communicate to me. These three observations perfectly agree in the latitude of the moon; whence it follows, that the duration of the eclipse was justly observed. And whereas the difference of longitude for Oxford, as determined by these observations, is only one second, and that of Loampit-Hill only two seconds, different from that which had been determined by means of the best time-keepers and other most exact observations, it follows, that these three observations may be considered as a single one, having a treble degree of accuracy. In effect, if the moment of conjunction found for Oxford and Loampit-Hill be reduced to the meridian of Greenwich, by adding to the latter  $5''{,}4$ , and to the former  $5' 0''$ , and a mean be taken, it will appear, that this mean differs only by  $0{,}6$  of a second from the conjunction deduced from the observation made at Greenwich only. This also clearly proves, that the eclipses of the sun, when accurately observed, give nearly the same exactness as the occultations of the stars, which from their nature are considered as the most exact.

The observation made at Dublin deserves our greatest attention, particularly since the establishment of a very excellent observatory there. Dr. USSHER confesses, that the longitude of that city has not been exactly determined (see the Transactions of the Royal Irish Academy for the year 1787, p. 86.). He supposes the longitude of Dublin to be  $24' 58''$  W. which he determined by means of a time-keeper, which Mr. ARNOLD happened to take with him to that city; whereas I find it to be  $25' 13''{,}4$ . This my determination I believe not to leave the uncertainty of two seconds;

seconds, because the latitude of the moon deduced from the observation is the same with that of Greenwich.

The observation of Mitau is likewise very interesting; it shews the situation of a distant country, where no observation had been made before M. BEITLER established an observatory there. The observation of this able astronomer is of such correctness, that it furnishes the same latitude of the moon as the preceding ones; consequently the difference of the meridians, which I thence deduce, may be considered to be as exact as that of Dublin. The difference is 1 h. 34' 54" E, which becomes of the greater importance to geography, because from Pomerania to Petersburg no one point had been accurately determined before.

The position of Berlin has been already determined by means of some eclipses; but the results do not agree. The difference of 53' 32", which I have deduced, not only agrees exactly with that mentioned in Vol. IV. of M. DE LA LANDE'S Astronomy, and which this famous astronomer had deduced from the occultation of Antares observed by himself in the year 1749; but it also comes so near to the longitude mentioned by Mess. LEXELL and BERNOULLI, in the Ephemeris of Berlin, as not to differ by more than two seconds.

The observation of Vienna gives for the difference of the meridians 1 h. 5' 31". Though this determination differs but 1" from that found in the Almanack of Milan, and in the *Requisite Tables*, yet the observations of the two phases had not been very accurately made.

Perinaldo in Italy is a place whose position has not been as yet well determined. The tables requisite for the Nautical Ephemeris lay down this place at 30' 40" to the East of Greenwich; some place it at 30' 20". The observation made by

M. MARALDI, nephew of the Paris Academician, which I have by me, gives  $30' 53''$  E. for the difference of the meridians, and this may be considered as the best hitherto known.

The observations made at Milan, by the astronomers DE CESARIS and REGGIO, were interrupted by intervening clouds. In fact, the latitude of the moon in conjunction comes out only equal to  $14' 32''$ , which shews that the duration of the eclipse was not properly observed. I have thence also calculated the conjunction separately for the beginning and for the end of the eclipse, and I have found out the following differences of meridians, *viz.* for the beginning  $36' 39'',6$ , and for the end  $36' 38''$ ; and for the end and beginning conjointly  $36' 37''$ . This last difference comes nearest to that mentioned in the Milan Ephemeris for the year 1789, which is  $36' 41''$ . The observation made at Bologna assigns  $45' 28''$  E. for the difference of the meridians. But the duration of the eclipse was not properly observed. However, notwithstanding this imperfection, it may happen that the result determined is exact.

The two observations of France, *viz.* that of Viviers, and that of Rouen, give almost the same difference which I find in the Requisite Tables; that of Rouen differing only  $1''$ , and that of Viviers  $2''$ . As the difference of the meridians between Paris and Rouen is known with the greatest precision to be  $4' 57''$  to the W. of Paris; if to this difference are added  $4' 22'',3$ , which is the difference I found between Rouen and Greenwich, there will result, for the difference of the meridians between the Observatory of Greenwich and Paris,  $9' 19'',3$ . This difference only differs by  $0'',7$  from that established by Dr. BRADLEY, which is  $9' 20''$ , as adopted by yourself, and lately confirmed by Major-general ROY.

The

The observations of Loampit-Hill, Greenwich, and Oxford, as they serve for the basis of all my calculations, I have calculated them two different ways, *viz.* by the method of parallactic angles, and by the method of the nonagesimal, and the results agreed together within a few tenths of a second. By these two different methods I have also calculated the observations of Vienna, Berlin, and Viviers, in order to shew, that the different latitudes of the moon, given by the various observations, were not owing to any error in my calculations. For these places, in which both the beginning and the end of the eclipse had been observed, I have deduced the time of conjunction from the two phases conjointly, which have also given the duration of the eclipse, which cannot be obtained from a single observation.

The error of the tables which results from the observation at Greenwich is  $+26''$  in longitude, and  $+11'',5$  in latitude, at 20 h.  $58' 47'',3$  of apparent time, taking for the longitude of the sun  $2s. 14^{\circ} 16' 54'',7$ , as I deduce from the Nautical Almanack, and that of the moon at the same time to be greater than the sun by  $26''$ , as deduced from the same Almanack. I suppose also the horary motion of the moon in the ecliptic, by taking it an half hour before and after the conjunction, to be  $36' 52'' + 0'',6$  for the hour following the conjunction, and  $-0'',6$  for the hour preceding the conjunction; the moon's horary motion in latitude is  $3' 24'',3$ ; the horizontal parallax of the moon *minus* that of the sun at Greenwich, to be  $60' 14'',4$  for the commencement of the eclipse, and  $60' 16'',4$  for the end; the sun's diameter  $31' 34'',6$ , less by  $3''$  than that given in the Almanack, according to the correction which you have found necessary to be made; the moon's diameter I have stated as in the Almanack. In the opinion of M. DE LA LANDE,

some correction ought to be made to the parallax and to the diameter of the moon, as well as to the diameter of the sun; but on the one hand this would not make any alteration in the difference of the meridians which I have found; and on the other I thought proper to make use of those elements the Nautical Almanack furnished me with, that being a work the most perfect of the kind that ever appeared, and to which all astronomers and navigators ought to pay the greatest attention.

In fine, I compared the moon's longitude in conjunction deduced from the eclipse with the new tables of the moon corrected by Mr. MASON, and found the longitude by those tables to be  $2\text{ s. } 14^{\circ} 17' 6'',4$ , and the latitude to be  $15' 1'',3$ . The error then of the new Tables is  $+11'',7$  in longitude, and  $+13'',1$  in latitude; but M. DE LA LANDE having lately sent to me from Paris the place of the sun, calculated with the new Solar Tables (a most useful improvement which M. DE LAMBRE has, with much ingenuity, deduced from your observations) I find the error in longitude to be  $+27'',4$ , the sun's place being  $2\text{ s. } 14^{\circ} 16' 39'',0$  at  $20\text{ h. } 58' 47'',3$ .

The following table contains the observations of the eclipse, and the results deduced from thence. The first vertical column shews the name and place of the observers; the two next vertical columns contain all those observations which have been made, in apparent time; the other columns shew the results, *viz.* the fourth column, contains the true conjunction in apparent time; the fifth column contains the longitude of the moon in conjunction, which being always the same, needs not to be repeated under every perpendicular column; the sixth column contains the latitude of the moon, which, as it depends upon the manner of observing the two phases, is subject to

some variety; the seventh or last column contains the difference between the various meridians and that of Greenwich.

This, Sir, in brief, is the result which I have been able to deduce from the various observations above mentioned, and which I intirely submit to your judgement. If you think that it deserves to be made public, and in that case would be pleased to present this Paper to the Royal Society, I shall esteem myself extremely honoured and obliged by it.

I have the honour to be, &c.

JOSEPH PIAZZI.

Table of the observations made at various places on the eclipse of the sun, which happened June 3, 1788, and of results deduced from the same.

	Beginning.	End.	Conjunction.	Longitude of the moon in conjunction.	Latitude in conjunction	Difference of meridians.
Greenwich, Dr. MASKE- LYNE.	h. ' '' 19 24 46,5	h. ' '' 21 1 24,0	h. ' '' 20 58 47,3		' '' 14 48,2	0
Loampit- Hill, Mr. AUBERT.	19 24 41,9	21 1 20,3	20 58 44,1		14 48,2	'' 3,2W
Oxford, Dr. HORNSBY.	19 20 36,1	20 54 40,0	20 53 46,2		14 48,7	' '' 5 1,1W
Dublin, Dr. USSHER.	19 5 46,5	20 27 42,1	20 33 33,9		14 48,3	25 13,4W
Mittau, M. BEITLER.	21 20 15,0	23 8 52,0	22 33 41,5		14 48,7	h. 1 34 54,2E
Berlin, M. BODE.	20 23 9,0	22 14 32,0	21 52 20,3		14 44,2	0 53 33E
Vienna, M. TRIES- NEKER.	20 25 49,0	22 32 40,0	22 4 18,8	S. ° 2 14 16 54,7	14 39,0	1 5 31,5E
Viviers, M. FLAU- GERGUAS.	19 26 38,0	21 25 41,0	21 17 29,0		14 33,0	18 41,7E
Perinaldo, M. MARALDI.	19 37 50,0	*	21 29 40		*	30 53,0E
Rouen, M. DU LAGNE.	*	21 7 15,0	21 3 9,6		*	4 22,3E
Milan, Mess. DE CESARIS and REGGIO.	19 48 23,0	21 51 14,0	21 35 24,7		14 32,0	36 37,4E
Bologna, M. MATTEUCCI.	19 55 10,5	22 3 45,5	21 44 15,3		14 31,0	45 28E
Padua, M. CHIMI- NELLO.	19 59 20,0	22 6 58,0	21 46 21,3		14 39,0	47 34E



## P O S T S C R I P T.

IN the month of February last, I was favoured by Count DE BRUHL with the observation of the eclipse of the 4th of June last, made at Warsaw by M. BYSTRZYSKI; about the same time I also received of M. DE LA LANDE some other observations of the same eclipse, *viz.* those made at Prague, Marfeilles, Crefmunster, and Bagdad in Mesopotamia, which I immediately calculated, in order to add them to the others, which Dr. MASKELYNE lately did me the honour of presenting to the Royal Society.

The observation of Marfeilles confirms in the best manner the difference of meridians set down in the *Requisite Tables*, differing from that only by a second. The observation of Warsaw gives a difference ten seconds greater, and that of Crefmunster fourteen seconds less; which differences ought not to surprize us, considering the observations upon which the longitudes of these two places had been established; but, on the other hand, the observation of Prague clearly proves, that the situation of that town had been much less accurately determined than one might have expected. The time for the conjunction, which results from this observation, is the very same as that which is deduced by M. GERSTNER's new method, described in the Berlin Ephemeris for the year 1791, p. 243. From this time of conjunction the difference of meridians comes out equal to  $57' 42'', 7$ , *viz.* one minute and seventeen seconds less than that of the *Requisite Tables*.

The calculation of the observation made at Bagdad seems to indicate that there is some mistake with regard to the end of the eclipse, having found, that the difference of apparent longitude

gitude at the end is 20'' greater than the sum of the semi-diameters of the sun and moon, increased in proportion of the apparent altitude of the moon: for this reason I do not give the moon's latitude in conjunction. As for the time of the conjunction, I deduce it both from the two phases together, and from the commencement only, having previously corrected the moon's latitude of the error which I discovered in the tables, *viz.* 11'',6. The time of conjunction which results from the first calculation is 23 h. 56' 11''; that which results from the second 23 h. 56' 16'': this last nearly agrees in the difference of meridians with the Ephemeris of Paris for the year 1789, and differs from the Requisite Tables by 2' 32''.

The following table represents, as the first, the observations and the results.

	Beginning.	End.	Conjunction.	Longitude.	Latitude.	Difference of meridians.
Warsaw, M. BYSTRZYSKI	h. ' '' 20 56 45	h. ' '' 22 57 33	h. ' '' 22 22 59,3		' '' 14 44	h. ' '' 1 24 12
Prague, M. STRNADT.	20 21 29	22 21 15	21 56 30		14 45	0 57 42,7
Marseilles, M. BERNARD.	19 26 42	21 29 23,5	21 20 17,5		14 40	0 21 30,2
Cresmunster, M. FIXL- MILLNER.	20 15 20	22 19 50,7	21 54 59		14 23	0 56 11,7
Bagdad, M. DE BEAU- CHAMP.	22 30 51	23 26 19	23 56 11		*	2 57 23,7



VII. *An Account of a bituminous Lake or Plain in the Island of Trinidad. By Mr. Alexander Anderson; communicated by Sir Joseph Banks, Bart. P. R. S.*

Read February 19, 1789.

**A** MOST remarkable production of nature in the island of Trinidad, is a bituminous lake, or rather plain, known by the name of Tar Lake; by the French called La Bray, from the resemblance to, and answering the intention of, ship pitch. It lies in the leeward side of the island, about half-way from the Bocas to the south end, where the Mangrove swamps are interrupted by the sand-banks and hills; and on a point of land which extends into the sea about two miles, exactly opposite to the high mountains of Paria, on the north side of the Gulf.

This cape, or head-land, is about fifty feet above the level of the sea, and is the greatest elevation of land on this side of the island. From the sea it appears a mass of black vitrified rocks; but, on a close examination, it is found a composition of bituminous scoriæ, vitrified sand, and earth, cemented together; in some parts beds of cinders only are found. In approaching this Cape, there is a strong sulphureous smell, sometimes disagreeable. This smell is prevalent in many parts of the ground to the distance of eight or ten miles from it.

This point of land is about two miles broad, and on the east and west sides, from the distance of about half a mile

from the sea, falls with a gentle declivity to it, and is joined to the main land on the south by the continuation of the Mangrove swamps; so that the bituminous plain is on the highest part of it, and only separated from the sea by a margin of wood which surrounds it, and prevents a distant prospect of it. Its situation is similar to a Savannah, and, like them, it is not seen till treading upon its verge. Its colour, and even surface, present at first the aspect of a lake of water; but I imagine it got the appellation of Lake when seen in the hot and dry weather, at which time its surface to the depth of an inch is liquid, and then from its cohesive quality it cannot be walked upon.

It is of a circular form, and I suppose about three miles in circumference. At my first approach it appeared a plane, as smooth as glass, excepting some small clumps of shrubs and dwarf-trees that had taken possession of some spots of it; but when I had proceeded some yards on it, I found it divided into areolæ of different sizes and shapes: the chasms or divisions anastomosed through every part of it; the surface of the areolæ perfectly horizontal and smooth; the margins undulated, each undulation enlarged to the bottom till they join the opposite. On the surface the margin or first undulation is distant from the opposite from four to six feet, and the same depth before they coalesce; but where the angles of the areolæ oppose, the chasms or ramifications are wider and deeper. When I was at it, all these chasms were full of water, the whole forming one true horizontal plane, which rendered my investigation of it difficult and tedious, being necessitated to plunge into the water a great depth in passing from one areola to another. The truest idea that can be formed of its surface will be from the areolæ and their ramifications on the back of

a turtle. Its more common consistence and appearance is that of pit-coal, the colour rather greyer. It breaks into small fragments, of a cellular appearance and glossy, with a number of minute and shining particles interspersed through its substance; it is very friable, and, when liquid, is of a jet black colour. Some parts of the surface are covered with a thin and brittle scoria, a little elevated.

As to its depth, I can form no idea of it; for in no part could I find a substratum of any other substance; in some parts I found calcined earth mixed with it.

Although I smelt sulphur very strong on passing over many parts of it, I could discover no appearance of it, or any rent or crack through which the steams might issue; probably it was from some parts of the adjacent woods: for although sulphur is the basis of this bituminous matter, yet the smells are very different, and easily distinguished, for its smell comes the nearest to that of pitch of any thing I know. I could make no impression on its surface without an axe: at the depth of a foot I found it a little softer, with an oily appearance, in small cells. A little of it held to a burning candle makes a hissing or cracking noise like nitre, emitting small sparks with a vivid flame, which extinguishes the moment the candle is removed. A piece put in the fire will boil up a long time without suffering much diminution: after a long time's severe heat, the surface will burn and form a thin scoria, under which the rest remains liquid. Heat seems not to render it fluid, or occupy a larger space than when cold; from which, I imagine, there is but little alteration on it during the dry months, as the solar rays cannot exert their force above an inch below the surface. I was told by one Frenchman, that in the dry season the whole was an uniform smooth mass; and by another, that the ravins

contained water fit for use during the year; but neither can I believe: for if, according to the first assertion, it was an homogeneous mass, something more than an external cause must affect it, to give it the present appearances: nor without some hidden cause can the second be granted. Although the bottoms of these ramified channels admit not of absorption, yet from their open exposure, and the black surface of the circumjacent parts, evaporation must go on amazing quick, and a short time of dry weather must soon empty them; nor from the situation and structure of the place is there a possibility of supply but from the clouds. To shew that the progress of evaporation is inconceivably quick here, at the time I visited it, there were, on an average, two-thirds of the time incessant torrents of rain; but from the afternoon being dry, with a gentle breeze (as is generally the case during the rainy season in this island), there evidently was an equilibrium between the rain and the evaporation; for in the course of three days I saw it twice, and perceived no alteration on the height of the water, nor any outlet for it but by evaporation.

I take this bituminous substance to be the *bitumen asphaltum* LINNÆI. A gentle heat renders it ductile; hence, mixed with a little grease or common pitch, it is much used for the bottoms of ships, and for which intention it is collected by many, and I should conceive it a preservative against the Borer, so destructive to ships in this part of the world.

Besides this place, where it is found in this solid state, it is found liquid in many parts of the woods; and at the distance of twenty miles from this about two inches thick, round holes of three or four inches diameter, and often at cracks or rents. This is constantly liquid, and smells stronger of tar than when  
indurated,

indurated, and adheres strongly to any thing it touches; greafe is the only thing that will divest the hands of it.

The soil in general, for some distance round La Bray, is cinders and burnt earths; and where not so, it is a strong argillaceous soil; the whole exceedingly fertile, which is always the case where there are any sulphureous particles in it. Every part of the country, to the distance of thirty miles round, has every appearance of being formed by convulsions of nature from subterraneous fires. In several parts of the woods are hot springs; some I tried, with a well graduated thermometer of FAHRENHEIT, were  $20^{\circ}$  and  $22^{\circ}$  hotter than the atmosphere at the time of trial. From its position to them, this part of the island has certainly experienced the effects of the volcanic eruptions, which have heaped up those prodigious masses of mountains that terminate the province of Paria on the north; and no doubt there has been, and still probably is, a communication between them. One of these mountains opposite to La Bray in Trinidad, about thirty miles distant, has every appearance of a volcanic mountain: however, the volcanic efforts have been very weak here, as no trace of them extend above two miles from the sea in this part of the island, and the greater part of it has had its origin from a very different cause to that of volcanos; but they have certainly laid the foundation of it, as is evident from the high ridge of mountains which surrounds its windward side to protect it from the depredations of the ocean, and is its only barrier against that over-powering element, and may properly be called the skeleton of the island.

From every examination I have made, I find the whole island formed of an argillaceous earth, either in its primitive state, or under its different metamorphoses. The bases of the  
mountains

mountains are composed of *schistus argillaceus* and *taicum lithomargo*; but the plains or low lands remaining nearly in the same moist state as at its formation, the component particles have not experienced the vicissitudes of nature so much as the more elevated parts, consequently retain more of their primitive forms and properties. As argillaceous earth is formed from the sediment of the Ocean, from the situation of Trinidad to the Continent, its formation is easily accounted for, granting first the formation of the ridge of mountains that bound its windward side, and the high mountains on the Continent that nearly join it: for the great influx of currents into the Gulf of Paria from the coasts of Brazil and Andalusia must bring a vast quantity of light earthy particles from the mouths of the numerous large rivers which traverse these parts of the Continent; but the currents being repelled by these ridges of mountains, eddies and smooth water will be produced where they meet and oppose, and therefore the earthy particles would subside, and form banks of mud, and by fresh accumulations added would soon form dry land; and from these causes it is evident such a tract of country as Trinidad must be formed. But these causes still exist, and the effect from them is evident; for the island is daily growing on the leeward side, as may be seen from the mud-beds that extend a great way into the Gulf, and there constantly increase. But from the great influx from the Ocean at the south end of the island, and its egress to the Atlantic again, through the Bocas, a channel must ever exist between the Continent and Trinidad.





VIII. *An Account of a particular Change of Structure in the human Ovarium.* By Matthew Baillie, M. D.; communicated by John Hunter, Esq. F. R. S.

Read February 26, 1789.

THE ovaria in women are subject to a great variety of changes from their natural structure. Many of these are exactly similar to what take place in other parts of the body; but there is one which seems peculiar to them, the nature of which has probably not been hitherto very well ascertained. The change of structure to which I allude, is a conversion of the natural substance of an ovarium into a fatty mass, intermixed with hair and teeth. This sort of change is rare, although it occurs sufficiently often to have been seen by most persons who are very conversant in the examination of dead bodies. There are many cases of it related in the different books of dissections, but, as far as I have discovered, most commonly without any remarks upon the mode of formation\*; or they have been considered as very imperfect attempts at the

\* It has been the opinion of some, that hair, teeth, nails, feathers, &c. are animal vegetables or plants; and, agreeably to this opinion, Dr. Tyson considers the growth of hair and teeth in the ovarium as a *lujus naturæ*, where nature endeavours to produce something, and being disappointed in forming an animal, produces a vegetable. Vide Hooke's Lectures and Collection, N<sup>o</sup> II. p. 11. and 15.

growth

growth of a foetus in the ovarium, in consequence of connection between a male and a female. This conjecture rests no doubt on strong circumstances of probability, and yet there are many powerful reasons which seem to oppose its being well founded. Generation is a process always depending on the action of a certain cause, *viz.* the usual connection between a male and a female; and, when effects similar to those in generation are perceived, it becomes very natural to conclude, that this cause has been employed. The bias to such an opinion will become the stronger, from reflecting on the passions that are known to influence so powerfully mankind, by which the agency of this cause is frequently excited. When a change, therefore, was observed in an ovarium, by which it was converted into a fatty mass with hair and teeth, this should seem to correspond so much with a change taking place in consequence of generation, that the mind would scarcely entertain a doubt of its arising from the same cause, and would readily infer, that it had been preceded by a connection between the sexes. This doubt would still be the less, from the circumstance of a complete foetus being sometimes formed in the ovarium, where the usual means of generation had been employed. The following case, however, exhibits many reasons why we should be led to believe, that the ovaria in women have some power within themselves of taking on a process which is imitative of generation, without any previous connection with a male; and it is with this view that I proceed to relate it.

In a female child, about twelve or thirteen years old, which was lately brought to Windmill-street for dissection, I found the right ovarium converted into a substance, doughy to the touch, and about the size of a large hen's egg. Upon cutting into the substance, I found an apparently fatty mass, inter-  
mixed

mixed with hair and an excrescence of bones. This startled me very much, as I had always been led to believe, that such appearances were a sort of imperfect conception. The circumstances altogether being very singular, I was led to pay considerable attention to the change in the ovarium.

The fatty mass was of a yellowish white colour, in some places more yellow than in others, was very unctuous to the feeling, and consisted of shortened or separated particles, not having the same coalescence which the fat has generally in the body. It became very soft when exposed to the heat of a fire, and sunk into a portion of paper, on which it was spread, so as to make it more transparent. When the paper to which it was applied was exposed to the flame of a candle, it burnt with considerable crackling.

The hair with which the fatty substance was mixed grew out of the inner surface of the capsule containing it, in some places in solitary hairs, but chiefly in small fasciculi, at scattered irregular distances. Besides these, there were loose hairs involved in the fatty mass. The hairs were, some of them, of considerable length, even to three inches, were fine, and of a light-brown colour. They resembled much more the hairs of the head, than what are commonly found on the pubis, and corresponded very much in colour to the hair of the girl's head.

There arose also from the inner surface of the capsule some vestiges of human teeth. One appeared to be a canine tooth, another to be a small grinder, two others to be incisors, and there was also a very imperfect attempt at the formation of another tooth. These were not fully formed, the fangs being wanting; but in two of them the bodies were as complete as they are ever found in the common circumstances. They were each

of them inclosed in a proper capsule, which arose from the inner surface of the ovarium, and consisted of a white thick opaque membrane. Attached to the capsules of three of the teeth, there was a white spongy substance. The membrane of the ovarium itself was of some considerable thickness, but unequal in the different parts, was very smooth in its inner surface, and more irregular externally. The uterus was smaller than it is commonly at birth, was perfectly healthy in its structure, and upon opening into its cavity it exhibited the ordinary appearances of a child's uterus at that period. The left ovarium was very small, corresponding to the state of the uterus. It appears clearly from this, that the uterus had not yet received the increase of bulk, which is usual at the age of puberty. The hymen was entire, such as is commonly found in a child of the same age; and there was just beginning a *lanugo* upon the labia, not more than what is often found on the upper lip of a boy of fifteen years old. Such are the circumstances attending this singular case, and they present to the mind various grounds of consideration.

The formation of hair and teeth is a species of generation, for in fact it makes a part of it, and strikes the mind as being very different from any irregular substance which is formed by disease. This formation too takes place in a part of the body which is subservient to generation, and where a complete foetus is sometimes formed. The whole of this looks very much as if the production of hair and teeth in the ovarium was a sort of imperfect impregnation. But when we take another view of it, there are reasons at least equally strong for believing that such productions may arise from an action in the ovarium itself, without any stimulus from the application of the male semen.

In the case before us, the uterus was as small as at birth, indeed more so, and the left ovarium (which was perfectly healthy) corresponded to the state of the uterus. It had not been at all stimulated, nor did appear capable of being stimulated by the application of the male semen. This seems to be a strong circumstance; for in a case where there was an ovum formed in one of the Fallopian tubes, the uterus was enlarged to more than twice its unimpregnated size; and, upon opening into its cavity, the decidua was observed to be formed as completely as in the impregnated uterus. This preparation is still preserved in the collection of Windmill-street. Nothing can be a stronger proof, that when an impregnation takes place out of the cavity of the uterus, the uterus still takes a share in the action, and undergoes some of the changes of impregnation. In another preparation, which is preserved in the same collection, where there was a foetus formed in the ovarium, the uterus was increased to more than twice its ordinary size, was very thick and spongy, and had its blood-vessels enlarged as in an impregnated uterus. This becomes another very strong proof of the action of the uterus in the formation of an extra-uterine foetus. In the case before us, however, the uterus had undergone no change, and does not seem to have arrived at that period, when it could be capable of undergoing such a change.

Besides, we are not to consider the formation of teeth in the ovarium to be a quicker process than it is commonly in the head of a foetus; but in the present case the teeth having advanced fully as far as they are at some months after birth, this process must have begun at least more than a twelvemonth before the death of the child. If then we consider it as an impregnation, since the appearances of the child do not warrant us to believe her to have been more than twelve or thirteen years

old, this brings the date of the impregnation to an earlier period than can well be believed. From all these circumstances we might be led to suppose, that the formation of the hair and teeth was not in consequence of any connection with a male, but arose from some action of the ovarium itself, in which the uterus did not participate. The existence of the hymen, especially in so young a girl, becomes a collateral confirmation of the same opinion, although much is not to be rested on it, when taken singly.

It will, perhaps, have some influence in removing the prejudices against this opinion, to make the following remarks. Hair is occasionally formed in parts of the human body, which are absolutely unconnected with generation. Encysted tumours are sometimes found containing hair. Mr. HUNTER has a preparation of this sort in his collection, which he cut out from under the skin of the eyebrow. This tumour was perfectly complete, and unconnected with the skin, except by the common intervention of cellular membrane, so as to have no communication whatever with the hair of the eyebrow. In this instance there was certainly a species of generation taking place in the encysted tumour itself, forming hairs as completely and fully as in the common progress of the formation of a child. Such encysted tumours have been found in other parts of the human body, and still more frequently in quadrupeds. Mr. HUNTER has in his collection many specimens of encysted tumours from cows and sheep containing hair and wool. These were perfectly complete, so as to have possessed a power of production within themselves, and were many of them found deeply seated at a considerable distance from the skin, which is the common parent of hair. In these tumours there is often the appearance of layers of cuticle, which is probably

probably a preparatory step to the formation of hair. All this shews most clearly, that hair may be formed without any species of generation as it is commonly understood.


But hair is in itself as distinct a consequence of generation as teeth, and as much a peculiar substance. If then the one be formed, there appears to be no reason why the other should not also be formed. The action producing the one is not better understood than that producing the other; nor does it appear to be really in itself less connected with that species of generation arising from the approach of a male, so that teeth may probably be formed by a peculiar action taking place in the ovarium itself, as well as the hair.

It will tend to add further weight to this opinion, to consider that many of the adult teeth are formed in a child after birth; and therefore their formation depends on an action taking place in the jaws at a particular period, and not on original growth. The same circumstance strikes more strongly in the occasional formation of teeth at an advanced time of life. Both of these processes take place after the animal has been formed, in consequence of a certain action being excited in a particular part of the body, and therefore there is less difficulty in believing that the same sort of process may go on in another part of the body not commonly employed in it. It seems reasonable also to suppose, that the ovaria should have a greater aptitude of taking on a process somewhat similar to generation than the other indifferent parts of the body, as they constitute a part of the organs which are so materially concerned in the real process itself\*. These circumstances, when taken collectively, would

\* As the formation of teeth and hair involved in a fatty mass may be said to be peculiar to the ovaria, and as there are strong reasons for believing, that this formation

would seem to render it very probable, that the formation of hair and teeth in the ovarium does not necessarily depend on a connection between a male and a female (as has been the common opinion), but arises from some action within the ovarium itself, which is imitative of generation.

formation may take place without an intercourse between the sexes, it becomes difficult to account for this peculiarity in them, unless by supposing, that they have a greater aptitude of running into such a process than the other parts of the body.





IX. *Some Account of the Vegetable and Mineral Productions of Boutan and Thibet. By Mr. Robert Saunders, Surgeon at Boglepoor in Bengal; communicated by Sir Joseph Banks, Bart. P. R. S.*

Read February 26 and March 5, 1789.

**R**OAD to Buxaduar, May 11 and 12, 1783. The tract of country from Bahar to the foot of the hills contains but few plants that are not common to Bengal. Pine-apples, mango-tree, jack and faul timber, are frequently to be met with in the forests and jungles. Find many orange-trees towards the foot of the hills, of a very good sort, and bearing much fruit. Saw a few lime-trees, and found three different species of the sensitive plant. One species is used medicinally by the natives of Bengal in fevers; it is a powerful astringent and bitter; another is the species from which Terra Japonica is made, a medicine the history of which we are but lately made acquainted with. The third species is well known as the sensitive plant, and common in Bengal.

The country, from Bahar to the foot of the mountains, to which we approach without any ascent, is rendered one of the most unhealthy parts of India, from a variety of causes.

The whole, a perfect flat, is at all times wet and swampy, with a luxuriant growth of reeds, long grass, and underwood, in the midst of stagnated water, numerous frogs and insects. The exhalations from such a surface of vegetable matter and swamps, increased by an additional degree of heat from the reflection

reflection of the hills, affect the air to a considerable extent, and render it highly injurious to strangers and European constitutions.

The thermometer at the foot of the hill, mid-day  $86^{\circ}$ , fell to  $78^{\circ}$  at two o'clock, the time we reached Buxaduar, and that hour of the day when it is generally highest.

The soil and appearance of the ground in ascending the hill are materially changed. See many loose sparry stones and rock containing iron. Two springs, conducted from a distant height by spouts, are very pure and good water, without any mineral impregnation. The mountains in view covered with forests of trees, rendered useless from their inaccessibility. Those peculiar to the country are known to the natives by the names Boumbshi, Toumbshi and Sindeshi, besides faul timber, bamboo, and plantains.

Buxaduar, May 12 to 21. Many of the plants peculiar to Bengal require nursing at Buxaduar. There is one very good banian tree. In the jungles, met with the ginger, and a very good sort of yam; saw some pomegranate-trees in good preservation; shallots in great perfection; a species of the *Lychnis*, *Arum*, and *Asclepias*, natives of more northern situations, and of little use; a bad sort of raspberry, and a species of the *Gloriosa*. The plantains in use below do not thrive here. In the jungles they have a plantain-tree producing a very broad leaf, with which they cover their huts; but the fruit is not eaten. See many weeds and long grass more common to Bengal than any other parts of Boutan.

From the 15th to the 22d, the rains were almost incessant at Buxaduar. Our People became unhealthy, and were attacked with fevers, which, if neglected in the beginning, proved obstinate quartans. This was the case with several of  
the

the natives whom I had an opportunity of seeing. They scarcely, however, admit that Buxaduar is unhealthy at any season of the year. After allowing for their prejudice, and the possibility of the natives suffering but little from the bad seasons, I cannot help thinking, that Buxaduar must be unhealthy, at least to strangers, from the month of May till towards the end of September. It lies high, but is overtopped by the surrounding mountains, covered with forests of trees and underwood. In all climates, where the influence of the sun is great, this is a never-failing cause of bad air. The exhalation that takes place from so great a surface in the day-time falls after sunset in the form of dew, rendering the air raw, damp, and chilly, even in the most sultry climates.

The thermometer at Buxaduar was never, at two o'clock in the afternoon, above  $82^{\circ}$ , or below  $73^{\circ}$ .

In the neighbourhood of Buxaduar there are several excellent springs of water, some of them with less impregnation of any sort than I ever met with; the nicest test scarcely produced the separation of a sensible quantity of earthy matter. Such waters are generally to be distinguished by the taste, which is insipid and unpleasant. When these springs could be traced to their source, they sunk the thermometer eight or ten degrees below the temperature of the atmosphere.

Road to Murishong, May 22 and 23. In ascending the hill from Buxaduar there is to be seen much of an imperfect quartz, of various forms and colour, having in some places the appearance of marble; but from chemical experiments, it was found to possess very different properties. This sort of quartz, when of a pure white, and free from any metallic colouring matter, is used as an ingredient in porcelain. I have not seen any that promises to answer that purpose better than what is to be met with in the mountains near

Buxaduar. It is known to mineralists in that state by the name of quartz gritstone. The rock which forms the basis of these mountains dips in almost every direction, and is covered with a rich and fertile soil, but in no place level enough to be cultivated. Many European plants are met with on the road to Murishong; many different sorts of mosses, fern, wild thyme, peaches, willow, chickweed, and grasses common to the more southern parts of Europe, nettles, thistles, dock, strawberry, raspberry, and many destructive creepers, some peculiar to Europe.

Murishong is the first pleasant and healthy spot to be met with on this side of Boutan. It lies high, and much of the ground about it is cleared and cultivated; the soil, rich and fertile, produces good crops. The only plant now under culture is a species of the Polygonum of LINNÆUS, producing a triangular seed, nearly the size of barley, and the common food of the inhabitants. It was now the beginning of their harvest; and the ground yields them, as in other parts of Boutan, a second crop of rice. Here are to be found in the Jungles two species of the Laurus of LINNÆUS; one known by the name of the bastard cinnamon. The bark of the root of this plant, when dried, has very much the taste and flavour of cinnamon; it is used medicinally by the natives. The Chenopodium, producing the semen fantonicum, or worm-feed, a medicine formerly in great character, and used in those diseases from which it is named, is common here.

Found in the neighbourhood of this place all the European plants we had met with on the road. The ascent from Buxaduar to Murishong is upon the whole great, with a sensible change in the state of the air.

Road to Chooka, May 25. On the road to Chooka find all the Murishong plants, cinnamon-tree, willow, and one or two firs;

firs; strawberries every where, and very good, and a few bilberry plants.

Much sparry flint, and a sort of granite with which the road is paved. There is a great deal of talc in the stones and soil, but in too small pieces to be useful. Frequent beds of clay and pure sand. Find two mineral wells, slightly impregnated with iron, with much appearance of that metal in this part of the country; and they are not unacquainted with the method of extracting it from the stones, but still despise its use in building. Towards Chooka there are many well cultivated fields of wheat and barley.

Road to Punekha, May 26. From Chooka the country opens, and presents to view many well cultivated fields and distant villages; a rapid change in climate, the vegetable productions, and general appearance of the country. Towards Punekha, pines and firs are the only trees to be met with; but they do not yet seem in their proper climate, being dwarfish and ill-shaped; peaches, raspberries, and strawberries, thriving every where; scarce a plant to be seen that is not of European growth. In addition to the many I have already mentioned, saw two species of the *Cratægus*, one not yet described. Saw two ash-trees in a very thriving state, the star-thistle, and many other weeds, in general natives of the Alps and Switzerland.

Much of the rock to-day is, I find on examination, pure limestone; a valuable acquisition if they did not either despise its use, or were unacquainted with its properties. It was most advantageously situated for being worked, and the purest perhaps to be met with. There is likewise abundance of fire-wood in this part of the country. In building they would derive great benefit from the use of it. Their houses are lofty, the timbers substantial, and nothing wanting

to make them durable, but their being acquainted with the use of lime. As a manure it might probably be used to great advantage. Many fields of barley in this part of the country; now the beginning of their harvest. The thermometer here fell, at four o'clock in the afternoon, to  $60^{\circ}$ : cold and chilly.

Road to Chepta, May 27. On the road to Chepta, the rock in general dips to the northward and eastward, in about an angle of sixty degrees. Much of limestone, and some veins of quartz, and loose pieces of sparry flint striking fire with steel.

Several springs, and one slightly impregnated with iron.

In addition to the plants of yesterday, find the *Coriandrum testiculatum*, *Inula montana*, and *Rhododendrum magnum*.

At Chepta met with a few turneps, one maple-tree, wormwood, goose-grass (*Galium aparine*), and many other European weeds; the first walnut-tree we had yet seen.

Chepta lies high, and not above six miles from the mountain of Lomyla, now covered with snow. The wind from that quarter, S.E. made it cold and chilly, and sunk the thermometer at mid-day to  $57^{\circ}$ . Here are some fields of wheat and barley not yet ripe.

Road to Pagha, May 29. Soon after leaving Chepta find a mineral well, which, on a chemical examination, gave marks of a strong impregnation from iron. I traced it to its source, where the thermometer, on being immersed, fell from  $68^{\circ}$  to  $56^{\circ}$ .

A little before we reach Pagha, met with some limestone, and a bed of chalk, which, near the surface, contained a great proportion of sand, but some feet under was much purer.

The forests of firs of an inferior growth, several ash-trees, dog-rose, and bramble.

Road to Taffesudon, May 30 and 31, June 1. The road from here to Taffesudon presents us with little that we have not met with; fewer strawberries, and no raspberries; some very good orchards of peaches, apricots, apples, and pears. The fruit formed, and will be ripe in August and September. Met with two sorts of cranberry, one very good. Saw the *Fragaria sterilis*, and a few poppies. At Wanakha found a few turneps, shallots, cucumbers, and gourds. Near to Taffesudon the road is lined with many different species of the rose, and a few jessamine plants. The soil is light, and the hills in many places barren, rocky, and with very little verdure. The rock in general laminated and rotten, with many small particles of talc in every part of the country incorporated with the stones and soil. Some limestone, and appearance of good chalk. Several good and pure springs of water.

Taffesudon and its neighbourhood abound with all the plants we have already mentioned. The hills are chiefly wood, with firs and aspen. I have not yet been able to find an oak-tree, and the ash is very seldom to be met with. The elder, holly, bramble, and dog-rose, are common. Found the birch-tree, cypress, yew, and delphinium. Many different species of the vaccinium, of which the bilberry is one, and the cranberry another. Towards the top of the adjacent mountains met with two plants of the *Arbutus uva ursi*, which is a native of the Alps, the most mountainous parts of Scotland, and Canada.

I have likewise seen a species of the rhubarb plant (*Rheum undulatum*) brought from a distance, and only to be met with near the summits of hills covered with snow, and where the  
soil

foil is rocky. The true rhubarb (*Rheum palmatum*) is likewise the native of a cold climate; and though China supplies us with much of this drug, it is known to be the growth of its more northern provinces, Tartary, and part of the Russian dominions. The great difficulty is in drying the root. People versant in that business say, that one hundred pounds of the fresh root should not weigh above six pounds and a half, if properly dried, and it certainly has been reduced to that. I have seen eighty pounds of fresh root produced from one plant; but, after drying it with much care and attention, the weight of the dried root could not be made less than twelve pounds. It was suspended in an oven, with an equal and moderate degree of heat. Little more than the same quantity of this powder produced a similar effect with the best foreign rhubarb.

The other plants common here are the service-tree, blessed thistle, mock orange, *Spiræa filipendula*, *Arum*, *Echites*, *Punica*, *Ferula communis*, *Erica*, and *Viola*. Of the rose-bush I have met with the five following species; *Rosa alpina*, *centifolia*, *canina*, *Indica*, *spinossissima*.

The culture of pot-herbs is every where neglected; turneps, a few onions and shallots, were the best we could procure. Mr. BOGLE left potatoes, cabbage, and lettuce-plants, all which we found neglected and dispersed. They had very improperly (from an idea most probably of their being natives of Bengal) planted them in a situation and climate which approaches very near to that of Bengal at all seasons, as we shall find afterwards. Melons, gourds, brinjals, and cucumbers, are occasionally to be met with. The country is fitted for the production of every fruit and vegetable common without the tropics, and in some situations will bring to perfection many of the tropical fruits.



There are two plants which I have to regret the not having had as yet an opportunity of seeing; one is the tree from the bark of which their paper is made; and the other is employed by them in poisoning their arrows. This last is said to come from a very remote part of the country. They describe it as growing to the height of three or four feet, with a hollow stalk. The juice is inspissated, and laid as a paste on their arrows. Fortunately for them, it has not all the bad effects they dread from it. I had an opportunity of seeing several who were wounded with these arrows, and they all did well, though under the greatest apprehension. The cleaning and enlarging some of the wounds was the most that I found necessary to be done. The paste is pungent and acrid, will increase inflammation, and may make a bad or neglected wound mortal; but it certainly does not possess any specific quality as a poison.

The fir, so common in this country, is perhaps the only tree they could convert to a useful and profitable purpose. What I have seen would not, from their situation, be employed as timber. The largest I have yet met with were near Wandepore; they measured from eight to ten feet in circumference, were tall and straight. Such near the Burrampooter, or any navigable river, might certainly be transported to an advantageous market. I am convinced that any quantity of tar, pitch, turpentine, and resin, might be made in this country, much to the emolument of the natives. Firs, which from their size and situation are unfit for timber, would answer the purpose equally well. The process for procuring tar and turpentine is simple, and does not require the construction of expensive works. This great object has been so little attended to, that they are supplied from Bengal with what they want of these articles.

The

The country about Taffesudon contains great variety of soil, and much rock of many different forms, but still an unpromising field for a mineralist. I have not found in Boutan a fossil that had the least appearance of containing any other metal than iron, and a small portion of copper. From information, and the reports of travellers, I believe it is otherwise to the northward. The banks of the Ticushu, admitting of cultivation for several miles above and below Taffesudon, yield them two crops in the year. The first of wheat and barley is cut down in June; and the rice, planted immediately after, enjoys the benefit of the rains. This country is not without its hot wells, as well as many numerous springs, some of which I have taken notice of. One hot well, near Wandepore, is so close to the banks of the river as to be overflowed in the rains, and we found it impossible to get to it: the heat of this well is great; but I could not learn that the ground about it was much different from the general aspect of the country. Another, several days journey from hence, is on the brow of a hill perpetually covered with snow. This hot well is held in great estimation by the people of the country, and resorted to by valetudinarians of every description. I gained but little satisfactory information respecting the degree of heat, or appearance of the ground about it, that could lead me to form a just opinion of either.

Taffesudon to Paraghon, Sept. 8 and 9. Left Taffesudon, and arrived next day at Paraghon. Much good rich soil, with more pasture, where the ground is not cultivated, than we had yet met with. Many fields of turneps in great perfection; a plant they seem better acquainted with the cultivation of than any other. Find on the road many large and well-thriving birch, willows, pines, and firs, some walnut-trees, the  
Arbutus

*Arbutus uva ursi*, abundance of strawberry, elderberry, bilberry, *Chrysanthemum* or greater daisy, and many European grasses. See the *Datura ferox* or thorn-apple, a plant common in China and some parts of Thibet, where it is used medicinally. They find it a powerful narcotic, and give the seeds where they wish that effect to be produced. It has been used as a medicine in Europe, and is known to possess these qualities in a high degree. See holly, dog-rose, and aspen. The present crop near Paraghon, on the banks of the Pachu, is rice, but not so far advanced as at Tasséfudon; the same may be said of their fruits. They say it is colder here at all seasons than at Tasséfudon, which is certainly below the level of this place.

Towards the summit of the mountain we crossed, found some rock of a curious appearance, forming in front six or seven angular semi-pillars, of a great circumference, and some hundred feet high. This natural curiosity was detached in part from the mountain, and projected over a considerable fall of water, which added much to the beautiful and picturesque appearance of the whole. Numerous springs, some degrees colder than the surrounding atmosphere, gushing from the rock on the most elevated part of the mountain, furnish a very ample and seasonable supply of excellent water to the traveller. The rock, in many places laminated, might be formed into very tolerable slate. Near to Paraghon iron stones are found, and one spring highly impregnated with this mineral.

Road to Dukaigun, Sept. 11. Our road to Dukaigun, nearly due north, is a continued ascent for eight miles, along the banks of the Pachu, falling over numerous rocks, preci-

pices, and huge stones. Here we begin to experience a very considerable change in the temperature of the atmosphere; the surrounding hills were covered with snow in the morning, which had fallen the preceding night, but disappeared soon after sunrise. The thermometer fell to  $54^{\circ}$  in the afternoon, and did not rise above  $62^{\circ}$  at noon.

The face of the mountains, in some places bare, with projecting rock of many different forms; quartz, flint, and a bad sort of freestone, common. Many very good springs, slightly impregnated with a selenitic earth.

The soil is rich, and near to the river in great cultivation. Many horses, the staple article of their trade, are bred in this part of the country. Found walnut-trees, peaches, apples, and pears.

Road to Sanha, Sept. 12. The road still ascending to Sanha, and near to the river for ten miles.

The thermometer falling some degrees, we found it cold and chilly. The bed of the river is full of large stones, probably washed down from the mountains by the rapidity of its stream; they are chiefly quartz and granite. Here is excellent pasture for numerous herds of goats.

Road to Chichakumboo. From Sanha the ascent is much greater, and, after keeping for ten miles along the banks of the Pachu, still a considerable stream, we reach its source (from three distinct rivulets, all in view, ramified and supplied by numerous springs), and soon after arrive at the most elevated part of our road.

Here we quit the boundary of Boutan, and enter the territory of Thibet, where nature has drawn the line still more strongly, and affords, perhaps, the most extraordinary contrast

traft that takes place on the face of the earth. From this eminence are to be feen the mountains of Boutan, covered with trees, fhubs, and verdure to their tops, and on the fouth fide of this mountain to within a few feet of the ground on which we tread. On the north fide the eye takes in an extenfive range of hills and plains, but not a tree, fhub, or fcarce a tuft of grafs to be feen. Thus, in the courfe of lefs than a mile, we bid adieu to a moft fertile foil, covered with perpetual verdure, and enter a country where the foil and climate feem inimical to the production of every vegetable. The change in the temperature of the air is equally obvious and rapid. The thermometer in the forenoon  $34^{\circ}$ , with froft and fnow in the night-time. Our prefent obfervations on the caufe of this change confirmed us in a former opinion, and incontestably prove, that we are to look for that difference of climate from the fituation of the ground as more or lefs above the general level of the earth. In attending to this caufe of heat or cold, we muft not allow ourfelves to be deceived by a comparifon with that which is immediately in view. We ought to take in a greater range of country, and where the road is near the banks of a river, we cannot well err in forming a judgement of the inclination of the ground. Pujukha and Wandepore, both to the northward of Taffesudon, are quite in a Bengal climate. The thermometer at the firft of thefe places, in the months of July and January, was within two degrees of what it had been at Rungpore for the fame periods. They feem in more expofed fituations than Taffesudon; and, were we to draw a comparifon of their heights from the furrrounding ground, I fhould fay they were above its level. The road, however, proves the reverfe. From Pujukha to

Taffefudon we had a continued and steep ascent for six hours and a half, with a very inconsiderable descent on the Taffefudon side. From the south side of the mountain dividing Boutan from Thibet, the springs and rivulets are tumbling down in cascades and torrents, and have been traced by us near to the foot of the hills, where they empty themselves to the eastward of Buxaduar. On the north side they glide smoothly along, and by passing to the northward as far as Tishoolumboo, prove a descent on that side, which the eye could not detect. This part of the country, being the most elevated, is at all times the coldest; and the snowy mountains, from their heights and bearings, notwithstanding the distance, are certainly those seen from Purnea.

The soil on the Thibet side of the mountain is sandy, with much gravel and many loose stones. On the road found the *Aconitum pyreneum*, and two species of the *Saxifraga*.

See a large flock of chowry tailed cattle; their extensive range of pasture seems to make amends for its poverty.

From Faro to Duina, Sept. 15. From Faro to Duina pass over an extensive plain, bounded by many small hills, oddly arranged; some of them detached and single, and all seem composed of sand collected in that form, having the plain for their general base.

At Duina found a few plots of barley, which they are now cutting down, though green, as despairing of its ripening. The thermometer, at six o'clock in the morning, below the freezing point, and the ground partially covered with snow.

Road to Chalu, Sept. 16. Continue on the plain; find three springs forcing their way through the ground with violence, and giving rise to a lake many miles in extent, stored

with millions of water-fowl and excellent fish. Of the first saw the cyrus, solan geese, many kinds of ducks, pintados, cranes, and gulls of different sorts. The springs of this lake are in great reputation for the cure of most diseases. I examined the water, and found it contain a portion of alum with the selenitic earth. On the banks of the lake I found a crystallization, which proves to be an alkaline salt; it is used by the natives for washing, and answers the purpose as well as pot-ash. The pasture which is impregnated with this salt is greedily sought after by sheep and goats, and proves excellent food for them. The hills are chiefly composed of sand incruusted by the inclemency of the weather and violent winds, seeming at first view composed of freestone.

Road to Simadar, Sept. 17. Pass a lake still more considerable than the former, with which it communicates by a narrow stream, about three miles long. There never was a more barren or unpromising soil; little turf, grass, or vegetation of any sort, except near the lake. See a few huts, mostly in ruins and deserted. The only grain in this part of the country is barley, which they are cutting down every where green.

Pass two springs, one of them slightly impregnated with alum. They form the principal source of a river, which empties itself in the Burrampooter near Tiffolumboo.

The wind from the eastward of south is now the coldest and most piercing; passing over the snowy mountains and dry sandy desert before described, it comes divested of all vapour or moisture, and produces the same effect as the hot dry winds in more southerly situations. Mahogany boxes and furniture, that had withstood the Bengal climate for years, were warped  
with

with considerable fissures, and rendered useless. The natives say, a direct exposure to these winds occasions the loss of their fore-teeth; and our faithful guide ascribed that defect in himself to this cause. We escaped with loss of the skin from the greatest part of our faces.

Road to Seluh, Sept. 18. Near our road to-day found a hot well, much frequented by people with venereal complaints, rheumatism, and all cutaneous diseases. They do not drink the water, but use it as a bath. The thermometer, when immersed in the water, rose from  $40^{\circ}$  to  $88^{\circ}$ . It has a strong sulphureous smell, and contains a portion of hepar sulphuris. Exposure to air deprives it, as most other mineral wells, of much of its property.

Road to Takui, Sept. 19. Pass some fields of barley and pease, and get into a milder climate. Find to-day a great variety of stone and rock, some containing copper, and others, a very pure rock crystal, regularly crystallized, with six unequal sides. The rock crystal is of different sizes and degrees of purity, but of one form. Find some flint and granite, several springs of water impregnated with iron, and nearly of the same temperature with the atmosphere. See a few ill-thriving willows planted near the habitations, and which are the only trees to be met with.

Road to Tiffolumboo, Sept. 20, 21, and 22. The remaining part of our journey is over a more fertile soil, enjoying a milder climate. Some very good fields of wheat, barley, and pease; many pleasant villages and distant houses, less sand and more rock, part flaty, and much of it a very good sort of flint. The soil in the valley a light-coloured clay and sand. They are every where employed in cutting down their crop. What



a happy climate! The sky serene and clear, without a cloud; and so confident are they of the continuance of this weather, that their crop is thrown together in a convenient part of the field, without any cover, to remain till they can find time to thresh it out,

Before we reached Tissoolumboo found some elms and ash-trees.

The hills in Thibet have, from their general appearance, strong marks of containing those fossils that are inimical to vegetation; such are most of the ores of metal and pyritical matter.

The country, properly explored, promises better than any I have seen to gratify the curiosity of a philosopher, and reward the labours of a mineralist. Accident, more than a spirit of enterprise and enquiry, has already discovered the presence of many valuable ores and minerals in Thibet. The first in this list is deservedly gold. They find it in large quantities, and frequently very pure. In the form of gold-dust it is found in the beds of rivers, and at their several bendings, generally attached to small pieces of stone, with every appearance of its having been part of a larger mass. They find it sometimes in large masses, lumps, and irregular veins; the adhering stone is generally flint or quartz, and I have sometimes seen a half-formed, impure sort of precious stone in the mass. By a common process for the purification of gold, I extracted 12 *per cent.* of refuse from some gold-dust, and on examination found it to be sand and filings of iron, which last was not likely to have been with it in its native state, but probably employed for the purpose of adulteration. Two days journey from Tissoolumboo there is a lead mine. The ore is much the same as that found in Derbyshire, mineralized by sulphur,

fulphur, and the metal obtained by the very simple operation of fusion alone. Most lead contains a portion of silver, and some in the proportion to make it an object to work the lead ore for the sake of the silver. Cinnabar, containing a large proportion of quicksilver, is found in Thibet, and might be advantageously employed for the purpose of extracting this metal. The process is simple, by distillation; but to carry it on in the great would require more fuel than the country can well supply. I have seen ores and loose stones containing copper, and have not a doubt of its being to be found in great abundance in the country. Iron is more frequently to be met with in Boutan than Thibet; and, was it more common, the difficulty of procuring proper fuel for smelting the less valuable ores, must prove an insuperable objection to the working them. The dung of animals is the only substitute they have for fire-wood, and with that alone they will never be able to excite a degree of heat sufficiently intense for such purposes. Thus situated, the most valuable discovery for them would be that of a coal mine. In some parts of China bordering on Thibet, coal is found and used as fuel.

Tincal, the nature and production of which we have only hitherto been able to guess at, is now well known, and Thibet, from whence we are supplied, contains it in inexhaustible quantities. It is a fossil brought to market in the state it is dug out of the lake, and afterwards refined into Borax by ourselves. Rock salt is likewise found in great abundance in Thibet.

The lake, from whence tincal and rock salt are collected, is about fifteen days journey from Tiffoolumboo, and to the northward of it. It is encompassed on all sides by rocky hills, without

without any brooks or rivulets near at hand; but its waters are supplied by springs, which being saltish to the taste are not used by the natives. The tincal is deposited or formed in the bed of the lake; and those who go to collect it, dig it up in large masses, which they afterwards break into small pieces for the convenience of carriage, exposing it to the air to dry. Although tincal has been collected from this lake for a great length of time, the quantity is not perceptibly diminished; and as the cavities made by digging it soon wear out or fill up, it is an opinion with the people, that the formation of fresh tincal is going on. They have never yet met with it in dry ground or high situations, but it is found in the shallowest depths, and the borders of the lake, which deepening gradually from the edges towards the center contains too much water to admit of their searching for the tincal conveniently; but from the deepest parts they bring up rock salt, which is not to be found in the shallows, or near the bank. The waters of the lake rise and fall very little, being supplied by a constant and unvarying source, neither augmented by the influx of any current, or diminished by any stream running from it. The lake, I am assured, is at least twenty miles in circumference, and standing in a very bleak situation is frozen for a great part of the year. The people employed in collecting these salts are obliged to desert from their labour so early as October, on account of the ice. Tincal is used in Thibet for soldering and to promote the fusion of gold and silver. Rock salt is universally used for all domestic purposes in Thibet, Boutan, and Naphaul.

The thermometer at Tissoolumboo during the month of October was, on an average, at eight o'clock in the morning  $38^{\circ}$ , at noon  $46^{\circ}$ , and six o'clock in the evening  $42^{\circ}$ . The

weather clear, cool, and pleasant, and the prevailing wind from the southward. During the month of November we had frosts morning and evening, a serene clear sky, not a cloud to be seen. The rays of the sun passing through a medium so little obscured had great influence. The thermometer was often below  $30^{\circ}$  in the morning, and seldom above  $38^{\circ}$  at noon in the shade; wind from the southward.

Of the diseases of this country, the first that attracts our notice, as we approach the foot of the hills, is a glandular swelling in the throat, which is known to prevail in similar situations in some parts of Europe, and generally ascribed to an impregnation of the water from snow. The disease being common at the foot of the Alps, and confined to a tract of country near these mountains, has first given rise to the idea of its being occasioned by snow water. If a general view of the disease, and situations where it is common, had been the subject of enquiry, or awakened the attention of any able practitioner, we should have been long since undeceived in this respect. On the coast of Greenland, the mountainous parts of Wales and Scotland, where melted snow must be continually passing into their rivers and streams, the disease is not known, though it is common in Derbyshire, and some other parts of England. Rungpore is about one hundred miles from the foot of the hills, and much farther from the snow, yet the disease is as frequent there as in Boutan. In Thibet, where snow is never out of view, and the principal source of all their rivers and streams, the disease is not to be met with; but what puts the matter past a doubt, is the frequency of the disease on the coast of Sumatra, where snow is never to be found. On finding the vegetable productions of Boutan the same as those of the Alps in almost every instance, it occurred to me, that  
the

the disease might arise from an impregnation of the water by these plants, or the soil probably possessing similar qualities, the spontaneous productions of both countries, with very few exceptions, being so nearly alike. It however appears more probable, that the disease is endemial, proceeding from a peculiarity in the air of situations in the vicinity of mountains with such soil and vegetable productions. I am the more inclined to think so, that I have universally found this disease most prevalent amongst the lower class of people, and those who are most exposed to the unguarded influence of the weather, and various changes that take place in the air of such situations. The primary cause in the atmosphere producing this effect is, perhaps, not more inexplicable than what we meet with in the low-lands of Essex and fens of Lincolnshire. An accurate analysis of the water used in common by the natives, where this disease is more or less frequent, and where it is not known in similar exposures, might throw some light on this subject.

This very extraordinary disease has been little attended to, from obvious reasons; it is unaccompanied with pain, seldom fatal, and generally confined to the poorer sort of people. The tumor is unsightly, and grows to a troublesome size, being often as large as a person's head. It is certainly not exaggerating to say, that one in six of the Rungpore district, and country of Boutan, has the disease.

As those who labour most, and are the least protected from the changes of weather, are most subject to the disease, we universally find it in Boutan more common with the women than men. It generally appears in Boutan at the age of thirteen or fourteen, and in Bengal at the age of eleven or twelve; so that in both countries the disease shows itself about the age

of puberty. I do not believe this disease has ever been removed, though a mercurial course seemed to check its progress, but did not prevent its advance after intermitting the use of mercury. An attention to the primary cause will first lead to a proper method of treating the disease; a change of situation for a short while, at that particular period when it appears, might be the means of preventing it.

The people of this happy climate are not exempt from the venereal disease, which seems to rage with unremitting fury in all climates, and proves the greatest scourge to the human race. It has been long a matter of doubt, whether this disease has ever been cured by any other specific than mercury and its different preparations. In defence of the opinion of other specifics being in use, it has always been urged, that the disease is frequent in many parts of the world, where it could not be supposed that they were acquainted with quicksilver, and the proper method of preparing it as a medicine. I must own, that I expected to have been able to have added one other specific for this disease to our list in the *Materia medica*, being informed that the disease was common, and their method of treating it successful; nor could I allow myself to think they were acquainted with the method of preparing quicksilver, so as to render it a safe and efficacious medicine. In this, however, I was mistaken.

The disease seems in this country to make a more rapid progress, and rage with more violence, than in any other. This is to be accounted for from the grossness of their food and little attention to cleanliness.

There is one preparation of mercury in common use with them, and made after the following manner. A portion of alum, nitre, vermilion, and quicksilver, are placed in the bottom of an earthen

earthen pot, with a smaller one inverted put over the materials, and well luted to the bottom of the larger pot. Over the small one, and within the large one, the fuel is placed, and the fire continued for about forty minutes. A certain quantity of fuel, carefully weighed out, is what regulates them with respect to the degree of heat, as they cannot see the materials during the operation. When the vessel is cool, the small inverted pot is taken off, and the materials collected for use. I attended the whole of the process, and examined the materials afterwards. The quicksilver had been acted on by the other ingredients, deprived of its metallic form, and rendered a safe and efficacious remedy.

A knowledge of chemistry has taught us a more certain method of rendering this valuable medicine active and efficacious; yet we find this preparation answering every good purpose, and by their guarded manner of exhibiting it perfectly safe. This powder is the basis of their pill, and often used in external application. The whole, when intimately mixed, formed a reddish powder, and was made into the form of pills by the addition of a plum or date. Two or three pills taken twice a day generally bring on, about the fourth or fifth day, a spitting, which is encouraged by continuing the use of the pills for a day or two longer. As the salivation advances, they put a stick across the patient's mouth, in the form of a gag, and make it fast behind. This, they say, is done to promote the spitting, and prevent the loss of their teeth. They keep up the salivation for ten or twelve days, during which time the patient is nourished with congee and other liquids. Part of this powder is often used externally by diffusing it in warm water, and washing sores and buboes. They disperse buboes frequently by poultices of turnip tops, in which they always  
put.

put vermillion, and sometimes musk. Nitre, as a cooler, is very much used internally by them in this disease, and they strictly enjoin warmth and confinement during the slightest mercurial course. Buboes advanced to suppuration are opened by a lancet, with a large incision, which they do not allow to close before the hardness and tumor are gone. In short, I found very little room for improving their practice in this disease. I introduced the method of killing quicksilver with honey, gave them an opportunity of seeing it done, and had the satisfaction of finding it successfully used by themselves before we left the country.

This happy climate presents us with but little variety in their diseases. Coughs, colds, and rheumatism, are more frequent here than in Bengal. Fevers generally arise here from a temporary cause, are easily removed, and seldom prove fatal. The liver disease is occasionally to be met with, and complaints in the bowels are not unfrequent; but the grossness of their food, and uncleanness of their persons, would in any other climate be the source of constant disease and sickness. They are ignorant (as we were, not many years ago) of the proper method of treating diseases of the liver and other viscera; this is, I believe, the cause of the most obstinate and fatal disease to be met with in the country, I mean the dropsy. As the Rajah had ever been desirous of my aid and advice, and had directed his doctors to attend to my private instructions and practice, I endeavoured to introduce a more judicious method of treating those diseases by mercurial preparations. I had an opportunity of proving the advantage of this plan to their conviction in several instances, and of seeing them initiated in the practice.

The Rajah favoured me with above seventy specimens of the medicines in use with them. They have many sorts of stones and  
petrifications



petrifications saponaceous to the touch, which are employed as an external application in swellings and pains of the joints. They often remove such complaints, and violent head-achs, by fumi-gating the part affected with aromatic plants and flowers. They do not seek for any other means of information respecting the state of a patient than that of feeling the pulse; and they confidently say, that the seat of pain and disease is easily to be discovered, not so much from the frequency of the pulse as its vibratory motion. They feel the pulse at the wrist with their three fore-fingers, first of the right, and then of the left hand; after pressing more or less on the artery, and occasionally removing one or two of the fingers, they determine what the disease is. They do not eat any thing the day on which they take physic, but endeavour to make up the loss afterwards by eating more freely than before, and using such medicines as they think will occasion costiveness.

The many simples in use with them are from the vegetable kingdom, collected chiefly in Boutan. They are in general inoffensive and very mild in their operation. Carminatives and aromatics are given in coughs, colds, and affections of the breast. The centaury, coriander, carraway, and cinnamon, are of this sort. This last is with them the bark of the root of that species of *Laurus* formerly mentioned as a native of this country. The bark from the root is in this plant the only part which partakes of the cinnamon taste; and I doubt very much if it could be distinguished by the best judges from what we call the true cinnamon. The bark, leaves, berries, and stalks of many shrubs and trees, are in use with them, all in decoction. Some have much of the astringent bitter taste of our most valuable medicines, and are generally employed here with the same view, to strengthen the powers of digestion, and mend the

the general habit. Their principal purgative medicines are brought by the Chinese to Lassa. They had not any medicine that operated as a vomit, till I gave the Rajah some ipecacuanha, who made the first experiment with it on himself.

In bleeding they have a great opinion of drawing the blood from a particular part. For head-achs they bleed in the neck; for pains in the arm and shoulder, in the cephalic vein; and of the breast or side, in the median; and if in the belly, they bleed in the basilic vein. They think pains of the lower extremity are best removed by bleeding in the ankle. They have a great prejudice against bleeding in cold weather; nor is any urgency or violent symptom thought at that time a sufficient reason for doing it.

They have their lucky and unlucky days for operating or taking any medicine; but I have known them get the better of this prejudice, and be prevailed on.

Cupping is much practised by them; a horn, about the size of a cupping glass, is applied to the part, and by a small aperture at the other end they extract the air with their mouth. The part is afterwards scarified with a lancet. This is often done on the back; and in pain and swelling of the knee it is held as a sovereign remedy. I have often admired their dexterity in operating with bad instruments. Mr. HAMILTON gave them some lancets, and they have since endeavoured, with some success, to make them of that form. They were very thankful for the few I could spare them. In fevers they use the Kuthullega nut, well known in Bengal as an efficacious medicine. They endeavour to cure the dropsy by external applications, and giving a compounded medicine made up of above thirty different ingredients: they seldom or never succeed in effecting a cure of this disease.

I explained to the Rajah the operation of tapping, and shewed him the instrument with which it was done. He very earnestly expressed a desire that I should perform the operation, and wished much for a proper subject; such a one did not occur while I remained, and perhaps it was as well both for the Rajah's patients and my own credit; for after having seen it once done, he would not have hesitated about a repetition of the operation. Gravelish complaints and the stone in the bladder are, I believe, diseases unknown here.

The small-pox, when it appears among them, is a disease that strikes them with too much terror and consternation to admit of their treating it properly. Their attention is not employed in saving the lives of the infected, but in preserving themselves from the disease. All communication with the infected is strictly forbidden, even at the risk of their being starved, and the house or village is afterwards erased. A promiscuous and free intercourse with their neighbours not being allowed, the disease is very seldom to be met with, and its progress always checked by the vigilance and terror of the natives. Few in the country have had the disease. Inoculation, if ever introduced, must be very general to prevent the devastation that would be made by the infection in the natural way; and where there could not be any choice in the subject fit to receive the disease, many must fall a sacrifice to it. The present Rajah of Thibet was inoculated, with some of his followers, when in China with the late Tishoo Lama.

The hot bath is used in many disorders, particularly in complaints of the bowels and cutaneous eruptions. The hot wells of Thibet are resorted to by thousands. In Boutan they substitute water warmed by hot stones thrown into it.

In Thibet the natives are more subject to fore eyes and blindness than in Boutan. The high winds, sandy soil, and glare from the reflection of the sun, both from the snow and sand, account for this.

I have dwelt long on this subject, because I think the knowledge and observations of these people on the diseases of their country, with their medical practice, keep pace with a refinement and state of civilization, which struck me with wonder, and no doubt will give rise to much curious speculation, when known to be the manners of a people holding so little intercourse with what we term civilized nations.

Dec. 1. Left Tishoolumbo, and found the cold increase every day as we advanced to the southward, most of the running waters frozen, and the pools covered with ice strong enough to carry. Our thermometer having only the scale as low as  $16^{\circ}$ , we could not precisely determine the degree of cold, the quicksilver being under that every morning. The frost is certainly never so intense in Great-Britain. On our return to the lakes the 14th, we found them deserted by the water fowl, and were informed that they had been one solid piece of ice since the 10th of November. Here we resumed our amusement of skating, to the great astonishment of the natives and Bengal servants.

On the 17th we re-entered Boutan, and in six days more arrived at Pudukha by Paraghon. No snow or frost to be met with in Boutan, except towards the tops of their highest mountains; the thermometer rising to  $36^{\circ}$  in the morning, and  $48^{\circ}$  at noon.

Took leave of the Debe Rajah, and on the 12th arrived at Buxaduar.

Calcutta, Feb. 17, 1784.

AS Lac is the produce of, and a staple article of commerce in Assam, a country bordering on and much connected with Thibet, some account of it may not be an improper supplement to the above remarks.

Lac is, strictly speaking, neither a gummy nor resinous substance, though it has some properties in common to both. Gums are soluble in water, and resins in spirits; lac admits of a very difficult union with either, without the mediation of some other agent.

Lac is known in Europe by the different appellations of stick lac, seed lac, and shell lac. The first is the lac in pretty considerable lumps, with much of the woody parts of the branches on which it is formed adhering to it. Seed lac is only the stick lac broke into small pieces, garbled, and appearing in a granulated form. Shell lac is the purified lac, by a very simple process to be mentioned afterward.

Many vague and unauthenticated reports concerning lac have reached the public; and though amongst the multiplicity of accounts the true history of this substance has been nearly hit on, little credit is given in Europe to any description of it hitherto published. My observations, as far as they go, are the result of what I have seen, from the lac on the tree, the progress of the insect now in my custody, and the information of a gentleman residing at Goalpara on the borders of Assam, who is perfectly versant in the method of breeding the insect, inviting it to the tree, collecting the lac from the branches, and forming it into shell lac, in which state much of it is received from Assam, and exported to Europe for various great and useful purposes. The tree on which this fly most commonly generates is known in Bengal by the name of the *Biber* tree, and is a species of

the Rhamnus. The fly is nourished by the tree, and there deposits its eggs, which nature has provided it with the means of defending from external injury by a collection of this lac, evidently serving the twofold purpose of a nidus and covering to the ovum and insect in its first stage, and food for the maggot in its more advanced state. The lac is formed into complete cells, finished with as much regularity and art as a honey-comb, but differently arranged. The flies are invited to deposit their eggs on the branches of the tree, by besmearing them with some of the fresh lac steeped in water, which attracts the fly, and gives a better and larger crop.

The lac is collected twice a year, in the months of February and August.

I have examined the egg of the fly with a very good microscope; it is of a very pure red, perfectly transparent, except in the centre, where there were evident marks of the embryo forming, and opaque ramifications passing off from the body of it. The egg is perfectly oval, and about the size of an ant's egg. The maggot is about the one-eighth of an inch long, formed of many rings (ten or twelve) with a small red head; when seen with a microscope, the parts of the head were easily distinguished, with six small specks on the breast, somewhat projecting, which seemed to be the incipient formation of the feet. This maggot is now in my custody, in the form of a nymph or crysalis, its annular coat forming a strong covering, from which it should issue forth a fly. I have never seen the fly, and cannot therefore describe it more fully, or determine its genus and species. I am promised a drawing of the insect in its different stages, and shall be able soon to add to a botanical description of the plant a drawing of the branch, with the different parts of fructification and lac on it.

The gentleman to whom I owe part of my information terms the lac the excrement of the insect. On a more minute investigation, however, we may not find it more so than the wax or honey of the bee, or silk of the silk-worm. Nature has provided most insects with the means of secreting a substance which generally answers the twofold purpose of defending the embryo, and supplying nourishment to the insect from the time of its animation till able to wander abroad in quest of food. The fresh lac contains within its cells a liquid, sweetish to the taste, and of a fine red colour, miscible in water. The natives of Assam use it as a dye, and cotton dipped in this liquid makes afterwards a very good red ink.

The simple operation of purifying lac is practised as follows. It is broken into small pieces, and picked from the branches and sticks, when it is put into a sort of canvas bag of about four feet long, and not above six inches in circumference. Two of these bags are in constant use, and each of them held by two men. The bag is placed over a fire, and frequently turned till the lac is liquid enough to pass through its pores, when it is taken off the fire, and squeezed by two men in different directions, dragging it along the convex part of a plantain-tree prepared for the purpose; while this is doing, the other bag is heating, to be treated in the same way. The mucilaginous and smooth surface of the plantain-tree seems peculiarly well adapted for preventing the adhesion of the heated lac, and giving it the form which enhances its value so much. The degree of pressure on the plantain-tree regulates the thickness of the shell, and the quality of the bag determines its fineness and transparency. They have learned of late, that the lac which is thicker in the shell than it used to be, is most prized in Europe. Assam furnishes us with the greatest quantity

tity of lac in use; and it may not be generally known, that the tree on which they produce the best and largest quantity of lac is not uncommon in Bengal, and might be employed in propagating the fly, and cultivating the lac, to great advantage. The small quantity of lac collected in these provinces affords a precarious and uncertain crop, because not attended to. Some attention at particular seasons is necessary to invite the fly to the tree; and collecting the whole of the lac with too great an avidity, where the insect is not very generally to be met with, may annihilate the breed.

The best method of cultivating the tree, and preserving the insect, being properly understood in Bengal, would secure to the Cofs possessions the benefit arising from the sale of a lucrative article, in great demand and of extensive use.

Stages and distance from Rungpore to Taffesudon and Tiffolumboo, in computed coffes and miles, two miles to a cof.

			Coffes.	M. F.
1783,	From Rungpore to			
May 6.	Calamaty Plains	- -	5½	11 0
8.	To Mongulhaut	- -	5½	11 4
9.	To Belladinga	- -	7	14 0
10.	To Bahar	- - -	4	8 0
11.	To Chichacotta in Boutan	-	13	26 0
12.	To Buxaduar	- -	12	24 0
22.	To Joogagoo	- -	5	10 0
23.	To Murishong	- -	5	10 0
25.	To Chooka	- -	9	18 0
			<hr/>	<hr/>
			66	132 4




*Productions of Boutan and Thibet.*

111

			Coffes.	M. F.
		Brought over	66	132 4
1783, May 26.	To Punuka	- -	7	14 0
	27. To Chepta	- -	5	10 0
	29. To Pagha	- -	5	10 0
	30. To Numloo	- -	4	8 0
	31. To Wanakha	- -	4	8 0
June 1.	To Taffesudon, capital of Boutan		3	6 0
			<hr/>	<hr/>
			94	188 4
Sept. 8.	To Pimitung	- -	7	14 0
	9. To Paraghon	- -	6	12 0
	11. To Dukaigun	- -	4	8 0
	12. To Sanha	- -	5	10 0
	13. To a tent on Thibet ground	-	8	16 0
	14. To Chichakumboo, Thibet	-	4	8 0
	15. To Duina	- -	10	20 0
	16. To Chalu	- -	15	30 0
	17. To Simadar	- -	9	18 0
	18. To Selu	- - -	17	34 0
	19. To Takui	- -	9	18 0
	20. To Dequini	- -	14	28 0
	21. To Sehundi	- -	15	30 0
	22. To Tiffolumboo, capital of Thibet		7	14 0
			<hr/>	<hr/>
			224	448 4
			<hr/>	<hr/>





A  
METEOROLOGICAL JOURNAL

KEPT AT THE APARTMENTS OF

THE ROYAL SOCIETY,

BY ORDER OF THE

PRESIDENT AND COUNCIL.

VOL. LXXIX.

Q

METEOROLOGICAL JOURNAL

for January 1788.

1788	Time.		Therm	Therm.	Barom.	Rain.	Winds.		Weather.
	H.M.		without	within.	Inches.	Inch.	Points.	Str.	
Jan. 1	8	0	43	49	29,97		SSE	1	Cloudy.
	2	0	40	51	29,91		S	1	Fine.
2	8	0	48	49	29,58	0,040	S	2	Cloudy.
	2	0	48	52	29,24		S	2	Rainy.
3	8	0	42	50	28,97	0,235	S	2	Fair.
	2	0	46	52	28,89		S	2	Rainy.
4	8	0	43	51	29,13	0,024	SW by W	1	Fair.
	2	0	44	56	29,23		WSW	1	Fine.
5	8	0	37	52	29,35		SW	1	Fine.
	2	0	40	56	29,32		SW	1	Fine.
6	8	0	42	53	29,16		SW	1	Cloudy.
	2	0	42	56	29,37		SW	1	Cloudy.
7	8	0	37	52	29,66		WSW	1	Cloudy.
	2	0	41	54	29,78		W	1	Cloudy.
8	8	0	39	52	29,96		ENE	1	Cloudy.
	2	0	43	53	30,01		ENE	1	Cloudy.
9	8	0	40	51	30,18	0,020	E	2	Rainy.
	2	0	40	52	30,17		E	2	Cloudy.
10	8	0	37	51	30,21		ENE	1	Cloudy.
	2	0	40	53	30,21		NE	1	Cloudy.
11	8	0	37	51	30,32		NE	1	Cloudy.
	2	0	39	53	30,36		NE	1	Cloudy.
12	8	0	36	52	30,38		W	1	Foggy.
	2	0	39	53	30,36		WNW	1	Fine.
13	8	0	39	52	30,21		WNW	1	Cloudy.
	2	0	41	53	30,11		WNW	1	Cloudy.
14	8	0	36	52	29,98		W	1	Fine.
	2	0	41	53	30,19		W	2	Cloudy.
15	8	0	27	50	30,48		WNW	1	Fine.
	2	0	33	52	30,56		WNW	1	Fine.
16	8	0	26	50	30,68		WNW	1	Fine.
	2	0	36	52	30,70		WNW	1	Fine.

METEOROLOGICAL JOURNAL

for January 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
Jan. 17	8	0	33	50	30,66		W	1	Foggy.
	2	0	40	53	30,63		W	1	Cloudy.
18	8	0	36	50	30,23		W	1	Cloudy.
	2	0	45	52	29,98		W	1	Cloudy.
19	8	0	37	50	29,68	0,060	WSW	2	Cloudy.
	2	0	43	53	29,85		WSW	2	Cloudy.
20	8	0	34	51	30,33		NW	1	Cloudy.
	2	0	40	53	30,38		WNW	1	Cloudy.
21	8	0	39	52	30,33		WNW	1	Cloudy.
	2	0	44	54	30,24		WSW	1	Cloudy.
22	8	0	37	52	29,97		W	1	Fine.
	2	0	44	54	30,00		WSW	2	Cloudy.
23	8	0	39	52	30,15		WNW	2	Cloudy.
	2	0	44	54	30,07		WNW	2	Cloudy.
24	8	0	44	52	29,86		W	2	Cloudy.
	2	0	48	54	29,84		W	2	Cloudy.
25	8	0	43	54	29,90	0,020	W	1	Cloudy.
	2	0	45	56	30,02		W	1	Fine.
26	8	0	39	55	30,06		W	1	Fine.
	2	0	46	55	29,98		WSW	1	Cloudy.
27	8	0	42	54	29,97	0,040	S by W	1	Cloudy.
	2	0	45	56	30,02		S	1	Cloudy.
28	8	0	38	54	30,27		SW	1	Cloudy.
	2	0	39	56	30,31		WSW	1	Cloudy.
29	8	0	32	53	30,33		WNW	1	Foggy.
	2	0	35	53	30,35		N	1	Foggy.
30	8	0	33	52	30,38		NE	1	Cloudy.
	2	0	38	53	30,41		NE	1	Cloudy.
31	8	0	36	51	30,30		E	1	Cloudy.
	2	0	38	54	30,21		ESE	1	Cloudy.

METEOROLOGICAL JOURNAL

for February 1788.

1788	Time.		Therm.	Therm	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
			Deg.	Deg.					
Feb. 1	8	0	32	50	30,00		ESE	1	Cloudy.
	2	0	33	50	29,87		ESE	1	Cloudy.
2	8	0	29	50	29,66		N	1	Cloudy.
	2	0	37	51	29,55		NW	2	Cloudy.
3	8	0	43	50	29,26	0,076	WSW	2	Cloudy.
	2	0	46	52	29,40		WSW	2	Cloudy.
4	8	0	35	50	29,73		SW	1	Foggy.
	2	0	42	53	29,77		SW	1	Fine.
5	8	0	40	50	29,70	0,112	E	1	Cloudy.
	2	0	46	52	29,71		E	1	Cloudy.
6	8	0	35	51	30,15		W	1	Foggy.
	2	0	45	53	30,18		W	1	Cloudy.
7	8	0	37	52	30,21		NE	1	Foggy.
	2	0	46	54	30,21		NE	1	Cloudy.
8	8	0	39	53	30,13		NE	1	Cloudy.
	2	0	38	53	29,98		NE	1	Rainy.
9	8	0	33	51	29,80	0,067	NNE	1	Cloudy.
	2	0	33	53	29,73		NNE	1	Cloudy.
10	8	0	33	51	29,73		N	1	Cloudy.
	2	0	35	51	29,77		NW	1	Cloudy.
11	8	0	38	50	29,89		SE	1	Cloudy.
	2	0	45	54	29,98		SSW	1	Cloudy.
12	8	0	40	51	30,21		SW	1	Foggy.
	2	0	48	52	30,16		SW	1	Cloudy.
13	8	0	43	52	29,94	0,040	SW	1	Small rain.
	2	0	49	54	29,97		W	1	Fair.
14	8	0	40	52	30,12		W	1	Cloudy.
	2	0	49	54	30,06		W	1	Cloudy.
15	8	0	44	53	29,86	0,146	WNW	1	Cloudy.
	2	0	50	55	29,85		NW	1	Fine.
16	8	0	41	54	29,76	0,053	WSW	1	Rain.
	2	0	47	56	29,65		WSW	1	Fair.

METEOROLOGICAL JOURNAL

for February 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
Feb. 17	8	0	46	56	29,71		W	I	Cloudy.
	2	0	47	56	29,75		SSW	I	Cloudy.
18	8	0	33	53	29,83	0,132	E	I	Cloudy.
	2	0	39	53	29,79		ESE	I	Fair.
19	8	0	39	52	29,45		E	I	Cloudy.
	2	0	44	54	29,33		SSE	I	Fine.
20	8	0	39	51	29,02	0,032	E	I	Cloudy.
	2	0	42	54	28,91		E	I	Cloudy.
21	8	0	41	52	28,71	0,235	E	I	Rainy.
	2	0	47	54	28,65		SSE	I	Rainy.
22	8	0	44	53	28,76	0,325	SW	I	Cloudy.
	2	0	47	54	28,94		WSW	I	Cloudy.
23	8	0	41	53	29,05		SW	I	Fine.
	2	0	48	54	29,08		SSW	I	Fair.
24	8	0	42	53	28,98		ESE	I	Cloudy.
	2	0	44	54	28,91		NE	I	Rain.
25	8	0	39	53	29,33	0,168	NW	I	Cloudy.
	2	0	43	54	29,50		WNW	I	Cloudy.
26	8	0	30	52	29,62		WNW	I	Fine.
	2	0	45	54	29,56		W	I	Fair.
27	8	0	41	52	29,31	0,038	E	I	Rain.
	2	0	44	54	29,35		E	I	Cloudy.
28	8	0	41	52	29,50		E	I	Cloudy.
	2	0	44	53	29,47		E	I	Cloudy.
29	8	0	42	53	29,28	0,037	SE	I	Cloudy.
	2	0	48	54	29,20		SSE	I	Cloudy.

METEOROLOGICAL JOURNAL

for March 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
Mar. 1	7	0	35	53	29,43		SW by S	1	Fine.
	2	0	45	54	29,48		SSW	1	Fair.
2	7	0	41	52	29,52		E	1	Cloudy.
	2	0	42	53	29,63		ESE	2	Cloudy.
3	7	0	35	51	29,99		E	2	Cloudy.
	2	0	41	53	30,08		E	2	Fair.
4	7	0	36	51	30,03		NW	1	Foggy.
	2	0	42	53	29,92		W	1	Fair.
5	7	0	33	51	29,97		NNW	1	Cloudy.
	2	0	41	53	29,88		W	1	Fair.
6	7	0	39	51	29,34	0,075	WSW	1	Fair.
	2	0	43	53	29,45		WNW	2	Cloudy.
7	7	0	34	50	29,43		NW	1	Fair.
	2	0	40	51	29,51		NW	1	Fair.
8	7	0	28	48	29,52		NE	1	Fair.
	2	0	39	50	29,54		NE	1	Fair.
9	7	0	29	48	29,59		NE	2	Fair.
	2	0	39	49	29,64		ENE	2	Fair.
10	7	0	30	47	29,89		ENE	1	Fine.
	2	0	38	50	29,97		ENE	1	Fair.
11	7	0	28	46	30,06		ENE	1	Fine.
	2	0	38	50	30,08		ENE	1	Fine.
12	7	0	29	47	30,03		E	1	Fine.
	2	0	39	51	30,00		ENE	1	Fine.
13	7	0	29	47	29,84		ESE	1	Cloudy.
	2	0	32	48	29,73		ESE	1	Cloudy.
14	7	0	31	46	29,52		E	1	Fine.
	2	0	39	49	29,44		ESE	1	Fair.
15	7	0	36	46	29,44		ESE	1	Cloudy.
	2	0	40	49	29,46		ESE	1	Cloudy.
16	7	0	38	47	29,46	0,120	ESE	1	Cloudy.
	2	0	37	47	29,46		ESE	2	Cloudy.



METEOROLOGICAL JOURNAL

for March 1788.

1788	Time		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.			Points.	Str.	
			Deg.	Deg.	inches.	Inch.			
Mar. 17	7	0	33	46	29,50		ESE	2	Cloudy.
	2	0	33	47	29,52		ESE	2	Cloudy.
18	7	0	33	46	29,68		ESE	2	Cloudy.
	2	0	39	49	29,75		ESE	2	Cloudy.
19	7	0	36	47	30,00		SE by E	1	Fair.
	2	0	42	49	30,03		SE by E	1	Fair.
20	7	0	41	47	29,82		SE	1	Cloudy.
	2	0	50	50	29,74		SW	1	Fair.
21	7	0	44	49	29,76	0,070	SSW	1	Fair.
	2	0	52	53	29,72		SW	1	Fair.
22	7	0	43	50	29,57		SSW	1	Cloudy.
	2	0	47	53	29,48		SSE	1	Cloudy.
23	7	0	39	50	29,32	0,071	N	2	Cloudy.
	2	0	44	53	29,47		N	2	Cloudy.
24	7	0	40	49	29,61		SW	2	Fair.
	2	0	51	53	29,59		SSW	2	Hazy.
25	7	0	46	51	29,52		SSW	2	Cloudy.
	2	0	52	54	29,58		SSW	2	Fair.
26	7	0	40	52	29,60		SSE	2	Fine.
	2	0	53	55	29,52		SSE	2	Fine.
27	7	0	42	54	29,52		S	1	Fine.
	2	0	51	57	29,44		SSE	2	Fine.
28	7	0	46	55	29,45		SSW	2	Fair.
	2	0	56	57	29,57		SW	2	Fair.
29	7	0	46	55	29,92		WSW	1	Cloudy.
	2	0	56	58	30,01		W	1	Fair.
30	7	0	50	56	29,95		WSW	1	Cloudy.
	2	0	59	58	29,90		SW	2	Fine.
31	7	0	48	58	29,63		W	2	Fine.
	2	0	49	59	29,68		W	2	Fair.

METEOROLOGICAL JOURNAL

for April 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
April 1	7	0	47	56	29,72	0,390	W	2	Rainy.
	2	0	55	60	29,74		WNW	2	Fair.
2	7	0	46	57	30,09	0,020	W	2	Fair.
	2	0	58	60	30,02		W	2	Fair.
3	7	0	47	58	29,72	0,020	W	2	Fair.
	2	0	54	60	29,50		W	2	Fair.
4	7	0	43	57	29,67	0,020	NW	2	Cloudy.
	2	0	40	58	29,75		NW by N	2	Hail.
5	7	0	40	55	29,97	0,020	NNW	2	Cloudy.
	2	0	41	57	30,10		N	2	Cloudy.
6	7	0	42	55	30,22	0,020	W	1	Fair.
	2	0	51	56	30,23		W	1	Cloudy.
7	7	0	48	55	30,27	0,020	W	1	Cloudy.
	2	0	51	56	30,30		WNW	1	Cloudy.
8	7	0	51	56	30,39	0,020	WNW	1	Cloudy.
	2	0	59	59	30,42		WNW	1	Cloudy.
9	7	0	48	56	30,48	0,020	WNW	1	Fair.
	2	0	59	59	30,47		NNW	1	Cloudy.
10	7	0	51	57	30,40	0,020	E	1	Cloudy.
	2	0	59	59	30,30		ESE	1	Fine.
11	7	0	49	57	30,16	0,020	E	1	Fine.
	2	0	58	59	30,07		SE	1	Fine.
12	7	0	48	57	29,97	0,020	SSW	1	Fine.
	2	0	56	58	30,09		SSW	1	Fine.
13	7	0	45	57	30,28	0,020	SW	1	Fine.
	2	0	59	61	30,25		SW by W	1	Fine.
14	7	0	45	58	0,06	0,020	W	1	Fine.
	2	0	53	61	30,00		W	1	Rainy.
15	7	0	43	57	30,03	0,097	W	1	Fine.
	2	0	51	59	30,00		W	1	Fair.
16	7	0	45	57	30,11	0,097	W	1	Fair.
	2	0	53	59	30,11		W	1	Fair.

METEOROLOGICAL JOURNAL

for April 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
Apr. 17	7	0	48	58	30,09		W	1	Cloudy.
	2	0	55	60	30,09		W	1	Cloudy.
18	7	0	50	59	30,21		W	1	Fair.
	2	0	62	61	30,26		N	1	Fair.
19	7	0	54	60	30,27		E	1	Fair.
	2	0	65	61	30,24		SE	1	Fine.
20	7	0	52	60	30,03		SSE	1	Fine.
	2	0	67	61	29,88		SSW	1	Fine.
21	7	0	45	59	29,76	0,100	SW	1	Fine.
	2	0	56	60	29,70		WSW	2	Fine.
22	7	0	46	59	29,95		SW	2	Fine.
	2	0	55	61	29,90		SW	1	Cloudy.
23	7	0	53	60	29,91		WSW	1	Cloudy.
	2	0	63	61	29,89		W	1	Cloudy.
24	7	0	47	60	29,95		W	1	Fine.
	2	0	56	60	30,01		WNW	2	Fine.
25	7	0	49	58	29,97		WNW	2	Fair.
	2	0	57	61	29,96		WNW	2	Fair.
26	7	0	54	59	30,15		W	1	Cloudy.
	2	0	61	60	30,18		W by S	2	Fair.
27	7	0	48	58	30,28		SW	1	Fine.
	2	0	63	61	30,28		WSW	1	Fine.
28	7	0	47	59	30,32		W	1	Fine.
	2	0	66	60	30,28		SE	1	Fine.
29	7	0	55	60	30,27		ESE	1	Fine.
	2	0	68	63	30,22		ESE	1	Fine.
30	7	0	55	60	30,22		ESE	1	Fine.
	2	0	68	63	30,20		ESE	1	Fine.

## METEOROLOGICAL JOURNAL

for May 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
May 1	7	0	54	60	30,18		ESE	1	Fine.
	2	0	69	63	30,16		ESE	1	Fine.
2	7	0	56	61	30,14		NE	1	Fine.
	2	0	71	64	30,14		E	2	Fine.
3	7	0	50	60	30,34		ENE	2	Cloudy.
	2	0	57	62	30,32		ENE	2	Fair.
4	7	0	50	60	30,29		NE	1	Cloudy.
	2	0	55	61	30,27		NNE	2	Fair.
5	7	0	51	60	30,22		ESE	2	Fine.
	2	0	64	61	30,14		ESE	2	Fine.
6	7	0	54	61	30,02		ESE	1	Fine.
	2	0	68	62	29,98		ESE	1	Fine.
7	7	0	57	61	29,88		SSE	1	Fair.
	2	0	64	63	29,86		SW by S	1	Cloudy.
8	7	0	54	61	29,96	0,025	SSW	1	Fine.
	2	0	66	64	29,93		SSW	1	Fine.
9	7	0	53	62	29,91		SSW	1	Cloudy.
	2	0	64	64	29,86		SW	2	Fair.
10	7	0	53	61	29,93	0,080	WSW	2	Fine.
	2	0	62	63	29,98		W	1	Fine.
11	7	0	49	60	30,18		WNW	1	Cloudy.
	2	0	62	62	30,17		WSW	1	Fair.
12	7	0	55	60	30,27		WSW	1	Fair.
	2	0	64	62	30,26		SSE	1	Fair.
13	7	0	51	60	30,31		E by S	1	Fine.
	2	0	59	62	30,28		ESE	1	Fine.
14	7	0	52	60	30,18		NE	1	Hazy.
	2	0	56	59	30,16		E	2	Cloudy.
15	7	0	52	59	30,12		ENE	2	Fair.
	2	0	58	60	30,05		ENE	2	Cloudy.
16	7	0	50	59	29,85	0,030	E	2	Rainy.
	2	0	61	60	29,92		E	2	Fine.

METEOROLOGICAL JOURNAL

for May 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
May 17	7	0	54	60	29,96	0,031	E	2	Fair.
	2	0	68	61	29,91		ESE	2	Fair.
18	7	0	55	59	29,87		E	1	Cloudy.
	2	0	59	61	29,87				Rainy.
19	7	0	54	60	29,89	0,260	NE	1	Cloudy.
	2	0	61	60	29,93		NNE	1	Cloudy.
20	7	0	55	60	30,13		NNE	1	Fine.
	2	0	70	61	30,15		N	1	Fine.
21	7	0	55	60	30,28		NW	1	Fine.
	2	0	70	61	30,28		WSW	1	Fine.
22	7	0	56	61	30,28		SW	1	Fine.
	2	0	68	63	30,27		WSW	1	Fine.
23	7	0	57	63	30,20		SW	1	Cloudy.
	2	0	68	63	30,16		SW	1	Cloudy.
24	7	0	57	63	30,08		SW	1	Fine.
	2	0	75	66	30,02		SSW	1	Fine.
25	7	0	63	65	30,00		S	1	Fine.
	2	0	75	69	30,02		SSW	2	Fine.
26	7	0	63	67	30,00		SE by S	1	Fine.
	2	0	73	72	30,00		SSE	1	Fine.
27	7	0	63	70	29,91		SE	1	Fine.
	2	0	80	73	29,85		SE	1	Fine.
28	7	0	67	70	29,71		E	1	Fine.
	2	0	73	71	29,66		SSE	1	Hazy.
29	7	0	63	70	29,58	0,071	SW	1	Cloudy.
	2	0	61	68	29,67			0	Cloudy.
30	7	0	52	66	29,71		E	1	Cloudy.
	2	0	55	65	29,72		E	1	Cloudy.
31	7	0	52	64	29,88		NNE	1	Cloudy.
	2	0	58	63	29,94		NE	1	Cloudy.

METEOROLOGICAL JOURNAL

for June 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.M.	Deg.	without	within.	Inches.	Inch.	Points.	Str.	
June 1	7	0	52	62	30,02		E	1	Fair.
	2	0	64	63	29,99		E	1	Cloudy.
2	7	0	54	62	29,98		SSE	1	Fine.
	2	0	68	63	29,98		SSE	1	Fine.
3	7	0	56	62	30,02		E	1	Fair.
	2	0	67	63	30,02		E	1	Fine.
4	7	0	57	63	30,00		WNW	1	Fine.
	2	0	62	63	30,04		WNW	1	Cloudy.
5	7	0	52	62	30,22			0	Fine.
	2	0	68	63	30,28		W	1	Fine.
6	7	0	60	63	30,20		W	1	Fine.
	2	0	72	60	30,08		SW	1	Fine.
7	7	0	60	65	30,00		W	1	Fair.
	2	0	72	66	29,97		W	1	Fair.
8	7	0	56	61	30,08	0,035	ESE	1	Fine.
	2	0	62	63	30,10		ESE	1	Fine.
9	7	0	52	62	30,17		ENE	2	Cloudy.
	2	0	64	63	30,19		NE	2	Fine.
10	7	0	55	62	30,21		N	2	Fine.
	2	0	69	64	30,18		ESE	2	Fine.
11	7	0	55	63	30,15		NE	1	Fine.
	2	0	67	64	30,08		NE	1	Fine.
12	7	0	54	63	30,06		NNE	1	Fine.
	2	0	70	66	30,04		NE	1	Fine.
13	7	0	57	63	30,06		NE	2	Fine.
	2	0	70	66	30,06		NNE	2	Fine.
14	7	0	58	64	30,07		NNE	1	Cloudy.
	2	0	67	65	30,06		NNE	1	Cloudy.
15	7	0	59	65	30,05		NNE	1	Fine.
	2	0	72	66	30,03		N	1	Fine.
16	7	0	59	59	30,00	0,120	N	1	Rain.
	2	0	68	66	29,96		N	1	Fine.

METEOROLOGICAL JOURNAL

for June 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
			Deg.	Deg.					
June 17	7	0	60	65	29,99		N	I	Fine.
	2	0	80	67	29,95		N	I	Fine.
18	7	0	65	69	29,87		E	I	Cloudy.
	2	0	70	70	29,84		SSE	I	Cloudy.
19	7	0	63	69	29,81	0,023	SSE	I	Cloudy.
	2	0	66	68	29,87		S	I	Rain.
20	7	0	56	67	29,94	0,060	SE	I	Cloudy.
	2	0	60	67	29,94		SE	I	Rain.
21	7	0	60	66	29,98	0,030	NNE	I	Cloudy.
	2	0	60	67	29,98		N	I	Cloudy.
22	7	0	63	66	29,98		NNW	I	Fair.
	2	0	72	68	29,96		NNW	I	Fair.
23	7	0	62	66	29,80		W	I	Fine.
	2	0	65	67	29,75		SSW	I	Cloudy.
24	7	0	58	66	29,70		SSW	2	Fair.
	2	0	63	66	29,68		SSW	2	Cloudy.
25	7	0	58	66	29,63	0,245	SW	I	Fair.
	2	0	65	67	29,60		SW	I	Cloudy.
26	7	0	58	66	29,61	0,050	WSW	I	Cloudy.
	2	0	63	66	29,57		SW by S	I	Cloudy, heavy rain with thund.
27	7	0	58	65	29,53	2,116	SW by S	I	Fair.
	2	0	71	67	29,49		S	I	Fair.
28	7	0	59	66	29,54	0,285	SSE	I	Rain.
	2	0	63	66	29,54		SE	I	Rain.
29	7	0	57	65	29,73	0,311	SE	I	Cloudy.
	2	0	64	65	29,79		SE	I	Cloudy.
30	7	0	59	64	29,86		W	I	Cloudy.
	2	0	67	65	29,93		W	I	Cloudy.

METEOROLOGICAL JOURNAL

for July 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
July 1	7	0	60	64	30,07		W	2	Cloudy.
	2	0	63	64	30,07		W	2	Cloudy.
2	7	0	61	64	30,07		SW	2	Fair.
	2	0	69	65	30,07		SW	2	Fair.
3	7	0	63	65	30,04		S	1	Fair.
	2	0	73	66	29,93		SSW	1	Fine.
4	7	0	58	65	29,74		SSW	2	Rain.
	2	0	70	66	29,77		SSW	2	Fine.
5	7	0	57	65	29,77	0,040	SSW	1	Fine.
	2	0	67	66	29,73		SSW	1	Fine.
6	7	0	57	64	29,81		SW	1	Fine.
	2	0	65	65	29,89		SW	1	Fair.
7	7	0	56	64	29,85		SSW	2	Fair.
	2	0	63	65	29,84		WSW	2	Rain.
8	7	0	57	65	29,90	0,250	SW	1	Fair.
	2	0	62	65	29,80		SSW	1	Cloudy.
9	7	0	56	64	29,76	0,210	SSW	1	Fair.
	2	0	62	65	29,76		S	1	Cloudy.
10	7	0	61	65	29,81		SSW	2	Cloudy.
	2	0	70	66	29,82		SSW	2	Cloudy.
11	7	0	62	65	29,90		S	2	Cloudy.
	2	0	68	66	29,92		S	1	Cloudy.
12	7	0	64	66	29,93	0,250	S	1	Rain.
	2	0	75	67	29,77		SSW	1	Fair.
13	7	0	61	66	29,73		SSW	2	Cloudy.
	2	0	67	66	29,78		SSW	2	Cloudy.
14	7	0	58	66	29,87	0,132	SSW	1	Fair.
	2	0	64	67	29,89		S	1	Fair.
15	7	0	58	66	29,82	0,111	SSW	2	Rain.
	2	0	67	68	29,77		SSW	2	Cloudy.
16	7	0	57	66	29,75	0,082	SSW	2	Cloudy.
	2	0	64	67	29,78		SSW	2	Cloudy.



METEOROLOGICAL JOURNAL

for July 1788.

1788	Time.		Therm without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
July 17	7	0	57	66	29,83		SW	2	Rain.
	2	0	67	67	29,85		W	1	Fair.
18	7	0	57	66	30,15	0,480	W	1	Fine.
	2	0	71	67	30,16		W	1	Fine.
19	7	0	62	66	30,14		SSW	2	Fine.
	2	0	69	68	30,08		SSW	2	Cloudy.
20	7	0	59	66	30,05		WSW	1	Cloudy.
	2	0	68	67	30,13		W	1	Cloudy.
21	7	0	60	66	30,20	0,065	ESE	1	Fair.
	2	0	67	66	30,22		NW	1	Cloudy.
22	7	0	60	66	30,21		W	1	Fair.
	2	0	71	67	30,20		W	1	Fine.
23	7	0	61	66	30,17		WNW	1	Hazy.
	2	0	71	67	30,11		W	1	Cloudy.
24	7	0	60	66	30,05		W	1	Fine.
	2	0	67	67	30,00		W	1	Cloudy.
25	7	0	60	66	30,11		W	1	Fair.
	2	0	65	67	30,10		W	1	Fair.
26	7	0	55	65	30,15		W	1	Cloudy.
	2	0	67	66	30,15		W	1	Cloudy.
27	7	0	55	65	30,14		W	1	Fine.
	2	0	69	66	30,12		W	1	Cloudy.
28	7	0	60	65	30,10		W	1	Fair.
	2	0	71	66	30,10		W	1	Fine.
29	7	0	59	65	30,11		WNW	1	Fair.
	2	0	72	66	30,10		NW	1	Fine.
30	7	0	62	66	30,16		SE by S	1	Cloudy.
	2	0	76	68	30,17		SW	1	Fine.
31	7	0	60	66	30,22		SW	1	Fine.
	2	0	77	69	30,20		SW	1	Fine.

METEOROLOGICAL JOURNAL

for August 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
			Deg.	Deg.					
Aug. 1	7	0	65	68	30,23		SW	1	Cloudy.
	2	0	72	69	30,29		W	1	Cloudy.
2	7	0	64	68	30,45		N	1	Fine.
	2	0	73	70	30,44		NE	1	Fine.
3	7	0	65	68	30,43		E	1	Fine.
	2	0	76	70	30,43		E	1	Fine.
4	7	0	60	66	30,43		ESE	1	Hazy.
	2	0	77	71	30,37				Fine.
5	7	0	61	67	30,30		NE	1	Hazy.
	2	0	69	69	30,25		N	1	Fair.
6	7	0	57	67	30,21		NNE	1	Fair.
	2	0	64	67	30,18		N	2	Cloudy.
7	7	0	58	66	30,12		NNE	1	Cloudy.
	2	0	62	66	30,12		NNE	1	Cloudy.
8	7	0	58	66	30,19		NNE	1	Cloudy.
	2	0	62	66	30,19		NNE	1	Cloudy.
9	7	0	58	66	30,19		NE	1	Cloudy.
	2	0	66	66	30,18		ENE	1	Cloudy.
10	7	0	58	65	30,14		NE	1	Cloudy.
	2	0	66	66	30,11		NE	1	Cloudy.
11	7	0	58	65	30,08		ENE	1	Cloudy.
	2	0	68	65	30,04		ENE	1	Cloudy.
12	7	0	57	65	29,95			0	Fair.
	2	0	71	66	29,83		SSW	1	Cloudy.
13	7	0	60	66	29,58	0,325	W	1	Cloudy.
	2	0	70	67	29,51		SW	1	Cloudy.
14	7	0	58	66	29,22	0,293	SW	1	Rain.
	2	0	66	66	29,37		SW	2	Rain.
15	7	0	57	65	29,60	0,230	WSW	2	Fine.
	2	0	67	66	29,60		WSW	2	Fair.
16	7	0	57	65	29,74	0,115	WSW	2	Fine.
	2	0	57	65	29,74		SW	1	Rain.

METEOROLOGICAL JOURNAL

for August 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
			Deg.	Deg.					
Aug. 17	7	0	58	65	29,72	0,673	S by W	1	Rain.
	2	0	61	65	29,64		S by W	1	Fair.
18	7	0	57	65	29,80	0,440	WSW	1	Fine.
	2	0	63	65	29,84		WSW	1	Rain.
19	7	0	59	65	29,68	0,130	SSW	2	Cloudy.
	2	0	65	66	29,66		SSW	2	Cloudy.
20	7	0	57	65	29,79	0,080	WSW	1	Fair.
	2	0	71	66	29,85		WSW	1	Fair.
21	7	0	57	66	29,99	0,023	SSW	1	Fair.
	2	0	70	66	29,98		SSW	1	Fair.
22	7	0	59	66	29,76	0,080	SSE	1	Rain.
	2	0	64	66	29,67		S	1	Rain.
23	7	0	59	66	29,67	0,045	SW	1	Cloudy.
	2	0	70	66	29,72		SW	1	Fine.
24	7	0	56	66	29,95		WSW	1	Fine.
	2	0	67	66	29,97		SW	1	Cloudy.
25	7	0	53	65	29,97	0,012	SW	1	Fine.
	2	0	69	66	29,94		SW	1	Fine.
26	7	0	57	65	29,82		SE	1	Cloudy.
	2	0	68	67	29,82		N	1	Fine.
27	7	0	57	65	29,93		W	1	Cloudy.
	2	0	63	65	29,96		W	1	Cloudy.
28	7	0	55	64	30,03		W	2	Cloudy.
	2	0	70	66	30,03		W	1	Fine.
29	7	0	58	65	29,97		SW	1	Cloudy.
	2	0	58	65	29,82		SSW	1	Rain.
30	7	0	55	64	29,82	0,230	W	2	Fine.
	2	0	67	65	29,84		W	1	Cloudy.
31	7	0	57	64	29,80	0,023	SW by S	1	Cloudy.
	2	0	62	64	29,74		SSW	1	Cloudy.

METEOROLOGICAL JOURNAL

for September 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
Sept. 1	7	0	59	64	29,87	0,220	NW	1	Fine.
	2	0	68	65	29,91		W	1	Cloudy.
2	7	0	58	64	29,98	0,905	WSW	1	Cloudy.
	2	0	67	65	29,98		WSW	1	Cloudy.
3	7	0	59	64	30,00	0,905	SW	1	Cloudy.
	2	0	67	65	30,00		SSW	1	Fair.
4	7	0	59	65	29,75	0,905	SE	1	Fine.
	2	0	74	67	29,71		SE	2	Fine.
5	7	0	60	66	29,78	0,905	SSE	1	Fair.
	2	0	70	67	29,80		SE	1	Cloudy.
6	7	0	55	66	29,90	0,905	SW	1	Fair.
	2	0	63	66	29,94		SSW	1	Cloudy.
7	7	0	55	66	30,15	0,905	SSE	1	Fine.
	2	0	72	67	30,17		SSE	1	Fine.
8	7	0	55	66	30,14	0,905	SSW	1	Fine.
	2	0	67	67	30,10		SW	1	Cloudy.
9	7	0	57	65	30,08	0,905	WSW	1	Cloudy.
	2	0	69	66	30,08		W	1	Cloudy.
10	7	0	53	65	30,07	0,905	W	1	Foggy.
	2	0	67	67	30,07		W	1	Fine.
11	7	0	58	66	30,08	0,905	SW	1	Cloudy.
	2	0	63	66	30,16		SSE	1	Fair.
12	7	0	51	65	30,25	0,070	ESE	1	Foggy.
	2	0	67	66	30,18		ESE	1	Fine.
13	7	0	56	65	29,98	0,070	ESE	1	Fine.
	2	0	67	66	29,88		W	1	Cloudy.
14	7	0	54	65	29,94	0,070	W	1	Fine.
	2	0	62	66	30,01		NNE	1	Fine.
15	7	0	50	64	30,12	0,070	NNE	1	Fine.
	2	0	61	64	30,05		E	1	Cloudy.
16	7	0	55	63	29,82	0,070	E	1	Cloudy.
	2	0	67	64	29,79		SSE	1	Fair.

METEOROLOGICAL JOURNAL

for September 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
Sept. 17	7	0	50	63	29,88	0,067	W	1	Fine.
	2	0	62	65	29,87		E	1	Fine.
18	7	0	56	64	29,49	0,105	ESE	2	Rain.
	2	0	62	64	29,50		E	2	Cloudy.
19	7	0	56	63	29,55	0,513	E	1	Rain.
	2	0	60	64	29,60		ESE	1	Cloudy.
20	7	0	54	63	29,57	0,280	ESE	1	Cloudy.
	2	0	60	64	29,47			0	Cloudy.
21	7	0	53	63	29,37	0,300		0	Cloudy.
	2	0	58	63	29,44		W	1	Cloudy.
22	7	0	45	60	29,55	0,128	W	1	Fine.
	2	0	53	60	29,50		SW	2	Cloudy.
23	7	0	52	61	29,75	0,145	W by S	1	Cloudy.
	2	0	60	62	29,74		SW	1	Cloudy.
24	7	0	52	61	29,67	0,302	W	1	Rain.
	2	0	58	61	29,68		SW	1	Cloudy.
25	7	0	45	59	29,81	0,062	W	1	Fine.
	2	0	59	60	29,88		WSW	1	Fair.
26	7	0	57	60	29,83		SSW	2	Cloudy.
	2	0	65	61	29,84		SSW	2	Fair.
27	7	0	52	60	29,95	0,058	SSW	2	Fine.
	2	0	61	62	30,00		SSW	2	Fine.
28	7	0	48	60	30,03		SW	1	Fine.
	2	0	60	62	29,90		SSW	2	Cloudy.
29	7	0	52	61	29,56	0,190	W	2	Fine.
	2	0	57	62	29,61		W	2	Fine.
30	7	0	50	60	29,86		W	2	Fine.
	2	0	56	60	29,92		W	2	Cloudy.

METEOROLOGICAL JOURNAL

for October 1788.

1788	Time.		Therm without	Therm within.	Barom.	Rain.	Winds.		Weather.
	H.M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.		
Oct. 1	7	0	53	59	29,92		W	2	Cloudy.
	2	0	58	60	29,90		W	1	Rain.
2	7	0	59	60	29,94	0,035	W	2	Cloudy.
	2	0	67	62	29,94		W	2	Cloudy.
3	7	0	57	61	29,98		W	2	Cloudy.
	2	0	63	62	30,02		W	2	Cloudy.
4	7	0	56	61	30,08		W	1	Fine.
	2	0	62	62	30,09		W	1	Cloudy.
5	7	0	56	61	30,00		SSW	1	Cloudy.
	2	0	58	62	29,95		SW	2	Cloudy.
6	7	0	55	61	29,70		SW	2	Cloudy.
	2	0	53	61	29,65		SW	2	Rain.
7	7	0	46	59	30,14	0,068	SSW	1	Rain.
	2	0	52	60	30,28		SSE	2	Fine.
8	7	0	43	58	30,55		SE by S	2	Fine.
	2	0	55	58	30,55		NE	2	Cloudy.
9	7	0	50	57	30,50		ENE	2	Fine.
	2	0	57	58	30,47		E	2	Fine.
10	7	0	47	57	30,44		E	2	Fine.
	2	0	59	59	30,44		E	2	Fine.
11	7	0	50	57	30,38		SE	2	Cloudy.
	2	0	59	61	30,35		E	2	Cloudy.
12	7	0	51	59	30,32		NE	2	Cloudy.
	2	0	59	61	30,32		NE	1	Fine.
13	7	0	51	60	30,23		NE	1	Cloudy.
	2	0	58	62	30,26		NE	1	Fine.
14	7	0	51	61	30,25		NE	1	Cloudy.
	2	0	56	62	30,22		NE	1	Cloudy.
15	7	0	42	60	30,05		NE	1	Fine.
	2	0	56	62	29,90		E	1	Fine.
16	7	0	42	60	29,66		NW	1	Foggy.
	2	0	52	60	29,64		W	1	Fair.

METEOROLOGICAL JOURNAL

for October 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	Inch.	Points.	Str.	
			Deg.	Deg.					
Oct. 17	7	0	50	59	29,74		W	I	Foggy.
	2	0	56	61	29,83		W	I	Cloudy.
18	7	0	38	57	30,15		N	I	Fine.
	2	0	50	60	30,22		NNE	I	Fine.
19	7	0	33	57	30,36		NNE	I	Foggy.
	2	0	43	57	30,36		NNE	I	Hazy.
20	7	0	36	54	30,32			0	Foggy.
	2	0	50	59	30,28		SW	I	Fine.
21	7	0	50	56	30,12		SW	I	Cloudy.
	2	0	60	60	30,06		WSW	I	Cloudy.
22	7	0	49	57	30,25		WSW	I	Cloudy.
	2	0	58	60	30,30		WSW	I	Cloudy.
23	7	0	53	58	30,27		WSW	I	Cloudy.
	2	0	60	62	30,20		W	I	Fair.
24	7	0	49	58	30,12		W	I	Cloudy.
	2	0	56	60	30,10		W by N	I	Fine.
25	7	0	44	58	30,11		WNW	I	Fair.
	2	0	53	60	30,15		NNW	I	Cloudy.
26	7	0	41	58	30,19		WNW	I	Fair.
	2	0	54	60	30,15		WNW	I	Fair.
27	7	0	47	58	30,08		WNW	I	Foggy.
	2	0	53	60	30,06		WNW	I	Cloudy.
28	7	0	42	58	30,02		WNW	I	Cloudy.
	2	0	52	60	30,00		WNW	I	Cloudy.
29	7	0	41	58	30,05			0	Foggy.
	2	0	48	60	30,06			0	Foggy.
30	7	0	42	58	30,15		N	I	Hazy.
	2	0	51	59	30,22		NE	I	Cloudy.
31	7	0	45	57	30,43		N	I	Cloudy.
	2	0	52	59	30,47		E	I	Cloudy.

## METEOROLOGICAL JOURNAL

for November 1788.

1788	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without Deg.	within. Deg.	Inches.	Inch.	Points.	Str.	
Nov. 1	7	0	41	57	30,50		E	1	Foggy.
	2	0	51	60	30,43		E	1	Fine.
2	7	0	43	57	30,18		E	1	Foggy.
	2	0	55	61	30,08		S	1	Fine.
3	7	0	55	59	29,92		SSW	2	Cloudy.
	2	0	58	61	29,77		SSW	2	Cloudy.
4	7	0	48	60	29,61	0,190	SW	2	Fine.
	2	0	55	61	29,81		WSW	2	Fair.
5	7	0	38	58	30,13		W	1	Fair.
	2	0	47	61	30,18		W	1	Fine.
6	7	0	37	57	30,29			0	Foggy.
	2	0	48	59	30,26		SSE	1	Fine.
7	7	0	43	57	30,05		SE	2	Fair.
	2	0	49	59	29,94		SSE	2	Cloudy.
8	7	0	49	58	29,94		W	1	Cloudy.
	2	0	50	59	29,97		NW	1	Cloudy.
9	7	0	41	58	29,91		E	1	Foggy.
	2	0	53	59	29,77		SSE	1	Fair.
10	7	0	44	57	29,73		E	1	Fair.
	2	0	53	60	29,84		E	1	Fair.
11	7	0	43	59	30,05		WSW	1	Fair.
	2	0	53	59	30,12		WSW	1	Fair.
12	7	0	47	59	30,20		WSW	1	Fair.
	2	0	55	60	30,18		WSW	1	Cloudy.
13	7	0	51	59	29,85		SW	2	Cloudy.
	2	0	54	60	29,74		SSW	1	Cloudy.
14	7	0	36	58	30,08	0,200	W	1	Fine.
	2	0	42	59	30,10		WNW	1	Fine.
15	7	0	35	56	30,01	0,120	WNW	1	Cloudy.
	2	0	40	58	30,11		NW	1	Fine.
16	7	0	32	54	30,43		NNW	1	Fine.
	2	0	38	57	30,42		N	1	Fine.



METEOROLOGICAL JOURNAL

for November 1788.

1788	Time.		Therm.	Therm	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	inc es.	Inch.	Points	Str.	
			Deg.	Deg.					
Nov. 17	7	0	42	54	30,17		WNW	I	Fair.
	2	0	48	57	30,17		WNW	I	Fine.
18	7	0	38	54	30,17		WNW	I	Fine.
	2	0	45	57	30,22		WNW	I	Fine.
19	7	0	42	55	30,21				Foggy.
	2	0	48	57	30,13		WNW	I	Cloudy.
20	7	0	42	54	30,20				Foggy.
	2	0	45	57	30,20				Foggy.
21	7	0	45	54	30,26				Foggy.
	2	0	49	57	30,25		W	I	Cloudy.
22	7	C	47	56	30,25		SW	I	Cloudy.
	2	0	49	58	30,25		SSW	I	Cloudy.
23	7	0	39	55	30,20		ESE	I	Fine.
	2	0	47	58	30,15		ESE	I	Fine.
24	7	0	36	55	30,11		E	I	Fine.
	2	0	43	58	30,15		E	I	Fine.
25	7	0	38	54	30,22		E	I	Cloudy.
	2	0	38	58	30,28		ENE	I	Fine.
26	7	0	29	52	30,34		ENE	I	Fine.
	2	0	35	54	30,29		ENE	I	Cloudy.
27	7	0	28	50	30,11		ENE	I	Cloudy.
	2	0	27	52	30,05		ENE	I	Cloudy.
28	7	0	27	49	29,96				Cloudy.
	2	0	31	50	29,96				Cloudy.
29	7	0	30	47	30,01		ESE	I	Cloudy.
	2	0	34	49	30,04		ESE	I	Cloudy.
30	7	0	35	48	30,18		N by W	I	Cloudy.
	2	0	35	49	30,20		NE	I	Cloudy.

METEOROLOGICAL JOURNAL

for December 1788.

1788	Time.		Therm. without	Therm. within.	Barom.	Rain.	Winds.		Weather.
	H.	M.	Deg.	Deg.	Inches.	Inch.	Points.	Str.	
Dec. 1	8	0	34	48	30,22		NE	1	Cloudy.
	2	0	36	49	30,20		ENE	1	Fine.
2	8	0	31	47	29,93		ENE	1	Fine.
	2	0	35	49	29,85		ENE	1	Fine.
3	8	0	24	46	29,68		W	1	Cloudy.
	2	0	31	48	29,61		W	1	Fair.
4	8	0	24	44	29,59		N	1	Fine.
	2	0	32	49	29,58		N	1	Fine.
5	8	0	34	45	29,68		NE	1	Cloudy.
	2	0	35	49	29,75		ENE	1	Cloudy.
6	8	0	33	45	29,84		NE	2	Cloudy.
	2	0	34	48	29,83		NE	2	Cloudy.
7	8	0	34	45	29,89		ENE	2	Cloudy.
	2	0	35	46	29,93		ENE	2	Cloudy.
8	8	0	33	44	30,00		ENE	1	Cloudy.
	2	0	38	48	30,00		E	1	Fine.
9	8	0	33	45	29,98		N	1	Fair.
	2	0	39	49	29,95		N	1	Fine.
10	8	0	34	47	29,86		N	1	Cloudy.
	2	0	40	51	29,88		ENE	1	Fine.
11	8	0	32	47	30,04				Foggy.
	2	0	35	50	30,07		WNW	1	Cloudy.
12	8	0	29	46	30,10		WNW	1	Cloudy.
	2	0	32	49	30,07		WNW	1	Fair.
13	8	0	27	45	29,96				Foggy.
	2	0	29	47	29,85		NE	1	Cloudy.
14	8	0	26,5	44	29,55		NE	1	Fair.
	2	0	30	46	29,50		NE	1	Cloudy.
15	8	0	22	42	29,61		NE	2	Cloudy.
	2	0	24	45	29,63		NE	2	Cloudy.
16	8	0	22	40	29,65		ENE	2	Fair.
	2	0	27	44	29,68		ENE	2	Fine.

METEOROLOGICAL JOURNAL

for December 1788.

1788.	Time.		Therm.	Therm.	Barom.	Rain.	Winds.		Weather.
	H.	M.	without	within.	Inches.	(inch.	Points.	Str.	
Dec. 17	8	0	26	41	30,00		ENE	2	Cloudy.
	2	0	28	44	30,08		ENE	1	Snow.
18	8	0	18	40	30,14		WNW	1	Cloudy.
	2	0	28	44	30,00		WNW	1	Cloudy.
19	8	0	26	41	29,90		NNW	1	Fine.
	2	0	32	44	29,94		NNW	1	Cloudy.
20	8	0	30	42	29,98		WNW	1	Cloudy.
	2	0	33	45	29,95		WNW	1	Fine.
21	8	0	36	42	29,66		NW	2	Cloudy.
	2	0	36	44	29,68		NW	2	Cloudy.
22	8	0	29,5	42	29,97		WNW	2	Fine.
	2	0	35	46	30,03		WNW	2	Fine.
23	8	0	23	42	30,22		WNW	1	Foggy.
	2	0	25	45	30,30		WNW	1	Foggy.
24	8	0	39	43	29,96		W	2	Cloudy.
	2	0	43	47	29,96		W	1	Cloudy.
25	8	0	41	45	29,94		W	1	Fair.
	2	0	46	48	29,80		W	2	Fine.
26	8	0	31	45	29,67		NNE	2	Cloudy.
	2	0	34	48	29,90		NNE	2	Fine.
27	8	0	30	45	29,97		NNW	2	Cloudy.
	2	0	32	47	29,90		NNW	2	Cloudy.
28	8	0	23	43	30,15		NNE	1	Fine.
	2	0	26	45	30,22		NNE	1	Fine.
29	8	0	19	41	30,30		NE	1	Fine.
	2	0	26	45	30,26		NE	1	Fine.
30	8	0	18	40	30,33		E	1	Fine.
	2	0	21	44	30,31		SE	1	Fine.
31	8	0	26	39	30,03		SSW	2	Cloudy.
	2	0	30	42	29,80		SSW	2	Snow.

1788	Thermometer without.			Thermometer within.			Barometer.			Rain.
	Greatest height.	Least height.	Mean height.	Greatest height.	Least height.	Mean height.	Greatest height.	Least height.	Mean height.	
	Deg.	Deg.	Deg.	Deg.	Deg.	Deg.	Inches.	Inches.	Inches.	Inches.
January	48	26	39,7	56	49	52,7	30,70	28,89	29,97	0,439
February	50	29	41,3	56	50	52,7	30,21	28,65	29,68	1,461
March	59	28	40,8	59	46	50,9	30,08	29,32	29,68	0,336
April	68	40	52,6	63	55	51,8	30,48	29,50	30,07	0,607
May	80	49	60,0	73	59	62,8	30,34	29,58	30,04	0,497
June	80	52	62,3	70	61	64,1	30,22	29,49	29,94	3,275
July	77	55	63,7	69	64	65,9	30,22	29,73	29,99	1,620
August	77	53	63,4	71	64	66,0	30,45	29,22	29,95	2,699
September	74	45	58,6	67	59	63,7	30,25	29,37	29,86	3,345
October	67	33	51,4	62	54	59,4	30,55	29,64	30,32	0,103
November	58	27	42,9	61	47	56,4	30,50	29,61	30,11	0,510
December	46	18	30,9	51	39	45,2	30,33	29,50	29,92	0,000
Whole year			50,6			57,6			29,96	14,892

PHILOSOPHICAL  
TRANSACTIONS,  
OF THE  
ROYAL SOCIETY  
OF  
LONDON.

VOL. LXXIX. For the Year 1789.

PART II.



*Bosch's sc.*

LONDON,  
SOLD BY LOCKYER DAVIS, AND PETER ELMSLY,  
PRINTERS TO THE ROYAL SOCIETY.

MDCCLXXXIX.



---

---

# C O N T E N T S

O F


## V O L. LXXIX. P A R T II.

- XI. *EXPERIMENTS on the Phlogistication of Spirit of Nitre.* By the Rev. Joseph Priestley, LL.D. F. R. S. Page 139
- XII. *Observations on a Comet. In a Letter from William Herschel, LL.D. F. R. S. to Sir Joseph Banks, Bart. P. R. S.* p. 151
- XIII. *Indications of Spring, observed by Robert Marsham, Esquire, F. R. S. of Stratton in Norfolk. Latitude 52° 45'.* p. 154
- XIV. *An Account of a Monster of the human Species, in two Letters; one from Baron Reichel to Sir Joseph Banks, Bart. and the other from Mr. James Anderson to Baron Reichel. Communicated by Sir Joseph Banks, Bart. P. R. S.* p. 157
- XV. *A supplementary Letter on the Identity of the Species of the Dog, Wolf, and Jackal; from John Hunter, Esq. F. R. S. addressed to Sir Joseph Banks, Bart. P. R. S.* p. 160
- XVI. *Abstract of a Register of the Barometer, Thermometer, and Rain, at Lyndon in Rutland; by Thomas Barker, Esq. Also of the Rain in Hampshire and Surrey. Communicated by Thomas White, Esq. F. R. S.* p. 162
- XVII. *On the Method of correspondent Values, &c. By Edward Waring, M. D. F. R. S. and Lucasian Professor of Mathematics at Cambridge.* p. 166

XVIII.

- XVIII. *On the Resolution of attractive Powers.* By Edward Waring, M. D. F. R. S. and Lucasian Professor of Mathematics at Cambridge. p. 185
- XIX. *Experiments on the Congelation of Quicksilver in England.* By Mr. Richard Walker; in a Letter to Henry Cavendish, Esq. F. R. S. p. 199
- XX. *Catalogue of a second Thousand of new Nebulæ and Clusters of Stars; with a few introductory Remarks on the Construction of the Heavens.* By William Herschel, L L. D. F. R. S. p. 212
- XXI. *An Attempt to explain a Difficulty in the Theory of Vision, depending on the different Refrangibility of Light.* By the Rev. Nevil Maskelyne, D. D. F. R. S. and Astronomer Royal. p. 256
- XXII. *Experiments and Observations on Electricity.* By Mr. William Nicholson; communicated by Sir Joseph Banks, Bart. P. R. S. p. 265
- XXIII. *Experiments on the Transmission of the Vapour of Acids through an hot earthen Tube, and further Observations relating to Phlogiston.* By the Rev. Joseph Priestley, LL.D. F. R. S. p. 289
- XXIV. *On the Production of nitrous Acid and nitrous Air.* By the Rev. Isaac Milner, B. D. F. R. S. and President of Queen's College, Cambridge. p. 300





P H I L O S O P H I C A L

T R A N S A C T I O N S.

---

XI. *Experiments on the Phlogistication of Spirit of Nitre.*  
By the Rev. Joseph Priestley, LL.D. F. R. S.

Read March 26, 1789.

**A**S the colouring of spirit of nitre has some connection with the doctrine of phlogiston, to which I propose to give my best attention, I have lately resumed my experiments on that subject, and beg leave to lay the result of them before the Society.

In my former experiments, vol. IV. p. 2. I found that the colourless acid became smoking, or orange-coloured, and emitted orange-coloured vapours, on being exposed to heat in long glass tubes, hermetically sealed; and I then concluded, that

this effect was produced by the action of *heat*, evolving, as it were, the phlogiston previously contained in the acid. Afterwards, having found that it was not *heat*, but *light* only, that was capable of giving colour to spirit of nitre, contained in phials with ground stoppers, in the course of several days; and that in this case the effect was produced by the action of *light* upon the *vapour*, which gradually imparted its colour to the liquor on which it was incumbent (see Vol. V. p. 342.), I was led to suspect, that as the glass tubes, in which I had formerly exposed this acid to the action of heat, were only held near to a fire, in the day-light, or candle-light, it might have been this *light*, which, in these circumstances, had, at least in part, contributed to produce the effect.

In order to ascertain whether the light had had any influence in this case, I now put the colourless spirit of nitre into long glass tubes, like those which I had used before, and also sealed them hermetically, as I had done the others; but, instead of exposing them to heat in the open air, from which light could not be excluded, I now shut them up in gun barrels, closed with metal screws, so that it was impossible for any particle of light to have access to them; and I then placed one end of the barrels so near to a fire as was sufficient to make the liquor contained in the tube to boil, which I could easily distinguish by the sound which it yielded. The consequence was, that in a short time the acid became as highly coloured as ever it had been when exposed to heat without the gun barrel. It was evident, therefore, that it had been mere *heat*, and not *light*, which had been the means of giving this colour to the acid, and which has been usually termed *phlogisticating* it.

When

When I made the former experiments, I had no suspicion that the *air* contained in the tube had any concern in the result of them; and, in those which I made in the phials in a moderate heat, I found that the acid received its colour when the best *vacuum* that I could make with an air pump was over it.

My friend Mr. KIRWAN, however, having always suspected, that the *air* was a principal agent in the business, I at this time gave particular attention to this circumstance; supposing that, if any part of the common air had been imbibed, it must have been the *phlogificated*, and that it was the phlogiston from this kind of air which had phlogificated the acid. The real result, however, was not so much in favour of this supposition as I had expected; for the principal effect of the process was the emission of dephlogificated air, so that the acid seems to become what we call phlogificated, by parting with this ingredient in its composition.

I put a small quantity of the colourless acid into a long glass tube, which besides the acid would have contained 1.23 ounce measures of common air, but that the vapour of the acid excluded about one-twentieth of the quantity. Having sealed the tube hermetically, I shut it up in a gun barrel, in the manner mentioned above, and exposed it to a boiling heat for several hours, and then opening it under water there came out of it 2.03 ounce measures of air, very turbid and white; and when it was examined, it appeared to be of the standard of 1.02, with two equal measures of nitrous air; when with one measure of the same nitrous air the standard of the common air was 1.07. The quantity of phlogificated air absorbed in this experiment I ascertained by the following computation.

As one measure of common air, and an equal quantity of nitrous air were reduced to 1.07 m. it is evident, that 0.93 m.

had disappeared; but as this was effected by the nitrous air uniting with all the dephlogisticated air contained in the common mass, and as they unite in the proportion of one measure of dephlogisticated air to two measures of nitrous air, one-third of the 0.93 m. *viz.* 0.31 m. will be the quantity of dephlogisticated air that was contained in the one measure of common air on which the experiment was made, the remainder, *viz.* 0.69, having been phlogisticated air. The common air contained in the tube would have been 1.23 oz. m.; but deducting from it one-twentieth in the whole, it will only be 1.17 oz. m. I then say, if one measure of this air contains 0.69 m. of phlogisticated air, 1.17 oz. m. will contain 0.8073 oz. m. of phlogisticated air. This, therefore, was the quantity of phlogisticated air which had been exposed to the action of the acid of nitre in the tube.

In order to find how much of the same kind of air was contained in the tube *after* the process, I examined the result above mentioned in the following manner. Since two measures of nitrous air, and one of this residuum, were reduced to 1.02 m. it is evident, that 1.98 m. had disappeared, and consequently one-third of this quantity, *viz.* 0.66 m. had been dephlogisticated air, and that the remainder of the measure, *viz.* 0.34, had been the proportion of phlogisticated air in one measure of this residuum. If then one measure of this residuum contains 0.34 m. of phlogisticated air, 2.03 oz. m. will contain 0.6902 oz. m. which is less than 0.8073 oz. m. the quantity contained in it before the process; so that a part of the phlogisticated air had been either absorbed or decomposed, its phlogiston having been imbibed by the acid at the same time that it had emitted the dephlogisticated air.

In another process, of the same kind, the glass tube contained 0.92 oz. m. of common air, and the air that came out of it after the process was one ounce measure, of the standard of 1.6 with two measures of nitrous air, and computing as I did before, the phlogisticated air in the tube before the process was 0.6072 oz. m., and after the process 0.54 oz. m.

In these computations it is supposed, that the air emitted by the acid was perfectly pure, so that all the phlogisticated air that is found after the process is supposed to have been contained in the common air confined in the tube before it was commenced. But I found, that the air emitted by the acid is by no means perfectly pure, so that much of the impurity must be ascribed to this circumstance.

In order to exclude all air from the contact of the acid, I made a quantity of it to boil in the tube, and when the vapour had expelled all the air, I sealed it hermetically, in the manner in which water hammers are made; and then exposing it to heat, found that it acquired as high a colour as when air had been confined along with it; so that it is evident, that *air* is not necessary to this effect. When the tube was opened under water, a quantity of dephlogisticated air rushed out, exceedingly white as before; but when I examined it, I found it to be of the standard of only 0.66. When this impurity is considered, it will appear, that when much air is yielded in this process, some phlogisticated air may have been imbibed, though, computing in the manner above mentioned, the phlogisticated air after the process should be in greater quantity than was contained in the tube before it, as was the case in the following experiment.

In a glass tube which, besides the acid, contained 1.13 oz. m. of common air, I exposed colourless spirit of nitre to heat

till it became of a deep orange colour; and when it was opened under water, there came out of it 2.83 oz. m. of air exceedingly turbid, of the standard of 0.66, with two equal quantities of nitrous air, when that of the common air, with one equal quantity of nitrous air, was 1.07. Computing in the manner above mentioned, there was in the tube before the process 0.7477 oz. m. of phlogificated air, and after the process 0.8792 oz. m. But the dephlogificated air, amounting to 1.7 oz. m. being of the standard of 0.66; will be found to contain 0.374 oz. m. of phlogificated air, which being deducted from 0.8792, there will remain only 0.5052 oz. m. which is considerably less than 0.7477 oz. m.

That the nitrous acid can become coloured, without imbibing any thing from phlogificated air, is evident not only from its becoming so when heated *in vacuo*, as described above, but also, when it was in contact with any other kind of air, as free from phlogificated air as I could make it. But from the manner in which these experiments were necessarily made, it was impossible intirely to exclude phlogificated air, either as part of the atmospheric air, or as contained in the impurities of the air that I made use of; for I first filled the tube with spirit of nitre, then plunging the orifice of it in a vessel of the same, I introduced a quantity of the air which I wished to expose to it. After this, putting my finger upon the orifice, I turned it upside down, and applying to it the closed end of a glass tube, of about the same diameter, I sealed it hermetically with a blow-pipe as expeditiously as I could. This is a necessary imperfection in the experiment; but I know not how to remedy it, if any of the acid is to be left in the tube. However, the phlogificated air introduced in this manner from the atmosphere must have borne a very small proportion to the air in the

the tube; and some objection will always remain to the experiment from the impurity of the air made use of.

Having repeatedly observed, that the acid became coloured in consequence of being exposed to heat in contact with any kind of air whatever, I exposed at the same time, and in the same circumstances, three equal quantities of the same colourless spirit of nitre, in three nearly equal tubes, one containing dephlogisticated, another phlogisticated, and a third inflammable air; that, if there should be any difference in the colouring of the acid in these cases, it might be the more easily perceived. But though I gave all the attention that I could, I did not perceive that there was any difference, except what arose from some of the tubes being placed a little nearer the fire than the rest; and, by changing their places, the colour was at length the very same in them all.

As in these three cases I examined the air before and after the process, in the manner above mentioned, I shall just recite the particulars.

Of the dephlogisticated air the tube contained before the process 1.46 oz. m. of the standard of 0.67, and after the process it contained 1.76 oz. m. of the standard of 0.77; a difference owing in part to the mixture of common air, which could not be excluded in the sealing of the tube, and in part to the air emitted from the acid not being pure.

Of the phlogisticated air, the tube contained 1.3 oz. m. and after the process 1.95 oz. m. of the standard of 1.38.

Of the inflammable air, the tube contained before the process 1.52 oz. m. and after the process 1.9 oz. m. of the standard of 1.8. They were all measured by a mixture of two equal quantities of nitrous air.

If

If these results be examined as that of the first experiment, with common air, it will be found that, in all these processes, there was less phlogificated air, or inflammable air, after the process than before; and this result being thus uniform, I cannot help concluding, that this kind of air is in part decomposed, and purified by this means; so that by this emission of dephlogificated air which the heat expels from the acid, something, and probably phlogiston, is at the same time imbibed from it; which proves that phlogificated air is no simple substance, but a compound, and that phlogiston is one constituent part of it; for this acid acquires the same colour, and all the same properties, by adding to it any thing that is supposed to contain phlogiston.

As the spirit of nitre can be rendered smoking, or phlogificated, by the mere expulsion of dephlogificated air, it is evident, that it contains two principles in close affinity with each other, and that nothing is necessary to render either of them conspicuous besides the absence of the other.

It is also natural to suppose, that, for the same reason that the *dephlogificating* principle (as it may be called) is expelled, the *phlogificating* principle should enter; so that the purification of the air in contact with the acid may be a necessary consequence of the expulsion of the pure air contained in it, the whole tending, as it were, to an equilibrium in this respect. It is therefore by no means difficult to conceive, that phlogiston should be extracted from the contiguous air at the same time that the dephlogificated air *not pure* (that is, containing a mixture of phlogificated air) is driven out of it; for the acid always containing phlogiston, whatever air is contained in it, and expelled from it, may necessarily contain phlogiston or phlogificated air; but the purer air may be emitted, and the  
less



less pure air be imbibed, till the whole come to be of the same quality. It may, however, perhaps follow from the emission of impure dephlogificated air, and the imbibing of phlogificated air at the same time, that the former does not consist of dephlogificated and phlogificated air loosely mixed, but of some intimate union of dephlogificated air with phlogiston, though they may be separated by a mixture of nitrous air, and other processes, in the very same manner as dephlogificated air may be separated from a loose mixture of phlogificated air.

It is evident from these experiments, that a red heat is not necessary to the conversion of nitrous acid into pure air, though this process, as appeared by my former experiments, produces this effect most quickly and effectually.

I cannot help considering the experiments above recited to be favourable to the doctrine of the phlogiston, and unfavourable to that of the decomposition of water, though not decisively so; for since the red vapour of spirit of nitre unquestionably contains the same principle that has been termed phlogiston, or the principal element in the constitution of inflammable air, and according to the antiphlogistians this is one constituent part of water, they must suppose, that the water in this acid is decomposed by a much more moderate heat than in most other cases. In general, I believe, they have thought a red heat to be necessary for this purpose. It is evident, that the conversion of water into steam by boiling, or by any heat that can be given to it under the strongest pressure, has no tendency whatever to decompose it. But if the mere boiling of water in nitrous acid could produce this effect, I do not see why the same should not be the case when water alone is boiled.

I think it will also be more difficult to explain the purification of the incumbent atmospherical air on the antiphlogistic

than on the phlogistic hypothesis, whatever be the constitution of phlogisticated air.

As, in the experiments above mentioned, *heat* without *light* gives colour to the nitrous acid, and the reflection or refraction of light is always attended with heat, it may perhaps be *heat* universally that is the means of imparting this colour, though the mode of its operation be at present unknown. And in these experiments, as well as the former, it is the *vapour* that first receives the colour, and imparts it to the liquid when it is sufficiently cold to receive it.

The rushing out of a quantity of turbid white air from a transparent tube, quite cold, is a striking phænomenon in these experiments. It may be worth while to examine of what it is that this remarkable cloudiness of the air consists. There is the same appearance, as I have more than once observed, in the rapid production of any kind of air, which is perfectly transparent as it passes along the glass tube through which it is transmitted, till it comes into contact with the water in which it is received.

P. S. Not to multiply my communications on the subject of *phlogiston* unnecessarily, I would beg leave to observe, at the close of this article (in reply to what has been objected to my former experiments, as being liable to exception from the phlogisticated air which could not be excluded from the dephlogisticated air when it was decomposed by means of inflammable air) that I have found the process I made use of to have no tendency whatever to decompose phlogisticated air. Indeed, nothing that we have hitherto known concerning this kind of air could make it probable, that mere *heat*, in contact with dephlogisticated or inflammable air, *could* have this effect. And

it is of no consequence whatever to say, that any particular substance, imagined to be decomposed, is *present* in a process, unless it can be shewn that, in that process, there are agents capable of decomposing it. If mere *heat* (which is all that my process requires) would decompose phlogisticated air, and reduce it to nitrous acid, the transmission of common air (which consists of dephlogisticated and phlogisticated air) through a red hot tube would have this effect, which it is well known not to have.

But what I have asserted above is a conclusion which I have drawn from comparing the decomposition of dephlogisticated air by the two processes with nitrous and inflammable air. That nitrous air, when mixed with dephlogisticated air, has no tendency to produce phlogisticated air, is evident from the almost total evanescence of both of them, when they are very pure, and mixed in due proportions; and that nitrous air has no effect on phlogisticated air is well known. If then the firing of dephlogisticated and inflammable air had a tendency to decompose any portion of phlogisticated air, which should happen to be mixed with them, less would remain after the firing of inflammable and impure dephlogisticated air than after mixing it with nitrous air; for as the impurities of dephlogisticated air consist of phlogisticated air, those would disappear in a greater proportion in the former process than in the latter. But by many careful trials I find, that I can reduce any kind of dephlogisticated air no farther by a mixture of inflammable air than I can by nitrous air. When the proportions are well managed, the diminution is as nearly as possible the same in both the cases.

I must observe, however, that it requires more nitrous air than inflammable air (from iron by steam) to produce this effect

in the proportion of about 10 to 9 ; so that nitrous air does not contain quite so much phlogiston as an equal bulk of inflammable air, as I had before thought to be the case.

In this Paper it will be observed, that I make the diminution of common air by nitrous air to be considerably less than I have usually done before. This has been the consequence of giving the two kinds of air a little agitation at the instant of mixing, which will generally make the diminution less by two tenths of a measure. But I have found, that when these mixtures of air, with and without agitation, have been kept some time, they approach to an equality of bulk.

At the same time I have observed, what I think not a little extraordinary, that agitation prevents the greatest diminution of dephlogisticated and nitrous air. I have found it to be 2.5 without agitation, and 6. with it.

The less diminution of the mixture of nitrous and common air is probably owing to the presence of so much phlogisticated air, which impedes the meeting of the nitrous air with the dephlogisticated air in the mixture ; because I find the same to be the case when I mix the same proportion of inflammable air with dephlogisticated air ; and when dephlogisticated air is agitated with nitrous air, the *water* may impede their union, as the phlogisticated air did before.

There is, therefore, no source of the *nitrous acid* which I find on the decomposition of dephlogisticated and inflammable air, besides the union of those two kinds of air, which therefore do not make *mere water*, as the antiphlogistians suppose.



XII. *Observations on a Comet. In a Letter from William Herschel, LL.D. F. R. S. to Sir Joseph Banks, Bart. P. R. S.*

Read April 2, 1789.

S I R,

Slough, March 3, 1789.

**T**HE last time I was in town, you expressed a wish to see my observations on the comet which my sister, CAROLINE HERSCHEL, discovered in the evening of the 21st of last December, not far from  $\beta$  Lyræ.

As she immediately acquainted the Rev. Dr. MASKELYNE, and several other gentlemen, with her discovery, the comet was observed by many of them. The Astronomer Royal, in particular, having, I find, obtained a very good set of valuable observations on its path, it will be sufficient if I communicate only those particulars which relate to its first appearance, and a few other circumstances that may perhaps deserve to be noticed.

December 21, 1788, about 8 o'clock, I viewed the comet which my sister had a little while before pointed out to me with her small Newtonian *sweeper*. In my instrument, which was a ten-foot reflector, it had the appearance of a considerably bright nebula; of an irregular, round form; very gradually brighter in the middle; and about five or six minutes in diameter. The situation was low, and not very proper for instruments with high powers.

December 22, about half after five o'clock in the morning, I viewed it again, and perceived that it had moved apparently in a direction towards  $\delta$  Lyræ, or thereabout. I had been engaged all night with the twenty-foot instrument, so that there had been no leisure to prepare my apparatus for taking the place of the comet; but in the evening of the same day, I took its situation three times, as follows:

Dec. 22.	at	<sup>h.</sup> 23	' 42	" 19	fidereal time, the comet passed the wire,
	at	23	49	24	$\beta$ Lyræ passed the same,
Difference		<hr/>			very accurate.
		7	5		
		<hr/>			

	at	23	52	52	the comet passed,
	at	23	59	58	$\beta$ Lyræ passed,
Difference		<hr/>			accurate.
		7	6		
		<hr/>			

	at	0	6	35	the comet passed,
	at	0	13	40	$\beta$ Lyræ passed,
Difference		<hr/>			very accurate.
		7	5		
		<hr/>			

I found in every observation the small star which accompanies  $\beta$  Lyræ\*, exactly in the parallel of the comet.

These transits were taken with a ten-foot reflector; and the difference in right ascension, I should suppose, may be depended upon to within a second of time. The determination


\* For this small star see my Catalogue of Double Stars, in the Philosophical Transactions for the year 1782, Part I. Class V. Star 3. where its distance and position are given, and consequently its parallel may be found.


also of the parallel can hardly err so much as fifteen seconds of a degree.

This, and several evenings afterwards, I viewed the comet again with such powers as its diluted light would permit, but could not perceive any sort of nucleus, which, had it been a single second in diameter, I think, could not well have escaped me. This circumstance seems to be of some consequence to those who turn their thoughts on the investigation of the nature of comets; especially as I have also formerly made the same remark on one of the comets discovered by M. MECHAIN in 1787, a former one of my sister's in 1786, and one of Mr. PIGOTT's in 1783; in neither of which any defined, solid nucleus could be perceived.

I have the honour to remain, &c.

WILLIAM HERSCHEL.








XIII. Indications of Spring, observed by Robert Marham, Esquire, F. R. S. of Stratton in Norfolk. Latitude 52° 45'.

Read April 2, 1789.

Dates.	Snow-drop flower.	Thrush fings.	Hawthorn leaf.	Hawthorn flower.	Frogs and Toads croak.	Sycamore leaf.	Birch leaf.	Elm leaf.	Mountain-ash leaf.	Oak leaf.	Beech leaf.	Horse-chestnut leaf.	Chestnut leaf.	Hornbeam leaf.	Ash leaf.	Ringdoves coo.	Rooks build.	Young Rooks.	Swallows appear.	Cuckoo fings.	Nightingale fings.	Churn Owl fings.	Yellow Butterfly appears.	Turnip in flower.	Lime leaf.	Maple leaf.	Wood Anemone flower.
1736		1735 Dec. 4.																	March 30.								
1738				Nifmes, France, Apr. 14, N.S.															Piacenza, Italy, Mar. 20, N. S.								
1739			Feb. 23.			Feb. 23.													April 13.	April 19.	April 25.			Feb. 21.			
1740		Feb. 27.	April 4.	May 28.		April 14.											Feb. 27.		March 31.	April 14.	May 2.	May 21.		May 2.			
1741				May 9.	March 12.											Feb. 2.	Feb. 13.	April 2.	April 9.	April 19.	April 14.						
1742		Jan. 31.		May 15.	April 11.													April 1.	April 17.	April 17.	April 21.			April 15.			
1743		Jan. 31.		May 8.	Feb. 28.														April 12.	April 17.	April 20.			March 23.			
1744			March 28.	Essex, May 12.	March 28.	March 30.											Feb. 21.	April 1.	March 31.	April 20.	April 9.	May 23.		April 8.			
1745	Jan. 6.	Jan. 26.	March 26.	May 13.	March 14.	March 29.	March 29.					March 29.				Jan. 3.	March 8.	April 8.	April 3.	April 22.	April 12.	May 10.		April 8.			
1746	Jan. 20.		March 26.	May 13.	March 15.	April 10.	April 10.			May 1.	May 1.						Feb. 25.	March 31.	April 8.	April 16.	April 19.	May 5.		April 16.	May 1.		
1747	Jan. 8.	Jan. 14.	Feb. 15.	April 26.	Feb. 24.	March 25.	March 29.			April 23.	April 26.					Jan. 13.	Feb. 13.	March 26.	April 2.	April 22.	April 15.	May 9.		March 18.			
1748	Jan. 5.	Jan. 29.	April 3.	Middlesex, May 22.	March 28.	April 14.	April 10.									Feb. 8.	Feb. 29.	April 3.	April 2.	April 16.	April 23.	May 28.		April 22.			
1749	Jan. 4.	Jan. 17.	Feb. 19.	Middlesex, May 1.	March 5.	March 18.	March 24.			April 22.	April 22.					Feb. 13.	Feb. 18.	March 29.	April 5.	April 13.	April 16.	May 20.		March 9.			
1750	Jan. 15.	Jan. 17.	Feb. 13.	April 13.	Feb. 20.	Feb. 22.	Feb. 21.			March 31.	April 15.					Jan. 22.	Feb. 13.	March 30.	April 8.	April 11.	April 9.	May 17.	Feb. 9.	Feb. 22.			
1751	Jan. 9.	Jan. 2.	March 10.	May 9.	March 27.		March 22.	March 6.		April 25.	April 24.	March 21.			April 16.	1750 Dec. 29.	March 3.	April 7.		April 19.	April 16.	May 3.		March 29.			
1752		Jan. 30.	Middlesex, Feb. 18.	May 14.	March 9.	April 6.	April 2.			April 20.	April 20.	March 19.				1751 Dec. 27.		March 29.	April 2.	April 9.	April 7.	May 12.		April 6.			
N. Style 1753	Feb. 1.	Feb. 1.	March 21.	May 11.	April 1.	April 3.	March 27.			April 24.	April 29.				May 11.	Feb. 22.	March 3.	April 15.	April 17.	April 24.	April 19.			March 30.			
1754	Jan 18.	Feb. 16.	April 7.	May 22.	April 6.	April 14.	April 14.			May 11.	May 7.						March 1.	April 11.	April 13.	April 24.	April 11.	May 16.		May 9.			
1755	Jan. 26.	Feb. 16.	March 31.	May 10.	April 1.	April 9.	April 1.	April 10.	April 9.	April 18.	April 21.	March 31.	April 16.	April 13.	April 22.	March 4.	March 14.	April 18.	April 6.	April 23.	April 14.	June 4.		April 15.	April 12.	April 18.	



Indications of Spring continued.

Dates.	Snow-drop flower.	Thrush fings.	Hawthorn leaf.	Hawthorn flower.	Frogs and Toads croak.	Sycamore leaf.	Birch leaf.	Elm leaf.	Mountain-ash leaf.	Oak leaf.	Beech leaf.	Horse-chestnut leaf.	Chestnut leaf.	Hornbeam leaf.	Ash leaf.	Ringdoves coo.	Rooks build.	Young Rooks.	Swallows appear.	Cuckoo fings.	Nightingale fings.	Churn Owl fings.	Yellow Butterfly appears.	Turnip in flower.	Lime leaf.	Maple leaf.	Wood Anemone flower.		
1756	Jan. 8.	Jan. 30.	Feb. 26.	May 25.	March 10.	April 1.	March 11	March 30.	April 5.	May 7.	May 7.					Feb. 22.	March 1.	April 8,	April 7.	April 18.	April 14.	May 16.		1755, Dec. 28. and March 4.	May 7.				
1757	Feb. 6.	Feb. 16.	March 20.	May 19.	March 29.	April 1.	March 29.	April 1.	March 31.	April 26.	April 29.				May 1.	Mar. 10.	March 13.	April 18.	April 16.	April 26.	April 16.	May 17		April 11.					
1758	Jan. 9.	Feb. 2.	March 19.	May 13.	March 16.	April 1.	April 1.	April 1.	April 13.	April 29.	April 27.	April 13.		March 15.	May 8.	March 3.	March 5.	April 13.	April 10.	April 30.	Middlefex, April 23.	May 18.	March 15.	April 14.					
1759	Jan. 9.	Jan. 17.	Feb. 11.	May 6.	March 1.	Middlefex, March 22.	Middlefex, March 31.	Middlefex, March 19.	April 1.	April 15.	April 25.	Middlefex, March 25.	April 21.	Middlefex, March 30.	April 23.	Feb. 11.	Feb. 28.	April 10.	April 10.	April 26.	April 21.	June 6.		Feb. 27	April 18.	April 18.			
1760	Jan. 24.	Feb. 10.	Kent, March 7.	April 27.	March 15.	April 3.	April 2.	April 9.	April 2.	April 19.	April 19.	April 12.	April 19.	April 13.	April 19.		Middlefex, March 12.	April 11.	April 11.	April 21.	April 21.	May 29.		April 13.	April 19.	April 13.			
1761		Jan. 4.	Surrey, Feb. 27.	May 4.	March 22.	March 25	March 24.	Surrey, Feb. 27,	March 25.	April 14.	April 20.	March 29.	April 10.	April 10.	May 3.	March 20.	Effex, March 11.	April 7.	April 5.	May 2.	April 17.	May 30.	March 12	Jan. 15.	April 4.	April 17.	March 29.		
1762	Jan. 12.	Jan. 22.	Surrey, March 7.	May 4.	April 19.	Middlefex, April 16.	April 6.	Middlefex, April 12.	April 19.	April 22.	April 21.	Middlefex, April 16.	April 20.	Herts. April 18	April 25.			April 20.	April 8.	April 20.	April 19.	May 31.	April 5.	April 19.	April 16.	April 21.			
1763		London, Feb. 11.	Surrey, Feb. 26.	May 7.	March 30.	March 28.	March 25.	April 1.	March 28.	April 24.	April 20.	Surrey, March 10.	April 18.	April 3.	May 7.			April 16.	April 5.	April 27.	April 20.	May 31.		April 17.	April 15.				
1764	Jan. 6.	Jan. 9.	Feb. 17.	Middlefex, May 17.				London, March 21.	Surrey, April 8.	Herts. April 24.	Herts. April 24.	Surrey, April 8.			Kent, May 5.	Feb. 22.	March 6.		London, April 8.	Herts. April 24.	Kent, May 2.			Middlefex, March 18.					
1765	Jan. 5.	Feb. 11.	March 16.	Kent, May 11.	March 23.	Middlefex, April 7.		Middlefex, March 29.		Herts. May 1.	Herts. May 1.	Middlefex, March 31.			Herts. May 1.	Feb. 25.	March 7.		Middlefex, April 14.	Middlefex, April 12.	Herts. May 3.	June 4.		Middlefex, April 10.		March 24.			
1766		Feb. 23.	March 16.		March 9.	Surrey, April 13.	March 11.	Middlefex, April 12.	Surrey, May 2.	Surrey, May 2.		Surrey, April 13.		Surrey, May 2.	Surrey, May 2.	March 3.	March 7.	Middlefex, April 24.	Middlefex, April 17.	Surrey, May 2.	Surrey, May 2.	May 26.	March 6		Surrey, April 12.				
1767	Jan. 28	Feb. 1.	Surrey, March 12.	May 29.		Surrey, April 19.	Surrey, April 19.	Middlefex, March 14.		Effex, May 3.	May 5.	Middlefex, April 8.							Surrey, April 23.	May 7.	Effex, May 3.			London, March 22.					
1768	Jan. 28.		Surrey, Feb. 28.	May 11.	March 17.	March 23.	March 21.	April 4.	April 4.	April 24.	April 24.	April 4.		April 19.	April 27.		Middlefex, Feb. 29.	April 10.	April 17.	April 24.	April 17.			March 28.	April 20.				
1769	Jan. 13.		Effex, March 4.	Effex, May 4.		Middlefex, April 17.		Kent, March 25.	Middlefex, April 6.	Kent, April 25.		Middlefex, April 6.	Kent, April 25.		Effex, May 4.				Surrey, April 19.	Kent, April 25.	Suffex, May 23.	June 10.		London, April 16.					
1770	Jan. 21.	Jan. 28.	Surrey, March 11.	May 22.		Middlefex, April 23.	May 1.	Middlefex, March 11.	May 1.	May 11.	May 6.	Middlefex, April 8.	May 12.	May 1.	May 12.		Feb. 23.		Middlefex, April 23.	May 1.	May 6.		Feb. 14.	April 23.	May 2.	May 7.			



Indications of Spring continued.

Dates.	Snow-drop flower.	Thrush fings.	Hawthorn leaf.	Hawthorn flower.	Frogs and Toads croak.	Sycamore leaf.	Birch leaf.	Elm leaf.	Mountain-ash leaf.	Oak leaf.	Beech leaf.	Horse-chestnut leaf.	Chestnut leaf.	Hornbeam leaf.	Ash leaf.	Ringdoves coo.	Rooks build.	Young Rooks.	Swallows appear.	Cuckoo fings.	Nightingale fings.	Churn Owl fings.	Yellow Butterfly appears.	Turnip in flower.	Lime leaf.	Maple leaf.	Wood Anemone flower.
1771	Jan. 27.	Feb. 21.	April 11.	May 25.	April 6.	May 4.	May 4.	Middlefex, April 24.	May 2.	May 15.	May 10.	May 2.	May 10.	May 7.	May 16.	Feb. 26.	March 15.		Middlefex, April 18.	May 5.	May 3.	May 20.	Feb. 18.	May 3.	May 7.	May 7.	April 11.
1772	Jan. 14.	Feb. 12.	March 22	May 22.	March 25	Middlefex, April 13.		Middlefex, April 8.	Middlefex, April 13.	May 13	May 3.	Middlefex, April 13.	May 11.	May 3.	May 26.	March 3.	Feb. 19.		Kent, April 15.	Surrey, April 26.	May 11.	May 17.	March 25.	April 17.	Middlefex, April 18.	May 5.	
1773	Jan. 11.	Jan. 26.	March 21.	May 6.	March 18.	April 10.	March 29.	April 6.	April 6.	April 23.	April 29.	April 16.		April 10.	May 20.	March 3.	March 3.	April 10.	April 16.	April 22.	Essex, May 20.	June 2.	March 3.	March 24.	April 21.	April 19.	April 8.
1774	Jan. 28	Herts. Feb. 17.	Herts. March 4.	May 10.	March 31.	April 3	March 31.	April 3.	March 31.	April 25.	April 24.	April 8.		April 16.	April 27.			April 11.	April 21.	April 26.	April 30.	May 6.	Herts. March 4	April 6.	April 8.	April 24.	March 26.
1775	Jan. 14.	Feb. 9.	Feb. 26.	May 1.	March 5.	March 25.	March 20	March 19.	March 21.	April 21.	April 25.	Suffolk, March 30.	April 30.	Cambridgefh. March 31.	May 2.	Feb. 15.	Feb. 26.	April 15.	Suffolk, April 14.	April 23.	May 7.	May 15.	Feb. 26.	March 15	Cambridgefh. March 31.	Herts. April 7.	March 9.
1776	1775 Dec. 30.	Feb. 21.	Herts. March 8.	April 28.	March 22.	March 26.	March 19.	March 19.	March 23.	April 15.	April 19.	March 30.	April 21.	April 2.	April 23.	Feb. 14.	March 13.	April 2.	April 8.	April 21.	April 17.	May 26.	March 22.	April 7.	April 12.	April 6.	April 3.
1777	Jan. 17.	Feb. 25.	March 20.	May 6.	March 25.	March 28.	March 26.	April 7.	March 29.	April 22.	April 22.	March 30.	April 25.	April 11.	May 4.	Feb. 28.	Feb. 26.	April 6.	April 13.	April 20.	May 8.	May 20.	Feb. 27.	April 7.	April 10.	April 22.	
1778	Jan. 26.	Feb. 9.	April 4.	Herts. May 14.	March 19.	April 6.	April 6.	April 12.	April 7.	April 30.	April 15.	April 11.	April 17.	April 12.	April 25.	Feb. 9.	March 5.	April 12.	April 21.	April 30.	May 5.	May 27.	March 18.	April 14.		April 13.	April 5.
1779	1778 Dec. 24.	Feb. 6.	Feb. 22.	April 16.	Feb. 25.	March 7.	March 4.	March 4.	March 5.	March 31.	April 5.	March 25.	April 5.	March 24.	April 2.	Feb. 25.	March 1.	April 4.	April 13.	April 25.	May 9.	May 19.	Feb. 18.	Feb. 28.	April 1.	March 31.	March 15.
1780	Feb. 9.	Feb. 16.	March 15.	May 10.	March 21.	March 25.	March 28.	March 26.	March 28.	April 26.	April 24.	March 22.	April 26.	March 29.	May 1.	March 8.	March 1.	April 6.	April 22.	April 22.		April 30.	March 8.	April 30.	April 2.	April 22.	March 29.
1781	Jan. 29.	Feb. 6.	March 9.	April 28.	March 16.	March 22.	March 24.	March 26.	March 24.	April 19.	April 15.	April 10.	April 20.	April 3.	April 19.	Feb. 9.	March 1.	April 6.	April 11.	April 16.	April 24.	April 29.	March 9.	April 7.	April 6.	April 15.	March 26.
1782	Jan. 4.	Feb. 21.	Feb. 20.	May 19.	March 13.	April 3.	April 10.	April 10.		May 14.	April 25.	April 10.	May 11.	April 22.	May 16.	Feb. 27.	Feb. 27.	April 10.	April 22.	April 22.	May 8.	May 27.	April 7.	April 10.	May 5.	April 27.	April 12.
1783	Jan. 26.	Jan. 31.	Feb. 23.	April 27.	March 10.	April 7.	April 1.	April 2.	April 1.	April 24.	April 19.	April 6.	April 19.	April 14.	April 20.	Feb. 14.	March 9.	April 12.	April 20.	April 28.	April 12.	May 15.	March 18	March 23.	April 2.	April 18.	April 1.
1784	Feb. 8.	March 4.	April 22.	May 14.	April 18.	April 23.	April 23.	May 6.	April 22.	May 9.	May 1.	April 23.	May 9.	April 27.	May 12.	March 7.	March 2.	April 9.	April 19.	April 26.	April 28.	May 16.	April 17.	May 14.	May 4.	April 27.	April 22.
1785	Jan. 24.	Jan. 24.	April 12.	May 2.	April 8.	April 19.	April 18.	April 19.	April 18.	May 5.	April 23.	April 19.	April 29.	April 19.	May 2.	March 18.	March 14.	April 16.	April 12.	April 27.	April 23.	May 18.	March 19.	April 27.	April 22.	April 24.	April 16.
1786	Jan. 22.	Feb. 15.	March 26.	May 15.	March 19.	April 18.	April 16.	April 19.	April 16.	May 6.	April 28.	April 18.	May 6.	April 18.	May 2.	March 12.	Feb. 3.	April 13.	April 21.	April 30.	April 22.	June 4.	March 12.	April 25.	April 20.	April 21.	April 11.
1787	Jan. 30.	Feb. 4.	Feb. 25.	May 1.	March 21.	March 22.	March 18.	March 18.	March 17.	April 26.	April 25.	March 21.	April 24.	March 18.	April 29.	March 9.	March 7.	April 15.	April 18.	May 2.	April 27.	June 4.	Feb. 18.	March 22.	March 29.	April 26.	March 16.
1788	Jan. 12.	Feb. 12.	March 2.	May 6.	March 24.	April 4.	April 3.	April 10.	April 4.	April 24.	April 18.	April 4.	April 20.	April 4.	April 20.	March 11.	Feb. 26.	April 14.	April 11.	April 21.	April 27.	May 23.	March 28.	April 13	April 11.	April 12.	March 30.

Year	Population	Area	Notes
1850	1000	100	
1860	2000	200	
1870	3000	300	
1880	4000	400	
1890	5000	500	
1900	6000	600	
1910	7000	700	
1920	8000	800	
1930	9000	900	
1940	10000	1000	
1950	11000	1100	
1960	12000	1200	
1970	13000	1300	
1980	14000	1400	
1990	15000	1500	
2000	16000	1600	
2010	17000	1700	
2020	18000	1800	

XIV. *An Account of a Monster of the human Species, in two Letters; one from Baron Reichel to Sir Joseph Banks, Bart. and the other from Mr. James Anderson to Baron Reichel. Communicated by Sir Joseph Banks, Bart. P. R. S.*

Read April 30, 1789.

TO SIR JOSEPH BANKS, BART.

S I R,

Fort St. George, Feb. 28, 1788.

I HAVE the pleasure to transmit to you the portrait of a Gentoo boy, an astonishing living subject, who being sent to me by a friend of mine residing in the environs of the native place of the boy, I made two drawings representing the alternate attitudes in which he can place half the body of his little brother, who adheres to his breast. See Tab. II. PERUNTALOO is a handsome well-made lad, possessing every due faculty of mind and body, rather more sagacious, and with a superior share of understanding, than young men in general of his age. In addition to the inclosed anatomical description of the boy by Mr. ANDERSON, you will observe in the drawings two circular dotted lines, about the lower part of the loins of the semi-monster. During the several fittings I had of PERUNTALOO, I observed an internal motion about these parts rather more conspicuous than any other of the body; and upon questioning the youth, he shewed me, that by retaining his breath, he could force a current of air into them, so as to swell the parts like two blown-up bladders, with a rumbling noise at the time of action. Whether there is a connection with the lungs of PERUNTALOO is

Vol. LXXIX. C c a question.

a question I cannot venture to determine; Mr. ANDERSON, however, thinks it well worth my mentioning this observation. The erection of the little penis in the semi-monster, and the command PERUNTALOO has of discharging the urine through it, are perfectly ascertained.

Such as this subject is, if with any merit, you may depend upon the correctness of the drawings.

I am, &c.

T. REICHEL.

TO BARON REICHEL.

S I R,

Fort St. George, Feb. 25, 1788.

AS you mean to send the elegant drawing of PERUNTALOO to Sir JOSEPH BANKS, you may acquaint him from me, that the little brother is suspended by the os pubis; an elongation of the sword-like cartilage of PERUNTALOO having anastomosed with that bone at the symphysis.

The lower orifice of the stomach seems to lie in the sac or cylindrical cavity between the two brothers on the right-side, and what may be reckoned the right hypochondre of the little one, as that part is tumid and full after eating.

The alimentary canal must be common to both, as the anus of the little one is imperforate.

There is a bladder of urine distinctly perceived, which occupies the left side of the sac, or left hypochondre of the monster.


Besides which, there remain only the sacrum, ossa innominata, and lower extremities perfect.


PERUNTALOO



Fig. 1.

Fig. 2.





PERUNTALOO says he has as complete a sense of feeling with every part of the body of his little brother as of his own proper body, and this may account for the erections you saw, and making water distinctly; but this volition does not extend to the legs or feet, which are cold in comparison with the rest.

I am, &c.

JAMES ANDERSON.

EXPLANATION OF THE PLATE.

PERUNTALOO, son of CHINDRAHPAH-NAYANDOO, of the Gentoo Cast. He was born at Popelpahdoo, 70 miles west from Mufilipatnam. He is 13 years of age, and measures 4 feet 6½ inches in height.

Fig. 1. Natural position.

Fig. 2. Reversed position.



XV. *A supplementary Letter on the Identity of the Species of the Dog, Wolf, and Jackal; from John Hunter, Esq. F. R. S. addressed to Sir Joseph Banks, Bart. P. R. S.*

Read April 30, 1789.

S I R,

I N the year 1787 I had the honour of presenting to this learned Society, a Paper to prove the Wolf, the Jackal, and the Dog to be of the same species. But as the complete proof of the Wolf being a Dog, which consisted in the half-bred puppy breeding again, had not been under my own inspection, although sufficiently well authenticated, I saved a female of one of the half-bred puppies, mentioned in that Paper, in hopes of being myself a witness of the fact; but when the period of impregnation arrived, we unluckily missed that opportunity. However, another half-bred puppy has had young, which is equally satisfactory to me as if my own had bred. JOHN SYMMONS, Esq. of Milbank, has had a female Wolf in his possession for some time, who was lined by a Dog, and brought forth several puppies, which I had the honour of seeing with you. This was a very short time after the brood had been produced by Mr. GOUGH's Wolf, the subject of my former Paper, therefore the puppies were nearly of an age with mine. These puppies Mr. SYMMONS has reared; only one of them was a female, and she had much more of the mother or Wolf in her than any of  
the

the rest of the same litter. I communicated my wish to Mr. SYMMONS, that either his puppy or mine should prove the fact to our own knowledge; which he immediately, with great readiness, acceded to. On the 16th, 17th, and 18th of December, 1788, this bitch was lined by a Dog, and on the 18th of February she brought eight puppies, all of which she now rears. If we reckon from the 16th of December, she went sixty-four days; but if we reckon from the 17th, the mean time, then it is sixty-three days, the usual time for a bitch to go with pup. These puppies are the second remove from the Wolf and Dog, similar to that given by my Lord CLANBRASSIL to the Earl of PEMBROKE, which bred again. (See Philosophical Transactions, Vol. LXXVII. p. 255.) It would have proved the same fact if she had been lined by either a Wolf, a Dog, or one of the males of her own litter.

I may just remark here, that the Wolf seems to have only one time in the year for impregnation natural to her, and that is in the month of December; for every time Mr. GOUGH's Wolf has been in heat was in this month, and it proves to be the same month in which Mr. SYMMONS's Wolf was in heat; for his half-bred Wolf is nearly of the same age with mine, and the time she was in heat was also the same with that of her own mother, and the present brood corresponds in time with the brood of Mr. GOUGH's Wolf.

I am, &c.

JOHN HUNTER.



XVI. *Abstract of a Register of the Barometer, Thermometer, and Rain, at Lyndon in Rutland; by Thomas Barker, Esq. Also of the Rain in Hampshire and Surrey. Communicated by Thomas White, Esq. F.R.S.*

Read April 30, 1789.

1788.

		Barometer.			Thermometer.						Rain.			
		Highest	Lowest	Mean.	In the House.			Abroad.			Lyndon	Surrey. S. Lambeth.	Hampshire.	
					Inch.	Inch.	Inch.	High.	Low.	Mean.			High.	Low.
					°	°	°	°	°	°	Inch.	Inch.	Inch.	Inch.
Jan.	Morn.	30,13	28,37	29,50	44	34	40	45	23 $\frac{1}{2}$	36	0,970	0,68	1,60	1,10
	Aftern.				44	35	40 $\frac{1}{2}$	49	30	41				
Feb.	Morn.	29,77	28,25	29,14	45	35	40 $\frac{1}{2}$	44	27 $\frac{1}{2}$	36	2,667	2,09	3,37	2, 6
	Aftern.				45	37	41	48	30	42				
Mar.	Morn.	29,65	28,84	29,23	51	34	40	50	22	35	1,072	0,64	1,31	1,36
	Aftern.				52 $\frac{1}{2}$	35 $\frac{1}{2}$	41	63	31	43				
Apr.	Morn.	30,02	28,94	29,59	56	42 $\frac{1}{2}$	50	54	35	45 $\frac{1}{2}$	0,588	0,47	0,61	0,50
	Aftern.				60	43 $\frac{1}{2}$	51	68 $\frac{1}{2}$	40	56				
May	Morn.	29,92	29,19	29,60	68 $\frac{1}{2}$	51	58	64	43 $\frac{1}{2}$	52 $\frac{1}{2}$	1,517	0,81	0,76	0,28
	Aftern.				72	53	60	82	51	66				
June	Morn.	29,85	29,10	29,52	65 $\frac{1}{2}$	56	60	64 $\frac{1}{2}$	50	56	0,608	1,94	1,27	1,36
	Aftern.				69	57 $\frac{1}{2}$	61	82 $\frac{1}{2}$	58	67				
July	Morn.	29,78	29,21	29,52	67 $\frac{1}{2}$	58	62	70 $\frac{1}{2}$	51	59	1,795	1,84	3,58	1,81
	Aftern.				70	59 $\frac{1}{2}$	63 $\frac{1}{2}$	83	58	72				
Aug.	Morn.	30,01	28,88	29,49	68	57 $\frac{1}{2}$	61 $\frac{1}{2}$	64	54	56	2,780	4,30	3,22	3,40
	Aftern.				70 $\frac{1}{2}$	59	63	77	62	68				
Sept.	Morn.	29,80	29,00	29,40	66	52 $\frac{1}{2}$	58 $\frac{1}{2}$	61 $\frac{1}{2}$	42	52	2,430	3,81	5,71	3,78
	Aftern.				66 $\frac{1}{2}$	53	59	75 $\frac{1}{2}$	50	63				
Oct.	Morn.	30,15	29,15	29,68	59	46	52	57	32	46	1,412	0,08	0, 0	0,03
	Aftern.				60	47	53	66	45	54 $\frac{1}{2}$				
Nov.	Morn.	30,01	29,06	29,62	53	37	45	51 $\frac{1}{2}$	25 $\frac{1}{2}$	39	0,453	0,62	0,86	0,74
	Aftern.				53	37 $\frac{1}{2}$	46	58	31	45				
Dec.	Morn.	29,85	29,12	29,47	39	27	34	40 $\frac{1}{2}$	15	27	0,890	0,00	0,21	0,42
	Aftern.				40 $\frac{1}{2}$	28	34 $\frac{1}{2}$	44 $\frac{1}{2}$	22 $\frac{1}{2}$	31 $\frac{1}{2}$				
Inches											17,182	17,28	22,50	16,84

THE year began open and mild, at first showery, afterward drier and stormy. The chief part of February was wet, but more so for frequency than quantity. After a few misty days there came in March above a fortnight's sharp frost, the longest this winter, and with severe east winds cut things more than all the winter before, which was in general an open one. The last twelve or thirteen days of March the spring set in pleasant, and continued forwarding all April, and proved a very dry spring. There were at times this year fits of exceeding hot weather, the end of April, the fourth week in May, the third in June, and second in July; but so much windy weather, with hot sun and cold winds, that bees which were forward the beginning of May, and some few swarms so early, seemed backward again at the end of the month. The grass was every where short, and began to burn; but a fine rain at the end of May strengthened the grain very much, and made the grass grow in some degree; but it soon began to burn again in a dry June, with almost constant north and north-east winds, so that the pasture was short, and very little hay.

The end of June, and two-thirds of July, were very frequent showers and wind. There were in some parts of England very heavy thunder-storms, and more rain than they wished for in hay time. The showers were light here; they made the grain ear well, pease and beans set thick, and brought the turnips past the fly. The grass also grew in some degree, but burnt again before July was out, and more in August, of which the first ten days were dry; but the showery latter part made the grass grow considerably, which was much wanted, and did not much hinder the harvest, which was in general well got, and was good. The autumn was very fine, and so much rain in one month, especially the third week in September, that there

was more grafs after that than there had been any part of the summer before, though not fuch quantities as there fometimes is; for, take the year throughout, I think I never knew lefs. But that was not the cafe in all parts of England: I believe it was in general a dry summer every where; but in fome places there was a great deal of grafs at times. So great a fruit year of moft forts, garden, orchard, and wild, I think, I hardly ever knew. After the first week in October it was dry again, and fo fine, mild, and clear of frosts, that the nasturtiums were not cut off till after the middle of November; and the ground and roads continued dry till the snow at Christmas, and there was in many places great want of water fo late in the year. Moft part of the last week in November, and the first third part of December, was a gentle frost; but then it fet in very fevere, and, except an imperfect thaw the 24th and 25th, has been an uncommonly cold and hard frost, freezing over many of the rivers, with a confiderable snow at times, chiefly the 26th and 27th, and continued to the end of the year, and beyond it.

*Account of a sinking-in of the ground.*

In a wet season, about Christmas 1787, a piece of apparently found ground on the north fide of a moderate hill, a mile and half south-west from Ketton in Rutland, funk down into the earth, leaving a great hollow. The ground was smooth before, and a waggon had lately gone over the place. There was nobody by when it fell in; but a labourer going home from his work was the first person who found it.

It was some time after the accident before I heard of it, and  
it



it was in spring time that I went to see it. I then found it to be an oval hole, five yards over one way, and four another, and about four yards deep in the middle; but some of the earth having lodged against the sides of the pit, it was not so deep there; yet the oval must upon the whole have sunk down about three yards, and gone directly downward, for the sides of the pit are left perpendicular. I found a little water at the bottom of the pit, and was told there had been a great deal more at first. The bottom half of the pit is a blue clay, and from a foot to a yard thick at the top is a stiff earth mixed with stones. There were plain signs that a drain from the ground above had in wet times run down near where the pit now is; some of it probably ran into and under the ground, and had, in a course of time, undermined it; and that seems to have been the reason that the pit sunk in as it has done.

A man of Ketton, who has freestone pits in the same lordship, but on the opposite side of the town, says, he sometimes meets with beds of clay in his pits, which are undermined, and have hollows in them. And to the northward of these stone pits there are many hollows, which they call the Swallow-pits; because, being hollow underneath, no water will lie in them, but runs through holes into the ground. These swallow-pits I know, and they seem to be clay at top; and he says, they do not appear to have been ever dug by men, but that the surface of the ground has sunk down into the hollow there was beneath it.



XVII. *On the Method of correspondent Values, &c.* By Edward Waring, M. D. F. R. S. and Lucasian Professor of the Mathematics at Cambridge.

Read May 28, 1789.

I.

1. **I**N the year 1762 I published a method of finding when two roots of a given equation  $x^n - px^{n-1} + qx^{n-2} - rx^{n-3} + \&c. = 0$  are equal, by finding the common divisors of the two quantities  $a^n - pa^{n-1} + qa^{n-2} - \&c.$ , and  $na^{n-1} - \overline{n-1}pa^{n-2} + \overline{n-2}qa^{n-3} - \&c.$ , and observed if they admitted only one simple divisor  $(a - A)$ , then two roots were only equal; if a quadratic  $(a^2 - Aa + B)$ , then two roots of the equation became twice equal; if a cubic  $(a^3 - Aa^2 + Ba - C)$ , then two roots became thrice equal; and so on: or, to express in more general terms what follows from the same principles, if the common divisor be  $\overline{a-b} \times \overline{a-c} \times \overline{a-d} \times \&c.$ , then  $r + 1$  roots of the given equation will be  $b$ ,  $s + 1$  roots will be  $c$ ,  $t + 1$  will be  $d$ , &c.; and it immediately follows, from the principles delivered in the second edition of the same Book, published in 1770, that to find when  $r + 1$ ,  $v + 1$ ,  $t + 1$ , &c. roots are respectively equal requires  $r + s + t$ , &c. equations of condition, which are deducible from the well known method of finding the common divisors of two quantities in this case of  $a^n - pa^{n-1} + qa^{n-2} - \&c.$ ,  $na^{n-1} - \overline{n-1}pa^{n-2} + \overline{n-2}qa^{n-3} - \&c.$  of the terms of their remainders, &c.

In



observed, in the before-mentioned Book, that (if the common divisor be  $(a - A)$ ) it will once only admit of 3, 4, 5, . . .  $r$  equal roots; if it be a quadratic, then it will twice admit of those equal roots; and so on.

4. If the roots of the equation of the least dimensions be substituted for  $a$  in the remaining equations, and each of the resulting values of the same equation be multiplied into each other, there will result the  $r - 1$  equations of condition: and the same may be deduced also from the several equations conjointly.

The equations of conditions found by the first method, if the divisions were not properly instituted, may admit of more rational divisors than necessary, of which some are the equations of conditions required.

## 2.

1. In the year 1776, I published in the *Meditationes Analyticæ* a new method of differences for the resolution of the following problem.

Given the sums of a swiftly converging series  $ax + bx^2 + cx^3 + dx^4 + \&c.$ , when the values of  $x$  are respectively  $\pi, \rho, \sigma, \&c.$ ; to find the sum of the series when  $x$  is  $\tau$ , that is, given  $S\pi = a\pi + b\pi^2 + c\pi^3 + d\pi^4 + \&c.$ ,  $S\rho = a\rho + b\rho^2 + c\rho^3 + \&c.$ ,  $S\sigma = a\sigma + b\sigma^2 + c\sigma^3 + \&c. \&c.$ ; to find  $S\tau = a\tau + b\tau^2 + c\tau^3 + \&c.$

To resolve this problem I multiplied the quantities,  $S\pi, S\rho, S\sigma, \&c.$  respectively into unknown co-efficients  $\alpha, \beta, \gamma, \&c.$  and there resulted

$$\alpha\pi a + \alpha\pi^2 b + \alpha\pi^3 c + \&c.$$

$$\beta\rho a + \beta\rho^2 b + \beta\rho^3 c + \&c.$$

$$\gamma\sigma a + \gamma\sigma^2 b + \gamma\sigma^3 c + \&c.$$

$$\&c. \quad \&c. \quad \&c.$$

and then made the sum of each of the terms respectively equal to its correspondent term of the quantity  $\tau a + \tau^2 b + \tau^3 c + \&c.$ , and consequently  $\alpha\pi + \beta\rho + \gamma\sigma + \&c. = \tau$ ,  $\alpha\pi^2 + \beta\rho^2 + \gamma\sigma^2 + \&c. = \tau^2$ ,  $\alpha\pi^3 + \beta\rho^3 + \gamma\sigma^3 + \&c. = \tau^3$ , &c. I assumed as many equations of this kind as there were given values  $\pi, \rho, \sigma, \&c.$  of  $x$ ; and consequently as many equations resulted as unknown quantities  $\alpha, \beta, \gamma, \&c.$ ; whence, by the common resolution of simple equations, or more easily from differences, can be found the unknown quantities  $\alpha, \beta, \gamma, \&c.$ , and thence the equation sought  $\alpha \times S\pi + \beta \times S\rho + \gamma \times S\sigma + \&c. = S\tau$  nearly.

3. In the Meditations are assumed for  $\pi, \rho, \sigma, \&c.$  the quantities  $p, 2p, 3p, 4p, \dots, \overline{n-2p}, \overline{n-1p}$ , and  $np$  for  $\tau$ ; which, if substituted for their values in the preceding equations, will give  $\alpha + 2\beta + 3\gamma + 4\delta + \&c. = n$ ,  $\alpha + 4\beta + 9\gamma + 16\delta + \&c. = n^2$ ,  $\alpha + 8\beta + 27\gamma + \&c. = n^3$ ,  $\alpha + 16\beta + 81\gamma + \&c. = n^4$ ; and if the sums of the series  $ax + bx^2 + cx^3 + \&c.$  which respectively correspond to the values  $p, 2p, 3p, \dots, \overline{n-1p}$  of  $x$  be  $S_1, S_2, S_3, S_4, \dots, \overline{S_{n-1}}$ , and the sum of the series  $ax + bx^2 + cx^3 + \&c.$  which corresponds to  $n$  value of  $x$  be  $S_n$ ; then will  $S_n = n\overline{S_{n-1}} - n \cdot \frac{n-1}{2} \overline{S_{n-2}} + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{2} \overline{S_{n-3}} \dots \pm n S_1$  nearly, which equation is given in the above-mentioned Book.

3. The logarithm from the number, the arc from the sine, &c. are found by serieses of the formula  $ax + bx^2 + cx^3 + \&c.$ ; and consequently this equation is applicable to them.

6.

4. In

4. In the same Book is assumed a series  $ax^r + bx^{r+1} + cx^{r+2} + dx^{r+3} + \&c.$  of a more general formula than the preceding, and in it for  $x$  substituted  $\alpha, \beta, \gamma, \delta, \&c., m$ ; and  $S\alpha, S\beta, S\gamma, S\delta, \&c.$ ;  $S_m$  for the resulting sums, and thence deduced

$$S_m = \frac{m^r \times m^s - \beta^s \cdot m^s - \gamma^s \cdot m^s - \delta^s \cdot \&c.}{\alpha^r \times \alpha^s - \beta^s \cdot \alpha^s - \gamma^s \cdot \alpha^s - \delta^s \cdot \&c.} \times S\alpha + \frac{m^r \times m^s - \alpha^s \cdot m^s - \gamma^s \cdot m^s - \delta^s \cdot \&c.}{\beta^r \times \beta^s - \alpha^s \cdot \beta^s - \gamma^s \cdot \beta^s - \delta^s \cdot \&c.} \\ \times S\beta + \frac{m^r \times m^s - \alpha^s \cdot m^s - \beta^s \cdot m^s - \delta^s \cdot \&c.}{\gamma^r \times \gamma^s - \alpha^s \cdot \gamma^s - \beta^s \cdot \gamma^s - \delta^s \cdot \&c.} \times S\gamma + \frac{m^r \times m^s - \alpha^s \cdot m^s - \beta^s \cdot m^s - \gamma^s \cdot \&c.}{\delta^r \times \delta^s - \alpha^s \cdot \delta^s - \beta^s \cdot \delta^s - \gamma^s \cdot \&c.} \\ \times S\delta + \&c. \text{ nearly.}$$

*Cor.* If for  $r$  and  $s$  be assumed respectively 1, the series becomes  $ax + bx^2 + cx^3 + \&c.$  of the same formula as the preceding: if  $r=0$  and  $s=1$ , the series becomes  $a + bx + cx^2 + \&c.$  The latter case will be the same as the former, when one of the quantities ( $\alpha$ ) substituted for  $x$  and its correspondent sum  $S\alpha$ , both become  $=0$ , and the equation deduced in both cases the same.

5. If  $\pi, \rho, \sigma, \&c.$  respectively denote  $r, r+p, r+2p, \dots r+n-2p, r+n-1p$ , and  $\tau = r+np$ ; and  $S, S_1, S_2, S_3, \dots S_{n-2}, S_{n-1}$ , be the sums either resulting from the series  $ax + bx^2 + cx^3 + \&c.$  or the series  $A + ax + bx^2 + cx^3 + \&c.$ , which respectively correspond to the values  $r, r+p, r+2p, \&c.$  of  $x$ ; and  $S_n$  the sum of the same series which corresponds to the value  $r+np$  of  $x$ ; then will  $S_n = nS_{n-1} - n \cdot \frac{n-1}{2} S_{n-2} + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} S_{n-3} - \dots \pm n \cdot \frac{n-1}{2} S_2 \mp nS_1 \pm S$  nearly; this equation differs from the preceding by the last term  $S$  not vanishing; in the preceding case  $S$  became  $=0$ , for it was the sum of the series  $ax + bx^2 + cx^3 + \&c.$ , which corresponded to  $x=0$ .

6. From

6. From the Meditations it appears, that  $r^m - n \times r \pm p^m + n \cdot \frac{n-1}{2} r \pm 2p^m - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} r \pm 3p^m + \&c.$  to the end of the series = 0, if  $m$  is less than  $n$ , and  $m$  and  $n$  are whole numbers; but if  $m = n$ , then it will =  $\pm 1 \cdot 2 \cdot 3 \cdot 4 \dots n-1 \cdot np^m$ ; whence it is manifest, that for the  $n$  first terms of the series  $A + ax + bx^2 + \&c.$  the equations are true; and for the  $n-1$  first terms of the series  $ax + bx^2 + cx^3 + \&c.$  and in the successive term of both the serieses they will err by a quantity nearly =  $\pm 1 \cdot 2 \cdot 3 \dots n \times p^n \times r^{-n} \times$  co-efficient of the term; and the errors of every subsequent term ( $x^{4+n}$ ) will be nearly as  $\pm m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3} \cdot \frac{m-3}{4} \dots \frac{m-b+1}{b} \times p^n \times r^{-n} \times$  co-efficient of the term  $x^{b+n}$ , if for  $r, r+p, r+2p, \&c.$  be substituted  $1, 1 + \frac{p}{r}, 1 + \frac{2p}{r}, \&c.$

7. Let the preceding equation  $S_n = n S_{n-1} - n \cdot \frac{n-1}{2} S_{n-2} + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot S_{n-3} - \&c. = n \times \log. r - p - n \cdot \frac{n-1}{2} \log. r - 2p + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \log. r - 3p + \&c. = \log. \frac{r \times r - 2p^s \times r - 4p^{s'} \times r - 6p^{s''} \times \&c.}{r - p^t - r - 3p^{t'} \times r - 5p^{t''} \times \&c.} = \log. K$ , where  $s, s', s'', \&c.$  denote the co-efficients of the alternate terms of the binomial theorem, viz.  $s = n \cdot \frac{n-1}{2}, s' = n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4}, \&c.$ , and  $t = n, t' = n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}, \&c.$  the co-efficients of the remaining alternate terms; the numerator  $r \times r - 2p^s \times r - 4p^{s'} \times r - 6p^{s''} \times \&c. = (\text{if } N = 2^{n-1}) r^N - Ppr^{N-1} + Qp^2r^{N-2} - Rp^3r^{N-3} \dots Lp^{n-1} \times r^{N-n+1} \pm Mp^n r^{N-n} \mp \&c.$ ; and the denominator  $r - p^t \times r - 3p^{t'} \times r - 5p^{t''} \times \&c. = r^N - Pp r^{N-1} + Qp^2 r^{N-2} -$

L Rp

$Rp^3r^{n-3} + \dots Lp^{n-1}r^{N-n+1} (\pm M + 1 \cdot 2 \cdot 3 \dots \overline{n-1}) p^n r^{N-n} \mp$   
 &c., whence the numerator and denominator have the  $n$  first  
 terms the same, and the next succeeding terms differ by  
 $1 \cdot 2 \cdot 3 \dots \overline{n-1} p^n r^{N-n}$ ; the numerator divided by the denomi-  
 nator =  $1 \pm \frac{1 \cdot 2 \cdot 3 \dots \overline{n-1}}{r^n} p^n$  nearly, if  $r$  be a great number in  
 proportion to  $p$ , &c. it would be  $+$  when  $n$  is an odd number,  
 and  $-$  when even.

8. The logarithm of the fraction  $K$  by the common series  
 $= K - 1 - \frac{\overline{K-1}^2}{2} + \frac{\overline{K-1}^3}{3} - \&c.$  has for its first term =  $\pm$   
 $\frac{1 \cdot 2 \cdot 3 \dots \overline{n-1}}{r^n} \times p^n$  nearly; for its second term the square of  
 the first divided by 2, &c.

9. The error of this equation not only depends on the loga-  
 rithm of  $K$ , which may be calculated to any degree of exact-  
 ness, but in the calculus on the errors of the given loga-  
 rithms.

10. If  $r$  be increased or diminished by any given number,  
 the  $n$  first terms of the numerator and denominator will still  
 result the same, and the next succeeding terms will differ by  
 $1 \cdot 2 \cdot 3 \cdot 4 \dots n-1 \times p^n \times r^{N-n}$ .

11. Let  $n \cdot \frac{n-1}{2}$  numbers be 2,  $n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4}$  num-  
 bers be 4,  $n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} \cdot \frac{n-4}{5} \cdot \frac{n-5}{6}$  numbers be 6, &c.;  
 their sum, the sum of the products of every two, the con-  
 tents of every three, four, five, &c. to  $n-1$  of them will be  
 equal to the sum, the sum of the products of every two, of  
 the contents of every three, four, five, &c. to  $\overline{n-1}$  of the  
 following numbers, *viz.*  $n$  numbers which are 1,  $n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3}$   
 numbers



numbers which are  $3, n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \cdot \frac{n-3}{4} \cdot \frac{n-4}{5}$ , which are 5, &c.; and the sum of the contents of every  $n$  of the former will be less than the sum of the contents of every  $n$  latter numbers by  $1 \cdot 2 \cdot 3 \cdot 4 \dots \overline{n-1}$ .

12. The method given in Art. 4. which I name a method of correspondent values, easily deduces and demonstrates the preceding equations, which cannot, without much difficulty, be done by the preceding method of differences; the method of correspondent values is much preferable to the method of differences, both for the facility of its deduction, and the generality of its resolution: for instance, from this method very easily can be deduced, &c. the subsequent and other similar equations.

Ex. 1.  $S_n = n \overline{S_{n-1}} - n \cdot \frac{n-1}{2} \overline{S_{n-2}} + n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \overline{S_{n-3}} \dots$   
 $\approx n S_1 = S$  nearly.

Ex. 2.  $S_{n+m} = \frac{\overline{m+n} \cdot \overline{m+n-1} \cdot \overline{m+n-2} \dots \overline{m+2}}{1 \cdot 3 \dots n-1} \times \overline{S_{n-1}} -$   
 $\frac{n-1}{1} \times A \times \frac{m+1}{m+2} \times \overline{S_{n-2}} + \frac{n-2}{2} \times B \times \frac{m+2}{m+3} \overline{S_{n-3}} - \frac{n-3}{3} \times C \times$   
 $\frac{m+3}{m+4} \times \overline{S_{n-4}} + \frac{n-4}{4} \times D \times \frac{m+4}{m+5} \times \overline{S_{n-5}} - \&c.$  nearly, where the letters A, B, C, D, &c. denote the preceding co-efficients, and the converging series is the same as in the preceding example.

Ex. 3. Let the converging series be of the formula  $ax + bx^3 + cx^5 + dx^7 + \&c.$ ; then will  $S_n = \overline{2n-2} \overline{S_{n-1}} - \overline{2n-1} \times$   
 $\frac{2n-4}{2} \overline{S_{n-2}} + \overline{2n-1} \times \frac{2n-2}{2} \times \frac{2n-6}{3} \overline{S_{n-3}} - \overline{2n-1} \cdot \frac{2n-2}{2} \cdot$   
 $\frac{2n-3}{2} \times \frac{2n-8}{4} \overline{S_{n-4}} + \&c.$  nearly, of which the general term is  
 $\overline{2n-1} \cdot \frac{2n-2}{2} \cdot \frac{2n-3}{3} \dots \frac{2n-l+1}{l-1} \times \frac{2n-2l}{l} \times \overline{S_{n-l}}$ .

Ex. 4. Let the series be of the formula  $A + ax^2 + bx^4 + cx^6 + \&c.$ ; then will  $S_n = \frac{n-1}{n} \times \frac{2n-2}{2n-1} S_{n-1} - \frac{n-2}{n} \times \frac{2n-4}{2n-1} \cdot \frac{2n-4}{2} \times S_{n-2} + \frac{n-3}{n} \times \frac{2n-2}{2n-1} \cdot \frac{2n-2}{2} \cdot \frac{2n-6}{3} S_{n-3} - \frac{n-4}{n} \times \frac{2n-4}{2n-1} \cdot \frac{2n-2}{2} \cdot \frac{2n-3}{3} \times \frac{2n-8}{4} S_{n-4} + \&c.$  nearly, of which the general term is  $\frac{n-l}{n} \times \frac{2n-1}{2n-1} \cdot \frac{2n-2}{3} \cdot \frac{2n-3}{3} \dots \frac{2n-l+1}{l-1} \times \frac{2n-2l}{l} \times S_{n-l}.$

Ex. 5. Let the given series be of the formula  $ax + bx^2 + cx^3 + \&c.$ , and in it for  $x$  be substituted  $p, -p, 2p, -2p, 3p, -3p, \dots, np, -np$  and  $mp$ , and for the sums of the resulting series be wrote respectively  $S^1, S^{-1}, S^2, S^{-2}, S^3, S^{-3}, \dots, S^n, S^{-n}$ , and  $S^m$ ; then will  $S^m =$

$$\frac{m \cdot m^2 - 1 \cdot m^2 - 4 \cdot m^2 - 9 \cdot m^2 - 16 \dots m^2 - n - 1^2 \cdot m - n}{n \cdot n^2 - 1 \cdot n^2 - 4 \cdot n^2 - 9 \cdot n^2 - 16 \dots n^2 - n - 1^2 \times 2n - 1 \cdot 2 \cdot 3 \cdot 4 \dots 2n} \times S^{-n} + A \times \frac{m+n}{m-n} S^{+n} - \frac{2n}{1} \times B \times \frac{m-n}{m+n-1} S^{-n+1} - C \times \frac{m+n-1}{m-n-1} \times S^{n-1} + \frac{2n-1}{2} \times D \times \frac{m-n-1}{m+n-2} \times S^{-n+2} + \frac{m+n-2}{m-n-2} \times E S^{n-2} - \frac{2n-2}{3} \times F \times \frac{m-n-2}{m+n-3} S^{-n+3} - G \times \frac{m+n-3}{m-n-3} \times S^{n-3} + \&c. \text{ nearly, where}$$

the letters A, B, C, D, &c. respectively denote the preceding co-efficients. In general, the co-efficients of the terms  $S^{-n+s}$

and  $S^{n-s}$  will be respectively  $M = \frac{2n-s+1}{s} \times L \times \frac{m-n-s+1}{m+n-s}$  and

$M \times \frac{m+n-s}{m-n-s}$ , where the letters L and M respectively denote their

preceding co-efficients; the co-efficients are to be taken affirmatively, or negatively, according as  $s$  is an even or odd number.

Ex. 6. If for  $x$  in the preceding series be substituted  $p, -p, 2p, -2p, 3p, -3p, \dots, n-1p, -n-1p, np$  respectively, then

$S^m$

$$S^m = \frac{m \cdot \overline{m^2-1} \cdot \overline{m^2-4} \cdot \overline{m^2-9} \cdot \overline{m^2-16} \dots (m^2-(n-1)^2)}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \dots (2n-1)} S^n - A \times$$

$$\frac{m-n}{m+n-1} \times S^{-n+1} - \frac{2n-1}{1} \times B \times \frac{m+n-1}{m-n-1} \times S^{n-1} + C \times \frac{\overline{m-n-1}}{m+n-2} \times$$

$$S^{-n+2} + \frac{2n-2}{2} \times D \times \frac{m+n-2}{m-n-2} \times S^{n-2} - \&c. \text{ nearly, when } A, B, C,$$

&c. denote as before the preceding co-efficients. The coefficients of the terms  $S^{-n+s}$  and  $S^{n-s}$  will be respectively  $L \times \frac{\overline{m-n-s+1}}{m+n-s}$  and  $\frac{2n-1}{s} \times M \times \frac{\overline{m+n-s}}{m-n-s}$ , L and M denoting the preceding co-efficients, which are to be taken negatively or affirmatively, as  $s$  is an even or an odd number. In this series when  $x=0$ , the correspondent sum = 0.

Ex. 7. Let the given series be of the formula  $a + bx + cx^2 + dx^3 + \&c.$ ; and in it for  $x$  be substituted  $0, p, -p, 2p, -2p, 3p, -3p \dots np, -pp$  and  $mp$ , and for the sums of the resulting series be wrote as before  $S^0, S^1, S^{-1}, S^2, S^{-2}, \dots S^n, S^{-n}$ , and

$$S^m; \text{ then will } S^m = \frac{m \cdot \overline{m^2-1} \cdot \overline{m^2-4} \cdot \overline{m^2-9} \dots (m^2-n-1^2) \times \overline{m-n}}{n \cdot \overline{n^2-1} \cdot \overline{n^2-4} \cdot \overline{n^2-9} \dots (n^2-n-1^2) \times 2n}$$

$$\times S^{-n} + A \times \frac{m+n}{m-n} \times S^{+n} - \frac{2n}{1} \times B \times \frac{m-n}{m+n-1} S^{-n+1} - \&c. \text{ this series}$$

observes the same law as the series given in Ex. 5. and only differs from it by the last term  $S_0$  not vanishing, that is, being = 0.

Ex. 8. Let the series be of the preceding formula  $a + bx + cx^2 + dx^3 + \&c.$ , and in it for  $x$  be substituted  $0; p, -p; 2p, -2p; 3p, -3p; \dots n-1p, -n-1p, np$ , and  $mp$ , and the sums resulting be  $S_0, S_1, S^{-1}, S^2, S^{-2}, \dots S^{n-1}, S^{-n+1}, S^n$  and  $S^m$ ;

$$\text{then will } S^m = \frac{m \cdot \overline{m^2-1} \cdot \overline{m^2-4} \dots \overline{m^2-(n-1)^2}}{1 \cdot 2 \cdot 3 \cdot 4 \dots 2n-1} S^n - A \times \frac{m-n}{m+n-1} \times$$

$S^{-n+1} - \&c.$  the same series as in Ex. 6. and differs from it only by the last term  $S_0$  not vanishing.

Ex. 9. Let the series be of the same formula  $a + bx + cx^2 + dx^3 + \&c.$  and in it for  $x$  be substituted  $p, -p, 3p, -3p, 5p, -5p, 7p, -7p, \dots np, -np$  and  $mp$ ; and the sums resulting be  $S^1, S^{-1}, S^3, S^{-3}, S^5, S^{-5}, S^7, S^{-7}, \dots S^n, S^{-n}$ , and  $S^m$ ; then will  $S^m =$

$$\frac{\overline{m^2 - 1} \cdot \overline{m^2 - 9} \cdot \overline{m^2 - 25} \cdot \overline{m^2 - 49} \dots \overline{m^2 - n - 2^2} \times m + n}{n^2 - 1 \cdot n^2 - 9 \cdot n^2 - 25 \cdot n^2 - 49 \dots n^2 - n - 2^2} \times 2n = 2^n \times 1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \dots n$$

$$\times S^n - A \times \frac{m-n}{m+n} \times S^{-n} - \frac{n}{1} \times B \times \frac{m+n}{m-n+2} \times S^{n-2} + C \times \frac{m-n+2}{m+n-2} \times$$

$$S^{-n+2} + \frac{n-1}{2} \times D \times \frac{m+n-2}{m-n+4} \times S^{n-4} - E \times \frac{m-n+4}{m+n-4} S^{-n+4} - \frac{n-2}{3} F$$

$$\times \frac{m+n-4}{m-n+6} S^{n-6} + \frac{m-n+6}{m+n-6} \times G \times S^{-n+6} + \frac{n-3}{4} \times H \times \frac{m+n-6}{m-n+8} \times$$

$S^{n-8} - \&c.$  nearly, where the letters A, B, C, D, E, &c. denote the preceding co-efficients of the terms  $S^n, S^{-n}, S^{n-2}, S^{-n+2}, S^{n-4}, S^{-n+4}, S^{n-6}, \&c.$  respectively. The co-efficients of the terms  $S^{2n-2s}$  and  $S^{-n+2s}$  will be  $M = \frac{n-s+1}{s} \times L \times$

$$\frac{m+n-2s+2}{m-n+2s} \text{ and } N = M \times \frac{m-n+2s}{m+n-2s}; \text{ where } L, M, \text{ and } N \text{ denote}$$

the co-efficients of the terms immediately preceding each other, that is, of the terms  $S^{-n+2s-2}, S^{n-2s},$  and  $S^{-n+2s}$ . The sign of the first co-efficient M will be + or -, according as  $s$  is even or odd; the second term N will have a contrary sign to the first.

These series may be made to begin from any term, which may be easily found by the method of correspondent values, and the subsequent terms from it by the given law; its preceding terms may be deduced from the same law reversed, that is, by putting the numerators of the fractions multiplied into it for the denominators, and the denominators for the numerators.

From these different serieses may be formed, by adding two or more terms of the given series together for a term of the required series; which method has been applied to converging series in general in the Meditations.

13. The method of correspondent values easily affords a resolution of the problems contained in Mr. BRIGG's or Sir ISAAC NEWTON's method of differences.

Ex. 1. Let the quantity be of the formula  $a + bx + cx^2 + dx^3 + \&c. \dots x^n = y$ , and  $n + 1$  correspondent values of  $x$  and  $y$  be given, viz.  $p, q, r, s, \&c.$  of  $x$ ;  $Sp, Sq, Sr, Ss, \&c.$  of  $y$ ; then

$$\text{will } y = \frac{\overline{x-q} \cdot \overline{x-r} \cdot \overline{x-s} \cdot \&c.}{p-q \cdot p-r \cdot p-s \cdot \&c.} \times Sp + \frac{\overline{x-p} \cdot \overline{x-r} \cdot \overline{x-s} \cdot \&c.}{q-p \cdot q-r \cdot q-s \cdot \&c.} \times Sq +$$

$$\frac{\overline{x-p} \cdot \overline{x-q} \cdot \overline{x-s} \cdot \&c.}{r-p \cdot r-q \cdot r-s \cdot \&c.} \times Sr + \frac{\overline{x-p} \cdot \overline{x-q} \cdot \overline{x-s} \cdot \&c.}{s-p \cdot s-q \cdot s-r \cdot \&c.} \times Ss + \&c.$$

The truth of this problem very easily appears by writing  $p, q, r, s, \&c.$  for  $x$  in the given series.

All the preceding examples may be applied to this case, by writing  $x$  for  $m$  in the given series; hence the resolutions of several cases of equi-distant ordinates by easy and not inelegant serieses, amongst which are included the two cases commonly given on this subject.

14. If a quantity be required, which proceeds according to the dimensions of  $x$ , reduce the above given value of  $y$  into a quantity proceeding according to the dimensions of  $x$ , and

there results  $y = \left( \frac{Sp}{p-q \cdot p-r \cdot p-s \cdot \&c. = A} + \frac{Sq}{q-p \cdot q-r \cdot q-s \cdot \&c. = B} \right.$

$$\left. + \frac{Sr}{r-p \cdot r-q \cdot r-s \cdot \&c. = C} + \frac{Ss}{s-p \cdot s-q \cdot s-r \cdot \&c. = D} + \&c. \right) \times x^{n-1}$$

$$\left( \frac{Sp \times q+r+s+\&c.}{A} + \frac{Sq \times p+r+s+\&c.}{B} + \frac{Sr \times p+q+s+\&c.}{C} + \right.$$

$$\left. \frac{Ss \times p+q+r+\&c.}{D} + \&c. \right) x^{n-1} + \left( \frac{Sp \times qr+qs+rs+\&c.}{A} + \right.$$

$$\frac{Sq \times pr + ps + rs + \&c.}{B} + \frac{Sr \times pq + ps + qs + \&c.}{C} + \frac{Ss \times pq + pr + qr + \&c.}{D} + \&c.) \\ \times x^{n-2} - \left( \frac{Sp \times qrs + \&c.}{A} + \frac{Sq \times prs + \&c.}{B} + \frac{Sr \times pqs + \&c.}{C} + \frac{Ss \times pqr + \&c.}{D} + \right. \\ \left. \&c. \right) x^{n-3} + \&c.$$

The law and continuation of this series is evident to any one versant in these matters from inspection.

These fractions may be reduced to a common denominator by substituting for  $Sp$  and  $A$  the products  $Sp \times P$  and  $A \times P$ , where  $P = \overline{q-r} \cdot \overline{q-s} \cdot \overline{r-s} \cdot \&c.$ ; for  $Sq$  and  $B$  the products  $Sq \times Q$  and  $B \times Q$ , where  $Q = \overline{p-r} \cdot \overline{p-s} \cdot \overline{r-s} \cdot \&c.$ ; for  $Sr$  and  $C$  the products  $Sr \times R$  and  $C \times R$ , where  $R = \overline{p-q} \cdot \overline{p-s} \cdot \overline{q-s} \cdot \&c.$ ; for  $Ss$  and  $D$  the products  $Ss \times S'$  and  $C \times S'$ , where  $S' = \overline{p-q} \cdot \overline{p-r} \cdot \overline{q-r} \cdot \&c. \&c.$

The fractions, in particular cases, will often be reducible to lower terms.

15. Let  $y = ax^b + bx^{b+1} + cx^{b+2} + \&c.$ , and the correspondent values of  $x$  and  $y$  be given as before, then will  $y =$

$$\frac{x^b \times \overline{x^l - q^l} \times \overline{x^l - r^l} \times \overline{x^l - s^l} \times \&c.}{p^b \times \overline{p^l - q^l} \times \overline{p^l - r^l} \times \overline{p^l - s^l} \times \&c.} \times Sp + \frac{x^b \times \overline{x^l - p^l} \times \overline{x^l - r^l} \times \overline{x^l - s^l} \times \&c.}{q^b \times \overline{q^l - p^l} \times \overline{q^l - r^l} \times \overline{q^l - s^l} \times \&c.} \times Sq \\ + \frac{x^b \times \overline{x^l - p^l} \times \overline{x^l - q^l} \times \overline{x^l - s^l} \times \&c.}{r^b \times \overline{r^l - p^l} \times \overline{r^l - q^l} \times \overline{r^l - s^l} \times \&c.} \times Sr + \frac{x^b \times \overline{x^l - p^l} \times \overline{x^l - q^l} \times \overline{x^l - r^l} \times \&c.}{s^b \times \overline{s^l - p^l} \times \overline{s^l - q^l} \times \overline{s^l - r^l} \times \&c.} \times Ss + \&c.$$

This series may in the same manner as the preceding be reduced to terms, proceeding according to the dimensions of  $x$ ; and the serieses given in the examples may (*mutatis mutandis*) be predicated of it.

16. A more general method of correspondent values is given in the Meditations, as also the subsequent  $y = \frac{\overline{x-q} \cdot \overline{x-r} \cdot \overline{x-s} \cdot \&c.}{\overline{p-q} \cdot \overline{p-r} \cdot \overline{p-s} \cdot \&c.}$

$$\begin{aligned} & \times Sp + \frac{\overline{x-p} \cdot \overline{x-r} \cdot \overline{x-s} \cdot \&c.}{q-p \cdot q-r \cdot q-s \cdot \&c.} \times Sq + \frac{\overline{x-p} \cdot \overline{x-q} \cdot \overline{x-s} \cdot \&c.}{r-p \cdot r-q \cdot r-s \cdot \&c.} \times Sr + \\ & \&c. \text{ as in Ex. I.} = Sp + (x-p) \left( \frac{1}{p-q} \times Sp + \frac{1}{q-p} \times Sq \right) + (x-p) \\ & (x-q) \left( \frac{1}{p-q} \times \frac{1}{p-r} \times Sp + \frac{1}{q-p} \times \frac{1}{q-r} \times Sq + \frac{1}{r-p} \times \frac{1}{r-q} \times Sr \right) + \\ & (x-p)(x-q)(x-r) \left( \frac{1}{p-q} \cdot \frac{1}{p-r} \cdot \frac{1}{p-s} \cdot \times Sp + \frac{1}{q-p} \cdot \frac{1}{q-r} \cdot \frac{1}{q-s} \right. \\ & \left. \times Sq + \frac{1}{r-p} \cdot \frac{1}{r-q} \cdot \frac{1}{r-s} \times Sr + \frac{1}{s-p} \cdot \frac{1}{s-q} \cdot \frac{1}{s-r} \times Ss \right) - \&c. \end{aligned}$$

The equality of these two different quantities will easily appear by finding the co-efficients of both, which are multiplied into the same given value of  $y$  as  $Sp, Sq, Sr, \&c.$  and the same power of  $x$ ; for with very little difficulty they will in general be found equal.

It is evident from this resolution that, giving the ordinates and their respective distances from each other, the value of any other ordinate at a given distance from the preceding, found by this method, will result the same, whatever may be the point assumed from which the absciss is made to begin.

3.

I. Let a series be  $Ax + Bx^2 + Cx^3 + Dx^4 + \&c.$  of such a formula that if in it for  $x$  be substituted  $a + b$ , there results a series  $A \times \overline{a+b} + B \times \overline{a+b}^2 + C \times \overline{a+b}^3 + D \times \overline{a+b}^4 + \&c. = (Aa + Ba^2 + Ca^3 + Da^4 + \&c.) \times (1 + qb + rb^2 + sb^3 + tb^4 + \&c.) + (1 + qa + ra^2 + sa^3 + ta^4 + \&c.) \times (Ab + Bb^2 + Cb^3 + Db^4 + \&c.)$  then will the series  $Ax + Bx^2 + Cx^3 + Dx^4 + \&c. = Ax + \frac{2B}{1 \cdot 2} x^2 + \frac{2 \cdot 3C}{1 \cdot 2 \cdot 3} x^3 + \frac{24ABC - 8B^3}{1 \cdot 2 \cdot 3 \cdot 4A^2} x^4 + \frac{36C^2A^2 + 24ACB^2 - 16B^4}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5A^3} x^5 + \frac{9 \cdot 24A^2BC^2 - 4 \times 24AB^3C}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6A^4} x^6 + \frac{216C^3A^3 + 432A^2B^2C^2 - 384ACB^4 + 64B^6}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7A^5} x^7 + \&c. ;$

$$\begin{aligned}
 & + \&c.; \text{ and the series } 1 + qx + rx^2 + sx^3 + tx^4 + \&c. = 1 + \frac{B}{A}x + \\
 & \frac{6CA - 2B^2}{1 \cdot 2A^2}x^2 + \frac{18CAB - 8B^3}{1 \cdot 2 \cdot 3A^3}x^3 + \frac{36C^2A^2 - 8B^4}{1 \cdot 2 \cdot 3 \cdot 4A^4}x^4 + \\
 & \frac{180C^2A^2B - 120ACB^3 + 16B^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5A^5}x^5 + \frac{216C^3A^3 + 216A^2C^2B^2 - 288ACB^4 + 64B^6}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6A^6}x^6 \\
 & + \&c.
 \end{aligned}$$

The terms of these two series can easily be deduced by the subsequent method. Let  $Kx^{n-2} + Lx^{n-1} + Mx^n$ , be successive terms of the series  $Ax + Bx^2 + Cx^3 + \&c.$ , and  $K^1x^{n-2} + L^1x^{n-1}$  successive terms of the series  $1 + qx + rx^2 + sx^3 + tx^4 + \&c.$ ; then will  $M = \frac{2A^2 \times B \times K^1 + 6CAK - 2B^2K}{n \cdot n - 1 \times A^2}$  and  $L^1 = \frac{n \times A \times M - B \times xL}{A^2}$ .

*Cor. 1.* Let  $B = 0$ , and the two series  $Ax + Bx^2 + Cx^3 + Dx^4 + \&c.$  and  $1 + qx + rx^2 + \&c.$  become respectively  $Ax + \frac{2 \cdot 3}{2 \cdot 3}Cx^3 + \frac{2^2 \cdot 3^2}{2 \cdot 3 \cdot 4 \cdot 5} \times \frac{C^2}{A}x^5 + \frac{2^3 \cdot 3^3}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} \times \frac{C^3}{A^2}x^7 + \frac{2^4 \cdot 3^4}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9} \times \frac{C^4}{A^3}x^9 + \&c.$ , and  $1 + \frac{2 \cdot 3}{1 \cdot 2} \times \frac{C}{A}x^2 + \frac{2^2 \cdot 3^2}{1 \cdot 2 \cdot 3 \cdot 4} \times \frac{C^2}{A^2}x^4 + \frac{2^3 \cdot 3^3}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} \times \frac{C^3}{A^3}x^6 + \&c.$

If in these series for  $A$  be substituted  $1$ , and for  $C$  be substituted  $-\frac{1}{2 \cdot 3}$ , there will result the series  $x - \frac{x^3}{2 \cdot 3} + \frac{x^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c.$ , and  $1 - \frac{x^2}{1 \cdot 2} + \frac{x^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.$  which give the sine and cosine in terms of the arc  $x$ .

*Cor. 2.* Let  $C = 0$ , and the above-mentioned series  $Ax + Bx^2 + \&c.$  becomes  $Ax + \frac{2}{1 \cdot 2}Bx^2 * - \frac{2^3}{1 \cdot 2 \cdot 3 \cdot 4} \times \frac{B^3}{A^2}x^4 - \frac{2^4}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} \times \frac{B^4}{A^3}x^5 * + \frac{2^6}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} \times \frac{B^6}{A^5}x^7 + \frac{2^7}{1 \cdot 2 \cdot 3 \cdot 4 \dots 8} \times \frac{B^7}{A^6}x^8 * - \frac{2^9}{1 \cdot 2 \cdot 3 \cdot 4 \dots 10} \times \frac{B^9}{A^8}x^{10} - \frac{2^{10}}{1 \cdot 2 \cdot 3 \dots 11} \times \frac{B^{10}}{A^9}x^{11} + \&c.$  The law of

this



this series is, first, that every third term vanishes; and, secondly, the signs of every two successive terms change alternately from + to - and - to +; and, lastly, the co-efficient

of the term  $x^n$  is  $\frac{2^{n-1}}{1 \cdot 2 \cdot 3 \dots n} \times \frac{B^{n-1}}{A^{n-2}}$ ; and the series  $1 + qx + rx^2$

$$+ \&c. \text{ becomes } 1 + \frac{B}{A}x - \frac{2B^2}{1 \cdot 2A^2}x^2 - \frac{2^3B^3}{1 \cdot 2 \cdot 3A^3}x^3 - \frac{2^3B^4}{1 \cdot 2 \cdot 3 \cdot 4A^4}x^4$$

$$+ \frac{2^4B^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5A^5}x^5 + \frac{2^6B^6}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6A^6}x^6 + \frac{2^6 \times B^7}{1 \cdot 2 \cdot 3 \dots 7A^7}x^7 -$$

&c. In this series the signs of three successive terms alternately change from + to - and - to +; and the co-efficient

of the term  $x^n$  is  $\frac{2^n \times B^n}{1 \cdot 2 \cdot 3 \dots nA^n}$  or  $\frac{2^{n-1} \times B^n}{1 \cdot 2 \cdot 3 \dots nA^n}$  according as  $n$  is

divisible by 3 or not.

2. Let a series  $1 + Px + Qx^2 + Rx^3 + Sx^4 + Tx^5 + \&c.$  be of such a formula, that if in it for  $x$  be substituted  $a + b$ , there

results a series  $1 + P \times \overline{a + b} + Q \times \overline{a + b}^2 + R \times \overline{a + b}^3 + S \times \overline{a + b}^4 + \&c. = (1 + Pa + Qa^2 + Ra^3 + Sa^4 + \&c.) \times (1 + Pb + Qb^2 + Rb^3 + Sb^4 + \&c.) + (Aa + Ba^2 + Ca^3 + Da^4 + \&c.) \times (Ab + Bb^2 + Cb^3 + Db^4 + \&c.)$ , then will the series  $Ax + Bx^2 + Cx^3 + Dx^4 +$

$$\&c. = Ax + Bx^2 + \left( \frac{2B^2}{3A} - \frac{PB}{3} + A \times \frac{A^2 + P^2}{6} \right) x^3 + \frac{2B^3 - 2PAB^2 + A^2 \times$$

$$\frac{A^2 + P^2 \times B}{6A^2} x^4 + \&c., \text{ and the series } 1 + Px + Qx^2 + Rx^3 + \&c. = 1 +$$

$$Px + \frac{A^2 + P^2}{2} x^2 + \frac{2AB + P \times \overline{A^2 + P^2}}{6} x^3 + \frac{4B^2 + \overline{A^2 + P^2}^2}{24} x^4 + \&c. \text{ Let}$$

$Kx^{n-2} + Lx^{n-1} + Mx^n$  be successive terms of the series  $Ax + Bx^2 + Cx^3 + \&c.$ , and  $K'x^{n-2} + L'x^{n-1} + M'x^n$  successive terms of the series  $1 + Px + Qx^2 + Rx^3 + \&c.$ ; then will  $A \times L + P \times L' = n$

$\times M'$  and  $B \times K + Q \times K' = n \cdot \frac{n-1}{2} \times M'$  expresses the law of the serieses.

Cor. Let  $B=0$ , then the series  $Ax + Bx^2 + Cx^3 + Dx^4 = A \times (x + \frac{P^2 \times A^2}{2 \cdot 3} x^3 + \frac{(P^2 + A^2)^2}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} x^5 + \frac{(P^2 + A^2)^3}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7} x^7 + \&c.)$ , and the series  $1 + Px + Qx^2 + Rx^3 + \&c. = 1 + Px + \frac{P^2 + A^2}{1 \cdot 2} x^2 + P \times \frac{P^2 + A^2}{1 \cdot 2 \cdot 3} x^3 + \frac{(P^2 + A^2)^2}{1 \cdot 2 \cdot 3 \cdot 4} x^4 + P \times \frac{(P^2 + A^2)^2}{1 \cdot 2 \cdot 3 \dots 5} x^5 + \frac{(P^2 + A^2)^3}{1 \cdot 2 \cdot 3 \dots 6} x^6 + \&c.$ ; the co-efficient of the term  $x^n$  will be  $(P^2 + A^2)^{\frac{n}{2}}$  or  $P \times (P^2 + A^2)^{\frac{n-1}{2}}$ , according as  $n$  is even or odd.

If in the equations before given for  $x$  be substituted  $a = b$  instead of  $a + b$ , then in the other quantities for  $b$  substitute  $-b$ .

3. If in Case 2. the difference between the two quantities  $(1 + Pa + Qa^2 + \&c.) \times (1 + Pb + Qb^2 + \&c.)$  and  $(Aa + Ba^2 + Ca^3 + \&c.) \times (Ab + Bb^2 + Cb^3 + \&c.)$  is assumed  $= 1 + P \times \overline{a + b} + Q \times \overline{a + b^2} + \&c.$ , then in the serieses before given for  $A, B, C, \&c.$  write respectively  $\sqrt{-1}A, \sqrt{-1}B, \sqrt{-1}C, \&c.$ , and there will result the corresponding serieses.

The same principles may be applied to many other cases.

4. Equations of these formulæ may be useful, when the sums of the serieses correspondent to a value ( $a$ ) of  $x$  are given, and the sums of the series correspondent to a value ( $a + b$ ) of  $x$  is required,  $b$  having a small ratio to  $a$ : for instance, let the given series be  $x - \frac{x^3}{2 \cdot 3} + \frac{x^5}{2 \cdot 3 \cdot 4 \cdot 5} - \frac{x^7}{2 \cdot 3 \dots 7} + \&c.$ ; the equation found in the first case is  $a + b - \frac{(a+b)^3}{2 \cdot 3} + \frac{(a+b)^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c. = (a - \frac{a^3}{2 \cdot 3} + \frac{a^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c.) \times (1 - \frac{b^2}{1 \cdot 2} + \frac{b^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.) + (1 - \frac{a^2}{1 \cdot 2} + \frac{a^4}{1 \cdot 2 \cdot 3 \cdot 4} - \&c.) \times (b - \frac{b^3}{2 \cdot 3} + \frac{b^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c.)$ ; but

but  $a - \frac{a^3}{2 \cdot 3} + \frac{a^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c.$ , and  $1 - \frac{a^2}{1 \cdot 2} + \frac{a^4}{2 \cdot 3 \cdot 4} - \&c.$  are the sine ( $s$ ) and cosine ( $c$ ) of an arc  $a$  of a circle whose radius is 1; and, consequently, if the sine  $s$  and cosine  $c$  of an arc  $a$  be given, the sine of an arc  $(a+b) = s \times (1 - \frac{b^2}{2} + \frac{b^4}{24} - \&c.) + c(b - \frac{b^3}{2 \cdot 3} + \frac{b^5}{2 \cdot 3 \cdot 4 \cdot 5} - \&c.)$ , which series, if  $b$  be very small in proportion to  $a$ , converges much faster than the common series for finding the sine from the arc: it has been given from different principles in the *Meditationes*, and is also easily deducible from the series for finding the sine and cosine from the arc by the propositions usually given in plane trigonometry: the cosine of the same arc  $(a+b) = c \times (1 - \frac{b^2}{1 \cdot 2} + \frac{b^4}{2 \cdot 3 \cdot 4} - \&c.) - s \times (b - \frac{b^3}{1 \cdot 2 \cdot 3} + \frac{b^5}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} - \&c.)$ ,

Ex. 2. Let the series be  $\overline{a+b} + \frac{\overline{a+b^3}}{2 \cdot 3} + \frac{\overline{a+b^5}}{2 \cdot 3 \cdot 4 \cdot 5} + \&c. =$   
 $(a + \frac{a^3}{2 \cdot 3} + \frac{a^5}{2 \cdot 3 \cdot 4 \cdot 5} + \&c.) \times (1 + \frac{b^2}{1 \cdot 2} + \frac{b^4}{1 \cdot 2 \cdot 3 \cdot 4} + \&c.) +$   
 $(1 + \frac{a^2}{1 \cdot 2} + \frac{a^4}{1 \cdot 2 \cdot 3 \cdot 4} + \&c.) \times (b + \frac{b^3}{1 \cdot 2 \cdot 3} + \frac{b^5}{2 \cdot 3 \cdot 4 \cdot 5} + \&c.);$   
 but  $a + \frac{a^3}{1 \cdot 2 \cdot 3} + \&c.) = x$ , and  $1 + \frac{a^2}{1 \cdot 2} + \frac{a^4}{1 \cdot 2 \cdot 3 \cdot 4} + \&c. =$   
 $\sqrt{1+x^2}$ , if  $a$  be the hyperbolic log. of  $x + \sqrt{1+x^2}$ ; therefore  
 $a + b + \frac{\overline{a+b^3}}{2 \cdot 3} + \frac{\overline{a+b^5}}{2 \cdot 3 \cdot 4 \cdot 5} + \&c. = x \times (1 + \frac{b^2}{2} + \frac{b^4}{2 \cdot 3 \cdot 4} + \&c.) +$   
 $\sqrt{1+x^2} \times (b + \frac{b^3}{2 \cdot 3} + \&c.)$

Let  $b + \frac{b^3}{2 \cdot 3} + \frac{b^4}{2 \cdot 3 \cdot 4 \cdot 5} + \&c. = y$ , and  $(x + \sqrt{1+x^2} \times$   
 $(y + \sqrt{1+y^2})) = V$ , then will  $\overline{a+b} + \frac{\overline{a+b^3}}{2 \cdot 3} + \frac{\overline{a+b^5}}{2 \cdot 3 \cdot 4 \cdot 5} + \&c. =$   
 $\frac{1}{2} V - \frac{1}{V}$ .

5. Let a quantity  $P$  be a function of  $x$ , or the fluent of a function of  $x \times \dot{x}$ , and the value  $X$  of it when  $x = a$  be known, and the value of it when  $x = a + b$  be required. Find a series of which the first term is  $X$ , and which proceeds according to the dimensions of  $b$ , if  $b$  be a very small quantity, and in general at least so small that the series from  $x = a$  to  $x = a + b$  neither becomes infinite or 0.

In the same manner, if an algebraical or fluxional equation or equations, expressing the relations between  $x, y, z, v, \&c.$  be given, find the correspondent values of  $y, z, v, \&c.$  to  $x = a$ , which let be  $Y, Z, V, \&c.$ ; then find serieses for  $y, z, v, \&c.$  of which the first terms let be  $Y, Z, V, \&c.$  respectively, and which proceed according to the dimensions of  $b$ , but subject to the same conditions as in the preceding case.

From fluxional equations may be deduced series which express the value of  $y, \&c.$  in terms of  $x$ , and always diverge, or always converge, whatever may be its value, as appears from the *Meditationes*.



XVIII. *On the Resolution of attractive Powers.* By Edward Waring, M. D. F. R. S. and Lucasian Professor of Mathematics at Cambridge.

Read May 28, 1789.

1. **A** FORCE acting at a given point may be resolved by an infinite number of ways into two, three, or more ( $n$ ) forces acting at the same point, either in the same or different planes with the given force and each other; and, *vice versa*, any number of such forces acting in the same or different planes may be reduced into one.

Ex. Fig. 1. Tab. III. Let a body A be acted on by three forces AB, AC, and AD, not being in the same plane; reduce any two of them AB and AC to one AE, by completing the parallelogram ABEC; then reduce the two forces AE and AD to one AF by completing the parallelogram AEDF, and the three forces AB, AC, and AD, are reduced to the one AF.

2. If  $n$  forces act on the body A at the same time, and any ( $n - 1$ ) of them be reduced to one, the force resulting will be situated in the same plane with the remaining, and force equivalent to the ( $n$ ) forces.

3. If one force  $a$  be resolved into several others  $x, y, z, v,$  &c. situated in different planes, and the sines of the angles, which the forces  $y, z, v,$  &c. contain with the plane made by the direction of the forces  $x$  and  $a$  be respectively  $s, s', s'',$  &c. then will  $sy \pm s'z \pm s''v \pm \&c. = 0.$

## P R O B L E M I.

Fig. 2. Given the law of attraction of each of the parts of a given line in terms of their distance from a given point P; to find the attraction of the whole line  $ab$  on the point P.

Find the attraction of the line  $ab$  on the point P in the two directions  $Pf$  and  $fb$  by the following method. Draw  $Px$  from the point P to any point  $x$  of the line  $ab$ , the force acting on the point P by the particle  $xy$  will be the given function (determined from the given law of attraction) of the distance into the particle; draw also  $Pb$  perpendicular from the point P to the line  $ab$ , and let  $Pf = a$ ,  $bf = b$ , and  $fx = y$ ; then will the distance  $Px = \sqrt{(a^2 \pm 2by + y^2)}$ , and the function of the distance into the particle  $xy = \phi(\sqrt{(a^2 \pm 2by + y^2)}) \times y = F(y) \times y$ ; let this be denoted by  $lx$  situated in the line  $Px$ , which resolve into two others  $nx = \frac{yy \times F : (y)}{Px = \sqrt{(a^2 \pm 2by + y^2)}}$  situated in the line  $ab$ , and  $ln$  (in a direction parallel to  $Pf$ )  $= \frac{ay \times F : (y)}{\sqrt{(a^2 \pm 2by + y^2)}}$ ; find the fluents of the fluxions  $\frac{yy \times F : (y)}{Px}$  and  $\frac{ay \times F : (y)}{Px}$  contained between the values  $af$  and  $fb$  of the line  $fx = y$ , which suppose Y and V respectively; through the point  $p$  draw  $Py$  parallel to  $fb = Y$ , and in the line  $Pf$  assume  $Pu = V$ ; complete the parallelogram  $Puzy$ ;  $Pz$  will be the force of the line  $ab$  on the point P.

Cor. If  $F : (y)$  varies as any power or root ( $2n$ ) of the distance  $Px = \sqrt{(a^2 \pm 2by + y^2)}$ , and  $n - \frac{1}{2}$  be an integer affirmative number or 0, the fluents Y and V of both the fluxions can be found in finite algebraical terms of  $y$ ; if  $n - \frac{1}{2}$  be an integer negative number, both the fluents can be found in the above-mentioned

mentioned finite terms together with the arc of a circle, whose radius is  $\sqrt{a^2 - b^2}$  and tangent  $y \mp b$ , unless  $n - \frac{1}{2} = -1$ , in which case the fluent  $Y$  involves that circular arc, and also the logarithm of  $y^2 \pm 2by + a^2$ . If  $n - \frac{1}{2}$  denotes a fraction whose denominator is 2, both the fluents can be expressed by the finite terms together with the log. of  $y \pm b + \sqrt{(y^2 \pm 2by + a^2)}$ . If the fluents be given, when  $n$  is a given quantity, and  $n - \frac{1}{2}$  not a whole affirmative number, from them can be deduced the fluents of any fluxions resulting by increasing or diminishing  $n$  by a whole number, unless in the above-mentioned case of  $n - \frac{1}{2} = -1$ . If  $b = 0$ , and consequently the line  $Pf$  is perpendicular to the given line  $ab$ , the fluent  $Y$  will be expressed by the finite terms, unless  $n - \frac{1}{2} = -1$ , in which case it will be as  $\frac{1}{2} \log. (y^2 + a^2)$  when properly corrected.

These fluxions  $\dot{Y}$  and  $\dot{V}$  may be transformed into others, whose variable quantity is  $Px = u$  the distance from  $P$ , by substituting in the fluxions for  $y$  and  $\dot{y}$  their respective values  $\sqrt{(u^2 - a^2 + b^2)} \mp b$  and  $\frac{u\dot{u}}{\sqrt{(u^2 - a^2 + b^2)}}$ , and consequently for  $\sqrt{(y^2 \pm 2by + a^2)}$  its value  $u$ .

#### P R O B L E M   I I .

Fig. 3. Given the attraction of each of the parts of a given surface in terms of their distance from a given point  $P$ , and an equation expressing the relation between an absciss  $Ap = x$ , and its correspondent ordinates  $pm = y$  of the surface; to find the attraction of the surface on the given point  $P$ .

First, by the preceding proposition find the attractions  $Y$  and  $V$  of any ordinate  $m p m'$  in the directions of the ordinate  $pm$  and of the line  $Pp$ ; and from the equation expressing the relation

relation

relation between the absciss and ordinates of the given curve, find the absciss in terms of the ordinates  $(pm) = \pi : (y)$ , and thence  $\dot{x} = \phi : (y) \times \dot{y}$  and  $\sqrt{(a'^2 \pm 2sa'x + x^2)} = \phi' : (y)$ , where  $PA = a'$  and  $s = \text{cofine}$  of the angle, which the absciss  $Ap$  makes with the line  $PA$ ; then find the fluents of the three fluxions  $\dot{x} \times Y = \dot{y} \times Y \times \phi : (y)$ ,  $\dot{x} \times \frac{V \times x}{\sqrt{(a'^2 \pm 2sa'x + x^2)}} = \phi : (y) \times \dot{y} \times \frac{\pi : (y)}{\phi : (y)} \times V$  and  $\dot{x} \times \frac{a'V}{\sqrt{(a'^2 \pm 2sa'x + x^2)}} = \dot{y} \times \frac{a'V}{\phi' : (y)}$  contained between the values of  $y$ , which correspond to the extreme values of  $x$ , which suppose  $Y'$ ,  $V'$ , and  $Z$ ; and draw through the point  $P$  the lines  $Py$  and  $Pz$  respectively parallel to the ordinates  $pm$  and to the absciss  $Ap$  and equal to  $r \times Y'$  and  $V'$ ; assume  $Pu$  in the line  $(PA) = t \times Z$ ,  $r$  and  $t$  denoting the fines of the angles, which the ordinates  $pm$  and line  $AP$  make with the absciss  $Ap$ : reduce these three forces  $Py$ ,  $Pz$ , and  $Pu$ , to one  $Pf$ , and  $Pf$  will be the force of the surface on the point  $P$ .

*Cor. 1.* If for  $y$  and  $\dot{y}$  be substituted their values in terms of  $x$  and  $\dot{x}$ , deduced from the equation expressing the relation between the absciss  $Ap$  and ordinate  $pm$  of the given curve, thence will be deduced the above-mentioned fluents  $Y$ ,  $V$ ,  $Y'$ ,  $V'$ , and  $Z$ , in terms of  $x$ ; and in the same manner, if for  $x$  and  $\dot{x}$  be substituted in the fluxions or fluents resulting their values  $\sqrt{(u^2 - a'^2 + 1^2 a'^2)} = sa'$ , and its fluxion, there will result the above-mentioned fluxions or fluents in terms of  $u$  the distance from the point  $P$ .

*Cor. 2.* Let the curve be a circle, of which  $A$  is the center,  $PA$  a line perpendicular to the plane of the circle, and the ordinate  $pm$  perpendicular to the absciss  $Ap$ ; the forces on each side of the absciss  $Ap$  will be equal, and the force in the direc-



tion of the abscifs  $Ap$  will be equal to that in the contrary direction; the force in the direction  $(PA) = 4 \times \int \frac{ai}{\sqrt{(u^2 - a^2)}} \times \int \frac{uy}{\sqrt{u^2 + y^2}} \times F : (\sqrt{(u^2 + y^2)}) = W$ , in which  $F : (\sqrt{u^2 + y^2})$  is the function of the distance, according to which the given force on the particles varies; the fluent  $\int \frac{uy}{\sqrt{(u^2 + y^2)}} \times F : \sqrt{(u^2 + y^2)}$  is contained between the values  $0$  and  $\sqrt{(r^2 + a^2 - u^2)}$  of the quantity  $y$ , and the fluent  $W$  is contained between the values  $a$  and  $\sqrt{(a^2 + r^2)}$  of the quantity  $u$ , where  $a = PA$  and  $r$  the radius of the circle; but the same force is  $= 2 \times 3,14159$  &c.  $\times \int ai \times F : (u)$ , where  $F : (u)$  denotes the given function of the distance  $(u)$ , and the fluent is contained between the values  $a$  and  $\sqrt{a^2 + r^2}$  of  $u$ .

P R O B L E M III.

To find the attraction of a given solid on a given point  $P$ . Find the attraction of every parallel section on that point by the preceding problem, and multiply it into the correspondent fluxion of the first abscissa  $AP$ , and also find the fluent of the resulting fluxion, which, properly corrected, multiply into the sine of the angle, which the first abscissa makes with the parallel sections, and the product will be proportional to the attraction of the solid on the given point  $P$ .

2. Fig. 4. Let the solid  $ABCH$  be generated by the rotation of a given curve round its axis  $AB$ , which passes through the point attracted  $P$ , and this solid be supposed to consist of small evanescent solids, whose bases are the surfaces  $EF, ef, \&c.$  of spheres, of which the center is  $P$ , and altitudes  $Ff, \&c.$  the

increments of the base AB contained between the two contiguous surfaces EF and *ef*: from the points E and *e* of the curve draw ED and *ed* perpendicular to the axis AB, and ES perpendicular to the arc E*e* of the given curve at the point E, and meeting the axis AB in S; then will the evanescent solid EF*f**e* =  $\rho \times PE \times FD \times Ff = \rho \times FD \times PS \times Dd$  (because  $Ff = \frac{PS \times Dd}{PE}$ ) =  $\rho \times \sqrt{(z^2 + y^2)} - z \times \frac{z\dot{z} - yy\dot{y}}{\sqrt{(z^2 + y^2)}}$ , where *z* and *y* denote respectively the absciss PD, and its correspondent ordinate DE of the given curve.

The increment of the attraction of the surface EF on the point P in the direction PD will be as the increment of the surface ( $\rho \times PE \times Dd$ )  $\times \frac{PD}{PE}$   $\times$  force of each particle =  $\rho \times PD \times Dd \times$  given force of the particle; but the fluent of the fluxion  $PD \times Dd$  contained between the points E and F is =  $\frac{1}{2}PE^2 - \frac{1}{2}PD^2 = \frac{1}{2}ED^2$ , whence the attraction of the evanescent solid EF*f**e* is as  $\frac{1}{2}\rho \times ED^2 \times Ff \times F : (\sqrt{x^2 + y^2})$  force of each given particle at the distance ( $PE = \sqrt{(x^2 + y^2)}) = \frac{1}{2}\rho \times ED^2 \times \frac{PS}{PE} \times Dd \times F : (\sqrt{z^2 + y^2}) = \frac{1}{2}\rho y^2 \times \frac{z\dot{z} - yy\dot{y}}{\sqrt{(z^2 + y^2)}} \times F : (\sqrt{(z^2 + y^2)})$ ; the fluent of which, properly corrected, is as the attraction of the solid on the point P;  $\rho$  denotes the circumference of a circle, whose radius is 1.

*Cor.* 1. The fluxion of this solid is  $\frac{1}{2}\rho y^2 \dot{z} = \dot{V}$  which deduced from the preceding principles =  $\rho \times (\sqrt{(z^2 + y^2)} - z) \times (z\dot{z} - yy\dot{y}) = \dot{V}$ , and consequently their fluents between two values of *z*, which correspond to two values of  $y = 0$ , will be equal to each other.

*Cor.*

Cor. 2. The increment of the attraction of this solid as given in this proposition  $\frac{1}{2} p \times y^2 \times \frac{z\dot{z} \mp y\dot{y}}{\sqrt{(z^2 + y^2)}} \times F : (\sqrt{(z^2 + y^2)}) = \dot{U}$ , but in the preceding proposition the force of a circle on the point P =  $p \times \int au \times F : (u)$ , where  $u = \sqrt{(z^2 + y^2)}$  and  $a = z$ , and  $y$  or  $u$  the only variable quantity contained in the fluxion; and consequently the fluxion of the attraction of the solid  $p \times \dot{z} \int z \frac{y\dot{y}}{\sqrt{z^2 + y^2}} \times F : ((z^2 + y^2)^{\frac{1}{2}}) = \dot{W}$ ; therefore, if for the fluent of  $\frac{zy\dot{y}}{\sqrt{(z^2 + y^2)}} \times F : ((z^2 + y^2)^{\frac{1}{2}})$  be substituted its fluent contained between the values  $a$  and the value of  $y$ , which in the given equation corresponds to  $z$ ; then the fluents of  $\dot{U}$  and  $\dot{W}$  contained between the two values of  $z$ , which corresponds to two values of  $y = 0$ , will be equal to each other.

The difference of the fluents of  $\dot{Y}$  and  $\dot{V}$ , &c. contained between any other two values of  $z$ , can easily be deduced from the difference of two segments of spheres.

1. It may not be improper to remark in this place, that from different methods of finding the sum of quantities, the fluents of fluxions, the integrals of increments, &c. quantities may often be deduced equal, which otherwise cannot without some difficulty; of which instances are contained in the Meditations, and I shall here subjoin one or two more to those already given in this Paper.

Ex. 1. Any curvilinear area ABC, &c. may be supposed to consist of evanescent areas EFef, of which the base EF is the arc of a circle, whose radius is PE =  $\sqrt{(z^2 + y^2)}$  and sine ED =  $y$ , and altitude Ff, and consequently the fluxion of the area =  $Ff \times \text{arc}(A)$  of a circle whose radius is PE and sine ED =

$\frac{PS}{PE} \times \dot{z} \times A = \frac{z\dot{z} - y\dot{y}}{\sqrt{(z^2 + y^2)}} \times A = \dot{V}$ ; the fluent of  $\dot{V}$  contained between the two values of  $z$  which correspond to two values of  $(y) = 0$  will be equal to the fluent of  $y\dot{z}$  contained between the same two values of  $z$ .

Ex. 2. The attraction of any circular arc  $EF$  in the direction  $PD$  on a point  $P$  ( $P$  being the center of a circle, of which  $EF$  is an arc, and  $ED$  the sine of that arc) will be as  $ED \times$  force at distance  $PE = ED \times F : (PE)$ ; for the attraction in the direction  $PD // F$  at the point  $x$  is as the increment of the arc  $xy \times$

$$F : (PE) \times \frac{Pl}{Px} \quad (xl' \text{ and } yl' \text{ being at right angles to } PF) = U \times \frac{Px}{xl'} \times \frac{Pl}{Px} \times F : (PE) = \frac{U' \times Pl}{ix} \times F : (PE) = \frac{ui}{\sqrt{PE^2 - u^2}} \times F : (PE),$$

if  $u = Pl$ ; and consequently the fluent of it is as  $\sqrt{PE^2 - u^2} \times F : (PE) = ED \times F : (PE)$ , and the attraction of the surface

$EFef$  will be as  $ED \times Ff \times F : (PE) = ED \times \dot{z} \times \frac{PS}{PE} \times F : (PE)$

$$= y \times \frac{z\dot{z} - y\dot{y}}{\sqrt{(z^2 + y^2)}} \times F : ((z^2 + y^2)^{\frac{1}{2}} = \dot{V}; \text{ the attraction of the curve}$$

will also vary as  $\int \dot{z} \int \frac{z\dot{z} \times F : ((u^2 + z^2)^{\frac{1}{2}})}{(z^2 + u^2)^{\frac{1}{2}}} = W$ , in which the

fluent of  $\frac{z\dot{z} \times F : ((z^2 + u^2)^{\frac{1}{2}})}{(z^2 + u^2)^{\frac{1}{2}}}$  is contained between  $u = 0$  and  $u = y$ ;

the fluents of  $\dot{V}$  and  $\dot{W}$  contained between two values of  $z$ , which correspond to two values of  $y = 0$ , will be equal to each other.

2. From a similar method may be deduced equalities between other like fluents, for the curve may be supposed to consist of other similar curve surfaces equally as circles, and the solid of similar segments of other solids equally as spheres.

3. From the same principles may innumerable serieses equal to each other be deduced; for by different converging serieses find

find the sum of the same quantity or quantities, and there will result serieses equal to each other: for instance (fig. 5), if the time of falling down the arcs AC and BC and their interpolations from the principles delivered in the *Meditationes Analyticæ*, of which the difference let be D; find the difference between the times of a body's falling through BC when it began to fall from A and from B by a series proceeding according to the dimensions of  $AB = o'$  a small quantity; and find, by a series of the same kind, the time of falling through AB; the sum of these two serieses will be equal to D. Similar propositions may be deduced from fluxional equations.

4. In some cases the ratios of the times of bodies falling through some particular distances to each other may be easily known; for instance, let the force vary as the  $m - 1$  power of the distance ( $x$ ), and  $a$  be the distance from which the body began to fall, then the velocity varies as  $\sqrt{(a^m - x^m)}$ , and the increment of the time as  $\frac{\dot{x}}{\sqrt{(a^m - x^m)}}$ ; but if the parts of different curves are proportional, then will  $a$ ,  $x$ , and  $\dot{x}$  vary in the same ratio as each other, and consequently the time through proportional parts of the distance will vary as  $a^{\frac{2-m}{2}}$ ; and if the bodies be resisted likewise by a force which varies as the  $\frac{2m-2}{m}$  power of the velocities, then the times through proportional parts will vary as before, that is, as  $a^{\frac{2-m}{2}}$ , where  $a$  denotes the proportional distances from the points where the forces and resistances are equal.

## P R O B L E M IV.

1. Fig. 6. Given an equation expressing the relation between the two abscissæ  $z = AP$  and  $x = Pp$  and their correspondent ordinates  $y = pm$  of a solid, to find its solid contents contained between two values of its first abscissæ  $z$ . Assume  $z$  as an invariable quantity, and from the equation resulting find the fluent  $Z$  of  $y\dot{x}$  contained between the extreme values of  $x$  or  $y$ ; then find the fluent of  $Z\dot{z}$  contained between the given values of  $z$ , and the fluent multiplied into the product of the sines of the angles, which the first abscissa makes with the plane of the ordinates and second absciss, and the second absciss makes with its correspondent ordinates, will be the solid content required.

2. Fig. 7. Let the first absciss  $z$  of a solid be perpendicular to the planes of the ordinates, and the second absciss  $Pp = x$  perpendicular to the ordinates themselves  $pm = y$ . First, assume the first absciss as invariable, and find the increment of the arc  $p'm = (\dot{x}^2 + \dot{y}^2)^{\frac{1}{2}}$ , then assume the second absciss  $Pp$  as constant, and let  $mu$  be the fluxion of the ordinate  $y$  or  $u$ , when the fluxion of the first absciss is  $\dot{z} = ul$ , where  $ul$  is perpendicular to the plane of the ordinates  $p'pm$ , and  $l$  a point of the surface of the solid; draw  $ub$  perpendicular to the arc  $p'm$ , and since  $ul$  is constituted at right angles to the plane  $pp'm$ ,  $lb$  will cut the arc  $p'm$  at right angles; but  $ub = \frac{um \times pp'}{p'm} = \frac{u\dot{x}}{\sqrt{(\dot{x}^2 + \dot{y}^2)}}$ ,

$lb = (bu^2 + lu^2)^{\frac{1}{2}} = \left(\frac{u\dot{x}}{\dot{x}^2 + \dot{y}^2} + \dot{z}^2\right)^{\frac{1}{2}}$  the fluxion of the surface will be  $lb \times \sqrt{(\dot{x}^2 + \dot{y}^2)}$ . From the given equation expressing the relation between the two abscissæ  $z$  and  $x$  and ordinates  $y$  find, by assuming  $z$  invariable  $p\dot{x} = \dot{y}$ , and by assuming  $x$  invariable  $q\dot{z} =$

$y' = u$ , which being substituted for their values in the quantity  $lh \times \sqrt{(x'^2 + y'^2)}$ , there will result  $(q^2 + p^2 + 1)^{\frac{1}{2}} \times x' \times z' = Ax'z' = \frac{(q^2 + p^2 + 1)^{\frac{1}{2}}}{p} \times y' \times z' = B'y'z'$ ; in A and B for  $y$  and  $x$  respectively substitute their value deduced from the given equation, and let the resulting quantities be  $A'x'z'$  and  $B'y'z'$ , where A' is a function of  $x$  and  $z$ , and B' a function of  $y$  and  $z$ ; find the fluent of  $A'x'z'$  from the supposition that  $x$  is only variable contained between the extreme values of  $x$  to a given value of  $z$ , which let be  $Lz$ , then find the fluent of  $Lz$  by supposing  $z$  only variable contained between given values of  $z$ , and it will be the surface of the solid contained between those values.

The same may be deduced by finding the fluent of  $B'y'z'$  on the supposition that  $y$  is the only variable quantity contained between the extreme values of  $y$  as before of  $x$  to a given value of  $z$ , which let be  $L'z$ ; then will the fluent of  $L'z$  contained between the given values of  $z$  be the surface required.

If the solid be a cone generated by the rotation of a rectangular triangle round a side containing the right angle as an axis;  $hu$  will be a given quantity, if  $z$  be given.

If the above-mentioned angles are given, but not right ones, the arc  $p'm$  and perpendicular  $lh$  can easily be deduced, and consequently the increment of the surface.

3. To define a curve of double curvature, it is necessary to have two equations expressing the relation between the abscissæ  $z$  and  $x$  and their ordinates ( $y$ ) given, and if the angles which they respectively make with each other be right ones; the fluxion of the arc as given in the *Proprietates Curvarum* is  $(z'^2 + x'^2 + y'^2)^{\frac{1}{2}}$ . Find its value from the two given equations in terms of  $x$ ,  $y$ , or  $z$ , multiplied into its respective fluxions, and its fluent, properly corrected, will be the length of the arc required.

If

If the angles are not right, they may easily be reduced to them.

4. The attractions of these surfaces, curves, &c. on a given point  $P$  may be deduced from the preceding principles of finding the attractions of each of the parts in the directions of the first abscissa, which passes through the point  $P$ , the second abscissa, and the ordinates, and then finding the integrals of these increments.

From the method which determines the attraction of a body, surface, &c. on a given point can be determined the attraction of a body, &c. on any number of points, and consequently the attraction of one body, &c. on another, &c.

It is sometimes advantageous to transform the first abscissæ, that it may pass through the point attracted: and the abscissæ and ordinates, that they may be at right angles to each other, &c.

#### P R O B L E M V.

1. Fig. 8. Given an equation expressing the relation between the two abscissæ  $AP$  and  $Pp$  of a solid, and their correspondent ordinates  $pm$ , or  $AP'$ ,  $P'p'$ , and  $p'm'$ ; to transform the first abscissa into any other  $Lb$ .

Let the abscissa  $Lb$  begin from a point  $L$  of the first abscissa  $AP$ , and meet an ordinate  $pm$  in the point  $b$ ; draw  $bp$ , and let the sines of the angles  $Ppm$ ,  $Pbp$ , and  $pPb$ ;  $LPb$ ,  $PbL$ , and  $PLb$ , be denoted respectively by  $r$ ,  $s$ , and  $t$ , &c.  $r'$ ,  $s'$ , and  $t'$ ; through a point  $b$  of the line  $Pb$  draw  $p'b'm'$  parallel to  $pm$ , and  $Lb = z$ ,  $bb' = x$ , and  $b'm' = y$ : in the given equation for  $AP$ ,  $Pp$ , and  $pm$  substitute respectively their correspondent



respondent values  $\frac{s'z}{r'} \pm AL(a)$ ,  $\frac{st'z}{rr'} \pm \frac{sx}{r}$  (for  $Pb = \frac{t'z}{r'}$  and  $Pb' = Pb \pm bb' = \frac{t'z}{r'} \pm x$ ), and  $y \pm \frac{tt'z}{rr'} \pm \frac{tx}{r'}$ ; there results an equation to the same solid expressing the relation between the two abscissæ  $z = Lb$  and  $x$ , and their correspondent ordinates  $y$ .

1. 2. If the absciss  $Lb$  does not begin from  $L$ , a point in the first given absciss  $AP$ , but from  $M$  a point given out of it, it may be reduced to the preceding case, by drawing from  $M$  a line  $MN = c$  to the plane of the first and second abscissæ parallel to the ordinates  $pm$ ; and from  $N$  to the first abscissa a line  $NO = b$  parallel to the second abscissæ, and substituting in the equation expressing the relation between  $AP$ ,  $Pp$ , and  $pm$  for  $AP$ ,  $Pp$ , and  $pm$  respectively  $z \pm AO (a)$ ,  $x \pm b$  and  $y \pm c$ ; and there results the equation required expressing the relation between the two abscissæ  $z$  and  $x$ , and their correspondent ordinates  $y$ , of which the first abscissa  $z$  passes through the point  $M$ .

2. To change the second abscissa  $Pp$  into any other  $Lb$ , the first abscissa and ordinates remaining the same. In the preceding figure let  $L$  be considered as a moveable point of the first absciss  $AL$ , and the sines of the respective angles denoted by the same letters as before, and  $Lb = x$ ,  $AL = z$ , and  $bm = y$ ; in the given equation for  $AP$ ,  $Pp$ , and  $pm$ , substitute  $z \pm \frac{s'x}{r'}$ ,  $\frac{st'x}{rr'}$ , and  $y \pm \frac{tt'x}{rr'}$ ; and there will result the equation required expressing the relation between  $z$  and  $x$  the abscissæ, and their correspondent ordinates  $y$ .

3. Fig 8. To change the ordinates, the abscissæ remaining the same, draw  $p'm$  an ordinate transformed,  $p'b$  parallel to the first abscissa  $AP$ , and meeting a second abscissa, of which

$pm$  is an ordinate in  $b$ : for the sines of the angles  $p'bp$ ,  $bpp'$ , and  $bp'p$ ;  $p'pm$ ,  $pm p'$ , and  $pp'm$  write  $r$ ,  $s$ , and  $t$ ,  $r'$ ,  $s'$ , and  $t'$ ; and for  $AP'$ ,  $P'p'$ , and  $p'm$  respectively  $z$ ,  $x$ , and  $y$ ; then substitute in the given equation for  $AP$ ,  $Pp$ , and  $pm$ , their respective values  $z (AP') = \frac{ss'}{rr'} \times y$ ,  $x (P'p') = \frac{ts'}{rr'} y$ , and  $\frac{t'}{r'} y$ ; and there results an equation to the solid expressing the relation between the two abscissæ  $AP'$  and  $P'p'$  and the transformed ordinates  $p'm$ .

From these cases, which are easily reducible to one, may be transformed any given abscissæ and their correspondent ordinates into any other containing given angles, &c. with the before-mentioned abscissæ and ordinates.

In the properties of curve lines, first published in 1762, is given a method of deducing the equation to any section of the solid, and in particular the case of deducing the equation to the projection of any curve on a given plane.

From the principles given in this, and the Paper on centripetal forces, which the Royal Society did me the honour to print, can be deduced the fluxional equations, whose fluents express the relations between the abscissæ and their correspondent ordinates of the curves described by bodies, of which the particles act on each other with forces varying according to given functions of their distances.



Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Fig. 7.




Fig. 8.





Fig. 8.





XIX. *Experiments on the Congelation of Quicksilver in England.*  
 By Mr. Richard Walker; in a Letter to Henry Cavendish,  
*Esq. F. R. S.*

Read May 28, 1789.

S I R,

I NOW beg leave to trouble you with the particulars of my experiments relative to the congelation of mercury; to which I shall add an account of a few experiments, relating to the production of artificial cold, made since my last Paper was written.

Exp. 1. On December 28th last, a favourable opportunity offered of beginning some experiments on the congelation of mercury, which I was desirous of effecting completely; how far I have succeeded will appear in the sequel.

For this purpose I prepared a mixture of diluted vitriolic acid (reduced by water till its specific gravity was to that of water as 1,5596 to 1) and strong fuming nitrous acid, of each equal parts. I preferred this mixture of acid because it has been found by Mr. M<sup>c</sup> NAB, in Hudson's Bay, to be capable of producing much greater cold, when the temperature of the materials at mixing is very low, than the nitrous acid alone; the former sinking a spirit thermometer to  $-54^{\circ}\frac{1}{2}$ , the latter never lower than  $-46^{\circ}$ .

The glass tube of a mercurial thermometer, with its bulb half filled with mercury, was provided, this occurring to me

as a convenient method of ascertaining when the mercury was congealed; for if, after being subjected to the cold of a frigorific mixture, the thermometer glass should be taken out and inverted, and the mercury found to remain completely suspended in that half of the bulb now uppermost, no doubt can remain of the success of the experiment; an hydrometer, with its lower bulb half an inch in diameter, and three-fourths full of mercury, was likewise provided, in case any accident should happen to the other.

It may be proper to premise here, that in all experiments of this kind I remove each vessel, when the liquor it contains is sufficiently cooled, out of the mixture in which it is immersed for that purpose, immediately previous to adding the snow or salts with intention to generate a still further increase of cold; and likewise prefer adding the snow or powdered salts to the liquor, instead of pouring the liquor upon these: it is necessary also to stir about the snow or salts, whilst cooling in a frigorific mixture, from time to time, otherwise it will freeze into a hard mass, and frustrate the experiment.

A half-pint glass tumbler, containing two ounces and a half of the above-mentioned diluted mixture of acids, being immersed in mixtures of nitrous acid and snow, until the liquor it contained was cooled to  $-30^{\circ}$ , was removed out of the mixture and placed upon a table; snow, likewise previously cooled in a frigorific mixture to  $-15^{\circ}$ , was added by degrees to the liquor in the tumbler, and the mixture kept stirring until a mercurial thermometer sunk to  $-60^{\circ}$ , where it remained stationary; the hydrometer was then immersed in the mixture (the thermometer glass having been broken in the course of the experiment), and stirred about in it for a short time, and on taking the hydrometer out, and gently shaking it, I  
perceived

perceived the mercury had already acquired the consistence of an amalgam, and after immersing it again for a few minutes, and then taking out and inverting it, I was gratified for the first time with the sight of mercury in a state of perfect congelation. I applied my hand to the inverted glass bulb; this soon loosened the solid mercury, which, on shaking the hydrometer, was distinctly heard to knock with force against the glass; it was then immersed a second time, and when taken out was found adhering to the glass as before. I now inverted the glass again, and kept it in that situation until the whole of the mercury melted, and dropped down globule after globule into the stem of the hydrometer. The interval of time from taking the mercury out of the frigorific mixture in a solid state, the last time, to its perfect liquefaction, was not noticed; but, upon recollection immediately afterwards, was supposed to be not less than three or four minutes. In a succeeding experiment this circumstance was attended to, and the frozen mercury, weighing seven scruples, was not entirely melted under seven minutes, the temperature of the air  $+30^{\circ}$ .

The experiment which follows I consider the most extraordinary, because it proves beyond a doubt, that mercury may be frozen not only here in summer, but even in the hottest climate, at any season of the year, by a combination of frigorific mixtures, in the way described in the Philosophical Transactions, Vol. LXXVII. p. 285. in which attempt to freeze mercury, made April 20, 1787, the temperature of the air and materials being  $+45^{\circ}$ , I certainly reached (without the assistance of snow or ice) the point of mercurial congelation; but had then no satisfactory proof that any part of the mercury was absolutely congealed.

Exp.

Exp. 2. On December 30. three ounces of a mixture composed of strong fuming nitrous acid two parts, and strong vitriolic acid and water each one part, were cooled in a half pint tumbler immersed in a frigorific mixture, till the temperature of the diluted mixture of acids was reduced to  $-30^{\circ}$ . The tumbler was then removed out of the mixture, and vitriolated natron (GLAUBER's salt) in very fine powder, previously cooled to  $-14^{\circ}$  by a frigorific mixture, added by degrees to the liquor in the tumbler, stirring it together until the mercury in the thermometer sunk to  $-54^{\circ}$ . The hydrometer used in the former experiment, with its lower bulb three-fourths full of mercury, was now immersed and stirred about in the mixture for a few minutes, when on taking it out, and inverting it, I had the satisfaction to find the same proof of the mercury being frozen as in the former instance. It was immediately shewn to the gentlemen present, who expressed likewise their entire satisfaction. Nearly four ounces of the powdered salt was added; but, I believe, some was added after the greatest effect was produced. I had no nitrated ammonia by me, otherwise I should have used upon this occasion, instead of vitriolated natron alone, a mixture of these two salts in powder, in the proportion of seven parts of the former to eight of the latter. The temperature of the room in which these experiments were made was  $+30^{\circ}$  each time, and the mercury taken from a jar containing several pounds.

Exp. 3. By an experiment made purposely on January 10. last, at which Dr. BOURNE was present, I have found that mercury may be congealed tolerably hard, by adding fresh fallen snow, at the temperature of  $+32^{\circ}$ , to strong fuming nitrous acid, previously cooled to between  $-25^{\circ}$  and  $-30^{\circ}$ , which may be very easily and quickly effected by immersing



the vessel containing the acid in a mixture of snow and nitrous acid.

I use the *fuming* nitrous acid upon all occasions, because that does not require to be diluted, cold being immediately produced on the smallest addition of snow.

Exp. 4. On January 12, at Dr. THOMSON'S request, I repeated the experiment of freezing mercury, at the Anatomy School in Christ Church, in the presence of the honourable Mr. WENMAN, the rev. Dr. HOARE, Dr. SIBTHORP, junior, Dr. THOMPSON, the rev. Mr. JACKSON of Christ Church, and Mr. WOOD of this place, a gentleman well known for his ingenuity in mechanics.

For this purpose were provided a spirit thermometer graduated very low, and a mercurial thermometer graduated to  $-76^{\circ}$ , two thermometer glasses, with bulbs very near, if not quite, an inch in diameter each, one filled with mercury nearly to the orifice of the tube, which was left open, the other with its bulb half filled, and an hydrometer with its lower bulb (considerably less than either of the others) likewise half filled with mercury; the temperature of the room at this time  $+28^{\circ}$ .

A pan, containing nine ounces of the mixture of acids prepared as in the first experiment, was placed in a larger pan, containing nitrous acid, and this, in a frigorific mixture of nitrous acid and snow, contained in another pan much larger. When the nitrous acid in the second pan was cooled by this mixture to  $-18^{\circ}$ , and the mixed acids in the smallest pan nearly as much, snow at somewhat between  $+20^{\circ}$  and  $+25^{\circ}$ , the temperature of the open air at that time, was added to the nitrous acid in the second pan, until the spirit thermometer sunk to near  $-43^{\circ}$ ; then the thermometer, with its bulb half filled, was immersed a sufficient time, and when taken  
out,

out, the mercury in it was found congealed, and adhering to the glass. The pan containing the mixed acids, and which had been removed whilst the snow was added to make the second mixture, was now replaced in it, in order to be cooled; and when the mixture of acids was reduced to the temperature of  $-34^{\circ}$ , snow previously cooled to  $-18^{\circ}$  was added, keeping the mixture stirred until the mercurial thermometer sunk to  $-60^{\circ}$ ; its temperature by the spirit thermometer was then found to be  $-51^{\circ}$ .

The three glasses containing the mercury to be frozen were now immersed in this mixture, and having been moved about in it for a considerable time, during which the spirit thermometer rose scarcely one degree, were then severally taken out and examined.

As the examination of the frozen mercury was more immediately under the inspection of Dr. THOMSON, I shall transcribe here that gentleman's account of the phenomena.

“When the freezing mixture was supposed to have produced its effect, the bulb which was completely filled was taken out, and broken on a flat stone by a moderate stroke or two with an iron hammer. This bulb was eleven or twelve lines in diameter.

The solid mercury was separated into several sharp and brilliant fragments, some of which bore handling for a short time before they returned to a fluid form. One mass, larger than the rest, consisting of nearly one-third of the whole ball, afforded the beautiful appearance of flat plates, converging towards a center. Each of these plates was about a line in breadth at the external surface of the ball, becoming narrower as it shot inwards. These facets lay in very different planes, as is common in the fracture of any crystallized ball, whether

whether of a brittle metal or of the earths, as in balls of calcareous stalactite. The solid brittle mercury in the present instance bore a very exact resemblance, both in colour and plated structure, to sulphurated antimony, and especially to the radiated specimens from Auvergne, before they are at all tarnished.

Instead of a solid center to this ball, it seemed as if there had been a central cavity, of about two lines in diameter, a considerable portion of which was evident in the fragment just described, at that part to which the radii converged. It is indeed possible, that this may have been merely the receptacle of some part of the mercury remaining fluid at the center. The hollow within was shining, but its edges were neither soft nor mouldering; on the contrary, they were sharp and well defined: nor was the brilliancy of the radii attributable to any exudation of mercury as from an amalgam.

In the two smaller bulbs, which were only half filled, the mercury preserved its usual lustre on the surface in contact with the glass, as well as on that surface which it had acquired in becoming solid. The latter was occupied by a conical depression, the gradations of which were marked by concentric lines.

One of these hemispheres was struck with a hammer, as in the former instance, but was rather flattened and crushed than broken. The other, on being divided with a sharp chisel, shewed a metallic splendour on its cut surface, but not equaling the polish of a globule of fluid mercury."

Thirteen ounces of snow in the whole were found to have been added to the mixed acids; but some was added to lower its temperature after the glasses containing the mercury were taken out, and the spirit thermometer had risen a few degrees.

This was a day remarkably favourable for such an experiment. My thermometer exposed to the open air stood, at three quarters past eight this morning, at  $+6^{\circ}$ , which is a very extraordinary degree of cold here; but this experiment was not begun till noon.

Exp. 5. On Jan. 14. I froze mercury at the Anatomy School again, in the presence of the rev. the Dean of Christ Church, the rev. Dr. HORNSBY, and Dr. THOMSON.

Four ounces now of the mixture of acids, prepared as in the first experiment, were cooled in a tumbler to  $-20^{\circ}$ , which required somewhat more than an equal weight of snow, cooled nearly to the same temperature, to produce the greatest effect. This was somewhat less than in the last experiment, the spirit thermometer sinking no lower than  $-46^{\circ}$ , owing chiefly to the weather having become much warmer, the temperature of the open air being now  $+36^{\circ}$ . The mercurial thermometer immersed in this mixture sunk to  $-55^{\circ}$ , where it became stationary; then two thermometer glasses, one half filled with mercury, and the other filled to a considerable height up the tube, after being immersed some time, were examined. Upon breaking the shell of glass from the former of these, the mercury was found in a perfectly solid state; but its upper surface, which was highly polished, and of the colour of liquid mercury, instead of being only slightly depressed, as had been seen in every other instance which afforded an opportunity for inspection, now formed a perfectly inverted hollow cone. This great depression, as well as the concentric circles mentioned in a former instance, I suppose, might be owing to a rotatory motion accidentally given to it whilst congealing. The solid mercury was beaten out, but having been suffered to lie some time on the table for inspection, very quickly melted

melted into liquid globules. The flexibility of solid mercury was clearly to be observed in this beautiful specimen; for the external surface, particularly the upper thin rim of the concave part, was evidently bent by the first gentle stroke of the hammer. The globe of mercury in the other glass, which was very small, exhibited nearly the same phenomena, as in the instances before mentioned.

It happened in these experiments of mine, contrary to what has generally occurred to others, that the mercury never sunk lower than  $-60^{\circ}$ , seldom so low, in the thermometer, and but little below the point of mercurial congelation in the tubes of the thermometer glasses filled nearly up to the orifice, with a view to shew the contraction of mercury in becoming solid by its great descent in the tube. On reflecting on this circumstance afterwards, it occurred to me, that the further descent of the mercury in these experiments was prevented not solely by the mercury freezing in the tube, the cause commonly assigned, but rather by the quick formation of a spherical shell of solid mercury within the bulb, by the sudden generation of cold.

Dr. BEDDOES expressing a desire to exhibit solid mercury at his Lecture before his Class, I undertook to freeze some at the Laboratory on March 12th last, and now resolved to satisfy myself respecting the cause which prevented the lower descent of the mercury in my former experiments. In this, as well as the former, the mercury in a thermometer graduated to  $-60^{\circ}$ , and likewise in a thermometer glass, filled nearly to the orifice, which lengthened its scale to near  $-250^{\circ}$ , sunk only a few degrees below the point of mercurial congelation, and then remained stationary. After waiting some time, I took the thermometer out of the mixture, and observed the bulb apparently full, and the short thread of mercury above unbroken.

I now embraced the lower part of the tube with my hand a few seconds, resting it upon the upper part of the bulb; and upon taking it away, I found that the whole of the mercury had subsided into the bulb, which it did not now quite fill, a small space at the top of the bulb remaining empty. I then took out the thermometer glass, and applied my hand to the tube; but the mercury remained stationary until I sunk my hand so as to communicate heat to that part of the bulb which is immediately connected with the tube, when the thread of mercury dropped entirely into the bulb. It was now immersed again for a short time, then taken out, and the shell of glass beaten off, which exposed a globe of solid mercury, nearly an inch in diameter. This bore several very smart strokes with a hammer before it began to liquify, but was not perfectly malleable.

In the course of these experiments, several fragments of the solid mercury were thrown into mercury in its ordinary liquid state, and were found to sink with considerable celerity.

In continuing my researches respecting the means of producing artificial cold, I have found that phosphorated natron produces rather more cold by solution in the diluted nitrous acid than the vitriolated natron.

At the temperature of  $+50^{\circ}$ , four parts of the diluted nitrous acid (prepared by mixing strong nitrous acid with half its weight of water) required eight parts of that neutral salt in fine powder to be added; in order to cause the thermometer to sink to  $-6^{\circ}$ ; and again, by the addition of five parts of nitrated ammonia in fine powder, the thermometer sunk so low as  $-16^{\circ}$ ; in the whole sixty-six degrees.

A mixture of this kind made the thermometer sink from  $80^{\circ}$  (the temperature of the materials before mixing) to  $0^{\circ}$ .

I was

I was directed to the trial of this salt, by the like remarkable sensation of coldness without pungency, which, with its other similar properties to ice, first induced me, whilst pursuing the subject of cold, to try the effect of dissolving the vitriolated natron in the mineral acids.

Equal quantities, by weight, of phosphorated natron and vitriolated natron, were evaporated separately over a gentle fire, until each was reduced to a perfectly dry powder. I then weighed them, and found the residuum of the phosphorated natron somewhat lighter than that of the vitriolated natron; from whence it is probable the former contains the greatest quantity of water of crystallization.

I have found, that each of the neutral salts which produce any remarkable degree of cold by solution in the mineral acids, *viz.* phosphorated natron, vitriolated natron, and vitriolated magnesia, lose this property entirely, when deprived by any means of their water of crystallization.

A short time after I had first succeeded in freezing water in summer, by one mixture composed of three different salts in water (having been induced to try the effect of such a method, from the consideration that water, already saturated with one kind of salt, will dissolve a portion of another, and after that a third, or even more), I met with the account of an experiment made by M. HOMBERG, related in one of the earlier Volumes of the Philosophical Transactions, in which it is said he produced an extraordinary degree of cold, by pouring a pint and a half of distilled vinegar upon two pounds of a powder composed of equal parts of crude sal ammoniac and corrosive sublimate, and shaking them well together. I immediately (July 30, 1786) prepared a mixture of this kind in smaller quantity, but found it produced only thirty-two degrees of cold,

cold, the temperature of the air and materials before mixing being  $63^{\circ}$ ; which is no more than I have found may be effected by a solution in water of crude sal ammoniac alone, previously dried and powdered.

By a trial made with great accuracy, I find, that even the mixture composed of diluted vitriolic acid and vitriolated natron is adequate to any useful purpose that may be required in the hottest country; for, by adding eleven parts of the salt in fine powder to eight parts of the vitriolic acid diluted with an equal weight of water, the thermometer sunk from  $80^{\circ}$ , the mean temperature of the hottest climate, and to which these materials were purposely heated before mixing, to rather below  $20^{\circ}$ .

Vitriolated natron, added to the marine acid undiluted, produces very nearly as great a degree of cold as when mixed with the diluted nitrous acid. At the temperature of  $50^{\circ}$ , two parts of the acid, require three parts of the salt in fine powder, which will sink the thermometer to  $0^{\circ}$ ; and if three parts of a mixed powder, containing equal parts of muriated ammonia and nitrated kali, be added afterwards, the cold of the mixture will be increased a few degrees more.

The frigorific mixture above described, composed of phosphorated natron and nitrated ammonia dissolved in the diluted nitrous acid, being the most powerful, it will probably be found most convenient for freezing mercury, when snow is not to be procured. The materials for this purpose may be previously cooled in mixtures made of marine acid with vitriolated natron, muriated ammonia, and nitrated kali, in the proportions mentioned above, this being much cheaper than those made with diluted nitrous acid, and very nearly equal in effect.



In my last Paper I mentioned a freezing mixture, made by dissolving a powder composed of equal parts of muriated ammonia and nitrated kali in water, and therein directed six parts of the mixed powder to be added to eight parts of water; but I have found since, that the best proportions are, five parts of the former to eight of the latter, by which I have sunk the thermometer from  $50^{\circ}$  to  $11^{\circ}$ .

Having now prosecuted my subject relative to mixtures for generating artificial cold without the use of ice, from a possible method proposed by Dr. WATSON (Essays, Vol. III. p. 139.), for freezing water in summer in this climate, and carried it on to a certain method of freezing, not only water, but even mercury, in the hottest climate, I now intend to take my leave of it.

I have the honour to be, &c.

RICHARD WALKER.



XX. *Catalogue of a second Thousand of new Nebulæ and Clusters of Stars; with a few introductory Remarks on the Construction of the Heavens.* By William Herschel, L L. D. F. R. S.

Read June 11, 1789.

**B**Y the continuation of a review of the heavens with my twenty-foot reflector, I am now furnished with a second thousand of new Nebulæ.

These curious objects, not only on account of their number, but also in consideration of their great consequence, as being no less than whole sidereal systems, we may hope, will in future engage the attention of Astronomers. With a view to induce them to undertake the necessary observations, I offer them the following catalogue, which, like my former one, of which it is a continuation, contains a short description of each nebula or cluster of stars, as well as its situation with respect to some known object.

The form of this work, it will be seen, is exactly that of the former part, the classes and numbers being continued, and the same letters used to express, in the shortest way, as many essential features of the objects as could possibly be crowded into so small a compass as that to which I thought it expedient to limit myself.

The method I have taken of *analyzing* the heavens, if I may so express myself, is perhaps the only one by which we can  
arrive

arrive at a knowledge of their construction. In the prosecution of so extensive an undertaking, it may well be supposed that many things must have been suggested, by the great variety in the order, the size, and the compression of the stars, as they presented themselves to my view, which it will not be improper to communicate.

To begin our investigation according to some order, let us depart from the objects immediately around us to the most remote that our telescopes, of the greatest *power to penetrate into space*, can reach. We shall touch but slightly on things that have already been remarked.

From the earth, considered as a planet, and the moon as its satellite, we pass through the region of the rest of the planets, and their satellites. The similarity between all these bodies is sufficiently striking to allow us to comprehend them under one general definition, of bodies not luminous in themselves, revolving round the sun. The great diminution of light, when reflected from such bodies, especially when they are also at a great distance from the light which illuminates them, precludes all possibility of following them a great way into space. But if we did not know that light diminishes as the squares of the distances encrease, and that moreover in every reflection a very considerable part is intirely lost, the motion of comets, whereby the space through which they run is measured out to us, while on their return from the sun we see them gradually disappear as they advance towards their aphe-  
lia, would be sufficient to convince us that bodies shining only with borrowed light can never be seen at any very great distance. This consideration brings us back to the sun, as a refulgent fountain of light, whilst it establishes at the same time beyond a doubt that every star must likewise be a sun,

shining by its own native brightness. Here then we come to the more capital parts of the great construction.

These suns, every one of which is probably of as much consequence to a system of planets, satellites, and comets, as our own sun, are now to be considered, in their turn, as the minute parts of a proportionally greater whole. I need not repeat that by my analysis it appears, that the heavens consist of regions where suns are gathered into separate systems, and that the catalogues I have given comprehend a list of such systems; but may we not hope that our knowledge will not stop short at the bare enumeration of phenomena capable of giving us so much instruction? Why should we be less inquisitive than the natural philosopher, who sometimes, even from an inconsiderable number of specimens of a plant, or an animal, is enabled to present us with the history of its rise, progress, and decay? Let us then compare together, and class some of these numerous sidereal groups, that we may trace the operations of natural causes as far as we can perceive their agency. The most simple form, in which we can view a sidereal system, is that of being globular. This also, very favourably to our design, is that which has presented itself most frequently, and of which I have given the greatest collection.

But, first of all, it will be necessary to explain what is our idea of a cluster of stars, and by what means we have obtained it. For an instance, I shall take the phenomenon which presents itself in many clusters: It is that of a number of lucid spots, of equal lustre, scattered over a circular space, in such a manner as to appear gradually more compressed towards the middle; and which compression, in the clusters to which I allude, is generally carried so far, as, by imperceptible degrees,

to end in a luminous center, of a resolvable blaze of light. To solve this appearance, it may be conjectured, that stars of any given, very unequal magnitudes, may easily be so arranged, in scattered, much extended, irregular rows, as to produce the above described picture; or, that stars, scattered about almost promiscuously within the frustum of a given cone, may be assigned of such properly diversified magnitudes as also to form the same picture. But who, that is acquainted with the doctrine of chances, can seriously maintain such improbable conjectures? To consider this only in a very coarse way, let us suppose a cluster to consist of 5000 stars, and that each of them may be put into one of 5000 given places, and have one of 5000 assigned magnitudes. Then, without extending our calculation any further, we have five and twenty millions of chances, out of which only one will answer the above improbable conjecture, while all the rest are against it. When we now remark that this relates only to the given places within the frustum of a supposed cone, whereas these stars might have been scattered all over the visible space of the heavens; that they might have been scattered, even within the supposed cone, in a million of places different from the assumed ones, the chance of this apparent cluster's not being a real one, will be rendered so highly improbable that it ought to be intirely rejected.

Mr. Michell computes, with respect to the six brightest stars of the Pleiades only, that the odds are near 500 000 to 1 that no six stars, out of the number of those which are equal in splendour to the faintest of them, scattered at random in the whole heavens, would be within so small a distance from each other as the Pleiades are\*.

\* Phil. Transf. vol. LVII, p. 246.

Taking it then for granted that the stars which appear to be gathered together in a group are in reality thus accumulated, I proceed to prove also that they are nearly of an equal magnitude.

The cluster itself, on account of the small angle it subtends to the eye, we must suppose to be very far removed from us. For, were the stars which compose it at the same distance from one another as Sirius is from the sun; and supposing the cluster to be seen under an angle of 10 minutes, and to contain 50 stars in one of its diameters, we should have the mean distance of such stars twelve seconds; and therefore the distance of the cluster from us about seventeen thousand times greater than the distance of Sirius. Now, since the apparent magnitude of these stars is equal, and their distance from us is also equal,—because we may safely neglect the diameter of the cluster, which, if the center be seventeen thousand times the distance of Sirius from us, will give us seventeen thousand and twenty-five for the farthest, and seventeen thousand wanting twenty-five for the nearest star of the cluster;—it follows that we must either give up the idea of a cluster, and recur to the above refuted supposition, or admit the equality of the stars that compose these clusters. It is to be remarked that we do not mean intirely to exclude all variety of size; for the very great distance, and the consequent smallness of the component clustering stars, will not permit us to be extremely precise in the estimation of their magnitudes; though we have certainly seen enough of them to know that they are contained within pretty narrow limits; and do not, perhaps, exceed each other in magnitude more than in some such proportion as one full-grown plant of a certain species may exceed another full-grown plant of the same species.

If we have drawn proper conclusions relating to the size of stars, we may with still greater safety speak of their relative situations, and affirm that in the same distances from the center an equal scattering takes place. If this were not the case, the appearance of a cluster could not be uniformly encreasing in brightness towards the middle, but would appear nebulous in those parts which were more crowded with stars; but, as far as we can distinguish, in the clusters of which we speak, every concentric circle maintains an equal degree of compression, as long as the stars are visible; and when they become too crowded to be distinguished, an equal brightness takes place, at equal distances from the center, which is the most luminous part.

The next step in my argument will be to shew that these clusters are of a globular form. This again we rest on the found doctrine of chances. Here, by way of strength to our argument, we may be allowed to take in all round *nebulæ*, though the reasons we have for believing that they consist of stars have not as yet been entered into. For, what I have to say concerning their spherical figure will equally hold good whether they be groups of stars or not. In my catalogues we have, I suppose, not less than one thousand of these round objects. Now, whatever may be the shape of a group of stars, or of a *Nebula*, which we would introduce instead of the spherical one, such as a cone, an ellipsis, a spheroid, a circle or a cylinder, it will be evident that out of a thousand situations, which the axes of such forms may have, there is but one that can answer the phænomenon for which we want to account; and that is, when those axes are exactly in a line drawn from the object to the place of the observer. Here again we have a million of chances of which all but one are against any other

hypothesis than that which we maintain, and which, for this reason, ought to be admitted.

The last thing to be inferred from the above related appearances is, that these clusters of stars are more condensed towards the center than at the surface. If there should be a group of stars in a spherical form, consisting of such as were equally scattered over all the assigned space, it would not appear to be very gradually more compressed and brighter in the middle; much less would it seem to have a bright nucleus in the center. A spherical cluster of an equal compression within,—for that such there are will be seen hereafter,—may be distinguished by the degrees of brightness which take place in going from the center to the circumference. Thus, when  $a$  is the brightness in the center, it will be  $\sqrt{a^2 - x^2}$  at any other distance  $x$  from the center. Or, putting  $a = 1$ , and  $x =$  any decimal fraction; then, in a table of natural sines, where  $x$  is the sine, the brightness at  $x$  will be expressed by the cosine. Now, as a gradual increase of brightness does not agree with the degrees calculated from a supposition of an equal scattering, and as the cluster has been proved to be spherical, it must needs be admitted that there is indeed a greater accumulation towards the center. And thus, from the above-mentioned appearances, we come to know that there are globular clusters of stars nearly equal in size, which are scattered evenly at equal distances from the middle, but with an increasing accumulation towards the center.

We may now venture to raise a superstructure upon the arguments that have been drawn from the appearance of clusters of stars and nebulae of the form I have been examining, which is that of which I have made mention in my “*Theoreti-*



*cal view—Formation of Nebulæ—Form I\*.*” It is to be remarked that when I wrote the paragraph I refer to, I delineated nature as well as I do now; but, as I there gave only a general sketch, without referring to particular cases, what I then delivered may have been looked upon as little better than hypothetical reasoning, whereas in the present instance this objection is intirely removed, since actual and particular facts are brought to vouch for the truth of every inference.

Having then established that the clusters of stars of the 1st Form, and round nebulæ, are of a spherical figure, I think myself plainly authorized to conclude that they are thus formed by the action of central powers. To manifest the validity of this inference, the figure of the earth may be given as an instance; whose rotundity, setting aside small deviations, the causes of which are well known, is without hesitation allowed to be a phænomenon decisively establishing a centripetal force. Nor do we stand in need of the revolving satellites of Jupiter, Saturn, and the Georgium Sidus, to assure us that the same powers are likewise lodged in the masses of these planets. Their globular figure alone must be admitted as a sufficient argument to render this point uncontrovertible. We also apply this inference with equal propriety to the body of the sun, as well as to that of Mercury, Venus, Mars, and the Moon; as owing their spherical shape to the same cause. And how can we avoid inferring, that the construction of the clusters of stars, and nebulæ likewise, of which we have been speaking, is as evidently owing to central powers?

Besides, the step that I here make in my inference is in fact a very easy one, and such as ought freely to be granted. Have I not already shewn that these clusters cannot have come to

\* Phil. Transf. vol. LXXV, p. 214.

their present formation by any random scattering of stars? The doctrine of chance, by exposing the very great odds against such hypotheses, may be said to demonstrate that the stars are thus assembled by some power or other. Then, what do I attempt more than merely to lead the mind to the conditions under which this power is seen to act?

In a case of such consequence I may be permitted to be a little more diffuse, and draw additional arguments from the internal construction of spherical clusters and nebulæ. If we find that there is not only a general form, which, as has been proved, is a sufficient manifestation of a centripetal force, what shall we say when the accumulated condensation, which every where follows a direction towards a center, is even visible to the very eye? Were we not already acquainted with attraction, this gradual condensation would point out a central power, by the remarkable disposition of the stars tending towards a center. In consequence of this visible accumulation, whether it may be owing to attraction only, or whether other powers may assist in the formation, we ought not hesitate to ascribe the effect to such as are *central*; no phænomena being more decisive in that particular, than those of which I am treating.

I am fully aware of the consequences I shall draw upon myself in but mentioning other powers that might contribute to the formation of clusters. A mere hint of this kind, it will be expected, ought not to be given without sufficient foundation; but let it suffice at present to remark that my arguments cannot be affected by my terms: whether I am right to use the plural number,—central powers,—or whether I ought only to say,—the known central force of gravity,—my conclusions will be equally valid. I will however add, that the idea of other

central powers being concerned in the construction of the sidereal heavens, is not one that has only lately occurred to me. Long ago I have entertained a certain theory of diversified central powers of attractions and repulsions; an exposition of which I have even delivered in the years 1780, and 1781, to the Philosophical Society then existing at Bath, in several mathematical papers upon that subject. I shall, however, set aside an explanation of this theory, which would not only exceed the intended limits of this paper, but is moreover not required for what remains at present to be added, and therefore may be given some other time, when I can enter more fully into the subject of the interior construction of sidereal systems.

To return, then, to the case immediately under our present consideration, it will be sufficient that I have abundantly proved that the formation of round clusters of stars and nebulae is either owing to central powers, or at least to one such force as refers to a center.

I shall now extend the weight of my argument, by taking in likewise every cluster of stars or nebula that shews a gradual condensation, or encreasing brightness, towards a center or certain point; whether the outward shape of such clusters or nebulae be round, extended, or of any other given form. What has been said with regard to the doctrine of chance, will of course apply to every cluster, and more especially to the extended and irregular shaped ones, on account of their greater size: It is among these that we find the largest assemblages of stars, and most diffusive nebulosities; and therefore the odds against such assemblages happening without some particular power to gather them, encrease exceedingly with the number of the stars that are taken together. But if the gradual accumulation either of stars or encreasing brightness has before

been admitted as a direction to the seat of power, the same effect will equally point out the same cause in the cases now under consideration. There are besides some additional circumstances in the appearance of extended clusters and nebulae, that very much favour the idea of a power lodged in the brightest part. Although the form of them be not globular, it is plainly to be seen that there is a tendency towards sphericity, by the swell of the dimensions the nearer we draw towards the most luminous place, denoting as it were a course, or tide of stars, setting towards a center. And—if allegorical expressions may be allowed—it should seem as if the stars thus flocking towards the seat of power were stemmed by the crowd of those already assembled, and that while some of them are successful in forcing their predecessors sideways out of their places, others are themselves obliged to take up with lateral situations, while all of them seem equally to strive for a place in the central swelling, and generating spherical figure.

Since then almost all the nebulae and clusters of stars I have seen, the number of which is not less than three and twenty hundred, are more condensed and brighter in the middle; and since, from every form, it is now equally apparent that the central accumulation or brightness must be the result of central powers, we may venture to affirm that this theory is no longer an unfounded hypothesis, but is fully established on grounds which cannot be overturned.

Let us endeavour to make some use of this important view of the constructing cause, which can thus model sidereal systems. Perhaps, by placing before us the very extensive and varied collection of clusters, and nebulae furnished by my catalogues, we may be able to trace the progress of its operation, in the great laboratory of the Universe.

If these clusters and nebulæ were all of the same shape, and had the same gradual condensation, we should make but little progress in this inquiry; but, as we find so great a variety in their appearances, we shall be much sooner at a loss how to account for such various phenomena, than be in want of materials upon which to exercise our inquisitive endeavours.

Some of these round clusters consist of stars of a certain magnitude, and given degree of compression, while the whole cluster itself takes up a space of perhaps 10 minutes; others appear to be made up of stars that are much smaller, and much more compressed, when at the same time the cluster itself subtends a much smaller angle, such as 5 minutes. This diminution of the apparent size, and compression of stars, as well as diameter of the cluster to 4, 3, 2 minutes, may very consistently be ascribed to the different distances of these clusters from the place in which we observe them; in all which cases we may admit a general equality of the sizes, and compression of the stars that compose them, to take place. It is also highly probable that a continuation of such decreasing magnitudes, and encreasing compression, will justly account for the appearance of round, easily resolvable, nebulæ; where there is almost a certainty of their being clusters of stars. And no Astronomer can hesitate to go still farther, and extend his surmises by imperceptible steps to other nebulæ, that still preserve the same characteristics, with the only variations of vanishing brightness, and reduction of size.

Other clusters there are that, when they come to be compared with some of the former, seem to contain stars of an equal magnitude, while their compression appears to be considerably different. Here the supposition of their being at different distances will either not explain the apparently greater

compression, or, if admitted to do this, will convey to us a very instructive consequence: which is, that the stars which are thus supposed not to be more compressed than those in the former cluster, but only to appear so on account of their greater distance, must needs be proportionally larger, since they do not appear of less magnitude than the former. As therefore, one or other of these hypotheses must be true, it is not all improbable but that, in some instances, the stars may be more compressed; and in others, of a greater magnitude. This variety of size, in different spherical clusters, I am however inclined to believe, may not go farther than the difference in size, found among the individuals belonging to the same species of plants, or animals, in their different states of age, or vegetation, after they are come to a certain degree of growth. A farther inquiry into the circumstance of the extent, both of condensation and variety of size, that may take place with the stars of different clusters, we shall postpone till other things have been previously discussed.

Let us then continue to turn our view to the power which is moulding the different assortments of stars into spherical clusters. Any force, that acts uninterruptedly, must produce effects proportional to the time of its action. Now, as it has been shewn that the spherical figure of a cluster of stars is owing to central powers, it follows that those clusters which, *ceteris paribus*, are the most compleat in this figure, must have been the longest exposed to the action of these causes. This will admit of various points of views. Suppose for instance that 5000 stars had been once in a certain scattered situation, and that other 5000 equal stars had been in the same situation, then that of the two clusters which had been longest exposed to the action of the modelling power, we suppose,

would be most condensed, and more advanced to the maturity of its figure. An obvious consequence that may be drawn from this consideration is, that we are enabled to judge of the relative age, maturity, or climax of a sidereal system, from the disposition of its component parts; and, making the degrees of brightness in nebulæ stand for the different accumulation of stars in clusters, the same conclusions will extend equally to them all. But we are not to conclude from what has been said that every spherical cluster is of an equal standing in regard to absolute duration, since one that is composed of a thousand stars only, must certainly arrive to the perfection of its form sooner than another, which takes in a range of a million. Youth and age are comparative expressions; and an oak of a certain age may be called very young, while a contemporary shrub is already on the verge of its decay. The method of judging with some assurance of the condition of any sidereal system may perhaps not improperly be drawn from the standard laid down page 218; so that, for instance, a cluster or nebula which is very gradually more compressed and bright towards the middle, may be in the perfection of its growth, when another which approaches to the condition pointed out by a more equal compression, such as the nebulæ I have called *Planetary* seem to present us with, may be looked upon as very aged, and drawing on towards a period of change, or dissolution. This has been before surmised, when, in a former paper, I considered the uncommon degree of compression that must prevail in a nebula to give it a planetary aspect; but the argument, which is now drawn from the powers that have collected the formerly scattered stars to the form we find they have assumed, must greatly corroborate that sentiment.

This

This method of viewing the heavens seems to throw them into a new kind of light. They now are seen to resemble a luxuriant garden, which contains the greatest variety of productions, in different flourishing beds; and one advantage we may at least reap from it is, that we can, as it were, extend the range of our experience to an immense duration. For, to continue the simile I have borrowed from the vegetable kingdom, is it not almost the same thing, whether we live successively to witness the germination, blooming, foliage, fecundity, fading, withering, and corruption of a plant, or whether a vast number of specimens, selected from every stage through which the plant passes in the course of its existence, be brought at once to our view?

WILLIAM HERSCHEL.

Slough near Windfor, May 1, 1789.

First Class. Bright nebulae.

I.	1785	Stars.	M.	S.	D.M.	Ob.	Description.
94	April 28	61 Ursa	f	0 6	n	2 17 2	cB. pL. E. spnf. vgmbM. 3½ l. 2' b.
95	—	—	f	35 0	n	2 7 2	cB. cL. E. np ff. bM. 4' l. 3' b.
96	May 1	14 Canum.	f	5 30	n	1 12 2	vB. cL. mE. sp nf. fmbM. 6' l. 1½ b.
97	—	—	f	7 58	n	0 47 1	vB. pL. E. nearly mer. gmbM.
98	—	—	f	36 50	f	0 12 1	cB. pL. R. vgmbM.
99	—	27 (γ) Bootis	p	13 46	f	1 46 2	vB. S. R. vfmB.
100	Sept. 10	41 Ceti	f	13 43	n	0 48 1	cB. pS. R. mbM. See III. 431.
101	—	67 —	p	17 19	n	0 25 2	cB. pL. E. near. mer. mbM. 5' l.
102	—	—	f	21 37	f	0 13 2	cB. pL. R. mbM.
103	24	14 Delphini	p	16 10	f	0 3 1	vB. L. gmbM. er. beautif. object.
104	28	93 (ψ) Aqua	f	1 8	n	0 42 1	cB. cL. E. near. mer. gmbM. F. rays.
105	Oct. 3	47 Ceti	f	26 24	f	0 37 1	cB. pL. iR. mbM.
106	—	89 (π) —	f	38 10	f	1 24 2	cB. cL. iR. bM. 3' dia.
107	6	20 Eridani	f	4 3	f	1 4 2	vB. R. BNM. 1½ dia.
108	8	111 (ξ) Pisc <sup>m</sup>	p	34 22	f	0 1 1	cB. vL. iR. p. vBft.
109	26	12 Eridani	p	7 17	n	2 54 3	cB. pS. lE. mer. mbM. r. 1½ l.



I.	1785	Stars.	M.	S.	D.M.	Ob	Description.
110	Nov. 27	9 Ceti	p	44 0	f 0 47	2	cB. cL. lE. gmbM. iF.
111	—	—	p	43 3	f 0 6	2	cB. cL. iR. gmbM.
112	29	5( $\gamma$ ) Arietis	f	5 48	f 0 17	1	vB. L. R. mbM. not er. 4' dia.
113	Dec. 7	60(4th $\sigma$ ) Can	f	18 22	n 1 34	2	cB. cL. lE. iF. mb foll. side.
114	—	18 Leo. min.	p	13 39	f 0 35	1	cB. cL. iF. mbM.
115	—	—	p	5 47	n 1 10	2	cB. pL. lE. iF. mbM.
116	}	— 37 —	f	11 5	n 1 1	1	{ Two; the 1st, cB. cL. iE; the 2d, pB. pL. iE. Dist. 1' at the vertex.
117							
118	—	46 Urfæ	p	3 41	f 1 32	1	cB. cL. iR. mbM.
119	28	31 (1st $d$ ) Vir	p	6 0	n 0 55	1	vB. pS.
120	31	30(n) Crateri	p	9 0	n 0 17	1	cB. L. iR. bM. 5' l. 4' b.
1786							
121	Jan. 1	13(n) Virgin	p	18 15	f 0 19	1	vB. cL. lE. mbM. 3' l. 2' $\frac{1}{2}$ b. bet. 2pBst.
122	Feb. 1	57( $\mu$ ) Eridani	p	4 0	n 0 22	1	cB. vL. iR. bM. er. 5 or 6' dia.
123	2	60 ( $\sigma$ ) Virg.	p	52 27	f 0 30	2	cB. S.
124	—	—	p	39 57	f 0 3	2	cB. cL. R.
125	—	—	p	39 12	f 1 6	2	cB. cL. E. mbM.
126	24	108	p	0 35	n 1 15	1	eB. mE. par. BN. 8 or 9' l.
127	—	110	p	1 47	f 0 23	1	cB. pS. mbM.
128	—	—	f	3 37	f 0 30	1	vB. pL. bM.
129	March 3	26 ( $\chi$ )	f	9 46	f 0 41	1	v brilliant. iR. vgbM.
130	—	—	f	26 35	f 0 3	2	vB. lE. mer. BN. and F. br. 2' l.
131	4	14 ( $\epsilon$ ) Crate.	f	0 29	n 1 3	1	cB. E. gbM. 5' l. 4' b.
132	19	26 Hydræ	f	1 44	n 0 4	2	cB. pL. lE. vgbM. 1' $\frac{1}{2}$ diam.
133	25	49( $g$ ) Virgin.	p	16 4	n 0 18	1	cB. vS. BN.
134	—	—	p	13 27	n 0 13	1	cB. 7 or 8' l. 3' b.
135	}	27 68 ( $\iota$ )	p	32 2	n 0 11	2	{ Two; both cB. cS. R. mbM. Dist. 1' near. mer. chev. mixed.
136							
137	28	41 Lyncis.	f	3 13	n 0 8	1	vB. R. vsmB. chev. 3' dia.
138	—	*1102 ( $e$ ) Hy	f	33 45	n 1 27	1	cB. R. psmB. * See note.
139	April 17	11 ( $s$ ) Virgin.	f	12 1	f 1 21	2	eB. vBN. r. 6 or 7' dia.
140	—	—	f	39 55	f 0 31	2	cB. pL. mbM.
141	—	—	f	45 50	f 1 32	1	vB. cL. E. np ff.
142	30	37	p	6 35	n 0 0	1	cB. pL. iR. gmbM.
143	—	43 ( $\delta$ ) Virgin	f	4 55	f 2 7	1	cB. np. pBst. and close to it.
144	—	109	p	25 58	n 0 54	1	cB. cL. R. gmbM.
145	}	—	p	25 14	n 1 27	1	{ Two; the p.pB. pL. E. Dist. 3 or 4' fp nf. The f cB. R. pL. Place of 2d.
146							
147	—	43 Ophiuchi	p	8 54	f 1 17	1	vB. R. gmbM. 2' $\frac{1}{2}$ dia.
148	May 1	24 ( $\alpha$ ) Serpen	p	22 26	f 1 16	1	cB. cL. iR. bM.
149	28	40 ( $e$ ) Ophiu	f	0 14	n 1 32	1	cB. pS. lE. er.
150	—	—	f	27 53	n 0 36	1	cB. R. vgbM. about 1' $\frac{1}{2}$ dia.
151	Sept. 4	71 ( $\epsilon$ ) Piscium	f	21 41	n 1 41	1	cB. cL. R. C. vgbM. N.
152	—	24 ( $\xi$ ) Arietis	p	16 23	n 0 20	2	vB. vS. R. or lE. vBN. 1' ff. est

I.	1786	Stars.	M.	S.	D.M.		Ob	Description.				
153	Sept. 20	59 (2d v) Ceti	p	23	16	f	0	6	1	cB. vL. E. sp nf. above 15' l.		
154	21	14 Triang.	f	1	23	n	0	59	2	cB pL. E. np ff. vgmbM. 3' l. 2' b.		
155	30	32 Eridani	f	7	49	f	1	1	2	cB. S. gmbM.		
156	Oct. 18	12 (q) Persei	p	1	41	f	1	10	2	cB. mE. 12° sp nf. vBN. near 10' l.		
157	26	90 (v) Piscium	f	28	9	n	0	13	1	cB. cL. E. par. mbM. 7' l. 3' b.		
158	Nov. 26	48 (v) Eridani	p	4	32	f	1	46	2	cB. pL. iR. vgmbM.		
159	Dec. 11	20 (π) Cassiop	f	8	30	n	0	33	3	vB. R. vgbM. 1½ dia.		
160	29	29 (γ) Virgin	p	6	17	f	2	19	2	vB. cL. E. sp nf. vgBN. F. bran.		
1787												
161	Jan. 14	6 Comæ	f	12	58	f	0	55	1	vB. pL. iR.		
162	—	29 —	f	10	35	n	0	2	1	vB. E. sp nf. Sft in it ½' p. N.		
163	Feb. 22	20 Sextantis	p	8	29	f	0	22	1	{ eB. cL. mE. 45 sp nf. N. 2' l. F. br. 5' l.		
164	Mar. 17	38 Leo. min.	p	2	54	f	0	36	3	cB. E. 30° np ff. mbM. er. 4' l. 2' b.		
165	—	6 Canum	p	15	42	n	0	25	2	{ vB. BN. not M. or 2 joined the n. N.		
166	—	—	p	1	20	n	0	23	2	vB. S. R. mbM.		
167	18	10 (n) Urfæ	f	13	43	f	1	40	1	cB. R. BN. 1½ dia.		
168	—	34 (μ) —	p	4	9	f	0	6	3	{ cB. R. vgbM. 8' dia. cft. in it, unconnected.		
169	—	6 Canum	p	16	16	n	0	53	1	cB. cL.		
170	—	20 —	f	28	12	n	1	6	2	cB. E. near par. SNM. 2' l.		
171	—	53 (2d v) Boot	p	49	57	n	1	10	2	cB. S. R. r. mbM.		
172	19	31 Leo. min.	f	25	2	f	0	3	1	{ cB. E. sp nf. few ft. in p. 1 in n. unconnected.		
173	—	—	f	86	19	n	0	23	1	vB. R. vgnM. 2½ dia.		
174	20	53 (ξ) Urfæ	f	46	14	n	0	24	1	cB. E. 5' l. 1½ b.		
175	—	13 Canum	p	46	3	n	2	28	1	vB. S. R. mbM.		
176	}	—	p	16	33	n	1	26	1	{ Two. The f. cB. E. mbM. The n. pB. E. sp nf. Both join and form the letter S.		
177												
178	}	April 9	8	—	f	7	36	f	0	12	1	{ Two. The n. vB. vmbM. The f. pB. Their uebul. run together.
179												
180	—	20	—	f	29	9	n	3	15	1	cB. mE. 60° np ff. vIBM.	
181	—	—	—	f	40	13	n	1	11	1	cB. cL. mbM.	
182	11	1 Serpentis	p	17	22	f	0	2	2	cB. pL. iR. mbM.		
183	—	—	p	11	19	n	0	1	2	cB. pL. iR. or lE.		
184	May 7	8 Librae	p	8	21	f	1	15	1	cB. pL. E. sp nf. mbM.		
185	11	19 (λ) Bootis	f	11	6	n	0	1	2	c or pB S. R. pmbM.		
186	12	—	p	47	14	n	1	20	2	{ cB. pL. R or lE. vgbM. 3' np. the 51st of the <i>Conn. des Temps.</i>		
187	—	—	p	20	15	n	1	14	1	cB. E. 30 sp nf. BN. vgF. branches.		
188	—	38 (2db) —	p	13	24	n	2	44	2	cB. lE. par. mbM. F. bran. 1½ l.		
189	15	24 (g) —	f	3	57	f	0	23	1	cB. cL. E. sp nf. broad.		

I.	1787	Stars.		M. S.		D.M.	Ob	Description.	
190	May 16	*Canum 6m.	f	11 32	f	1 11	1	{ Two. The f. cB. cL. The n. pB. S. dist. 1' 1/2. * See note.	
191									
192			Oct. 14	3 Lacertæ	p	80 46	n		2 32
193	Nov. 12	54 (φ) Andro	p	1 26	n	0 54	1	{ Two close together. Both vB. dist. 2'. fp nf. One is 76 of the Conn.	
1788									
194	Jan. 14	56 Urfæ	f	3 19	n	0 5	2	vB. cL. mE. mer. BN. 6' l. 2' b. chev.	
195	—	67 —	f	4 49	n	0 2	2	E. vBN. and F. branches.	
196	—	—	f	7 17	n	0 38	2	cB. cL. iF. vgbM. ff. ft.	
197	—	8 Canum	p	3 32	n	0 19	1	{ Two. The f. vB. vL. iE. The n. B. pS. iF. dist. 1' 1/2.	
198									
199			15	15 Leo. min.	f	32 1	f		0 24
200	Feb. 5	59 (2d σ) Can	p	4 29	n	0 29	1	v brilliant. mE. fp nf. 8' l. 3' b. beauti.	
201	—	63 (x) Urfæ	f	0 5	f	0 17	2	cB. mE. fp nf. near. mer. 5' l. 1' b.	
202	—	—	f	0 47	n	0 4	2	cB. S. lE.	
203	—	659 —	f	7 42	n	0 31	1	cB. cL. R. pBNM.	
204	March 9	9 (i) —	p	16 27	n	2 7	1	cB. vS. lE. m.	
205	—	—	f	22 18	n	3 1	1	{ vB. lbM. chev. bran. m. neb. 6' l. 4' b.	
206	—	3 Canum	p	14 39	n	1 35	3	{ cB. E. 45° np ff. 6' l. 4' b. al- most equally B.	
207	—	—	p	14 0	f	1 32	3	cB. mE. 70° fp nf. 6 or 7' l. 2' b.	
208	—	—	p	9 9	n	1 32	3	cB. mE. fp nf. SBNM. 5' l. 1' b.	
209	—	—	p	3 33	f	1 6	2	cB. cL. E. mbM.	
210	April 1	60 Urfæ	f	40 0	n	0 9	2	vB. S. lE. near. par. BN. eF. bran.	
211	—	11 Canum	f	5 47	f	1 58	3	cB. S. R. bM. f. vSft.	
212	10	60 Urfæ	f	50 50	f	1 58	1	cB. pL. E.	
213	27	19 (λ) Bootis	p	110 25	f	1 48	1	{ v brilliant. cL. E. fp nf. difficulty r. has 3 or 4 BN.	
214	May 1	17 (x) —	p	8 26	n	1 56	1	cB. cL. n. ends abruptly. f. vg.	
215	5	Neb. II. 757.	p	3 27	f	1 14	1	vB. cL. E. f. 2 ft.	

Second class. Faint nebulæ.

II.	1785	Stars.		M. S.		D.M.	Ob	Description.
403	April 26	1 Comæ	p	8 50	f	1 21	3	F. cL. iF. lbM.
404	27	5 —	p	11 40	f	0 29	1	pB. pL. R. C. mbM.
405	—	—	p	1 0	f	0 24	2	pB. pL. iF. lE. bM. p. pcft.
406	—	20 —	p	6 8	f	1 27	1	{ pF. pL. mbM. S neb. joined to it, or lb. in the n.
407	—	—	f	6 44	f	1 35	1	pB. pS. lE.
408	28	61 Urfæ	f	7 54	n	0 46	2	F. S. R. gbM. near 1/2' dia.
409	May 1	—	f	33 54	n	2 25	2	pB. pL. vgbM. r.
410	—	14 Canum	p	32 8	f	0 14	2	pB. cL. R. smbM. r.

II.	1785.	Stars.		M.	S.	D.M.	Ob	Description.
411	May 1	14 Canum.	p	24	25	f	0 43	2 pB. pL. R. lbM. 2' np. pBft.
412	—	—	p	17	8	f	0 28	2 F. S. lE. glbM. er.
413	—	—	p	0	50	f	0 36	2 pB. S.R. bM. and vF. on the edges.
414	—	—	f	5	58	n	0 27	1 F. S. lE.
415	—	—	f	48	34	n	0 15	1 F. S. iF.
416	—	—	f	58	10	f	1 8	2 pB. pL. iE. mbM.
417	—	—	f	58	18	f	0 47	1 pB. pL. iE. bM.
418	—	51 ( $\mu$ ) Bootis	p	69	38	f	1 48	1 pB. iR. mbM.
419	—	—	p	68	31	f	0 37	1 F. pL.
420	—	—	p	61	32	f	2 17	1 pB. vS. R. vgmbM.
421	—	—	p	55	14	f	1 53	1 F. pL. iF.
422	—	—	p	52	36	i	0 52	1 F. cL. iF. unequally B.
423	—	—	p	47	57	f	0 37	1 pF. pS. iF. bM.
424	2	49 ( $\delta$ ) —	p	83	12	n	0 31	1 F. pL. lbM.
425	5	34 ( $\omega$ ) Serpen	p	4	0	n	0 15	3 F. cS. iR. stellar.
426	} Aug. 12	1 Aquarii	f	7	50	f	0 12	1 { Two. Thep.F.S.iR.mbM. The f.
427								
428	} —	6 ( $\gamma$ ) Piscium	p	2	16	n	1 14	1 { Two. The f. pB.mE. par. mbM.
429								
430	} —	6 ( $\gamma$ ) Piscium	p	2	16	n	1 14	1 { dist. and p.
431								
432	—	—	f	22	5	n	1 9	4 pB. cL. E. 75° sp nf. 3' l.
433	—	41 Ceti	p	18	0	f	0 4	1 pB. pL. bM. i. parallelogram. mer.
434	—	—	p	14	23	n	1 18	1 F. S. iF. bM. r..
435	—	67 —	p	15	52	f	0 27	1 F. S. iR. bM.
436	—	—	f	1	45	f	0 14	1 F. pS. lE. f. 2 or 3 uneq. ft.
437	—	—	f	2	7	f	0 24	1 F. pS. lE.
438	—	—	f	4	33	n	0 54	2 pB. vL. iF. mbM. r..
439	26	59 ( $\rho$ ) Pegasi	f	8	34	f	0 30	1 pB. pS. mbM.
440	—	—	f	9	1	f	0 30	1 pB. pS. bM.
441	—	—	f	10	1	n	0 10	1 F. S.
442	Oct. 1	62 ( $\eta$ ) Aqua	f	9	4	f	0 5	3 F. S. r. lbM. or f. M.
443	—	—	f	15	19	f	1 29	2 F. S. iR. lbM. 1' $\frac{1}{2}$ f. S. ft.
444	—	20 Ceti	p	10	20	f	0 24	1 F. pL. lbM.
445	—	—	p	6	50	f	0 35	1 F. iF. er. 1' b.
446	—	—	p	2	16	f	0 45	2 pB. S. R. mbM. m.
447	—	34 —	f	1	3	n	2 0	2 { F. S. Two more near it. See
448	} —	43 —	f	3	28	f	0 53	1 { III. 592. 593.
449								
450	} —	371 (1st $\tau$ ) Aqu	f	11	10	n	0 45	2 { Nebulosities run together.
451								
452	} —	18 Ceti	p	5	33	f	0 59	1 { tions. er. 2 or 3' from each other.
453								
453	} —	563 ( $\kappa$ ) Aqua	f	13	50	f	1 19	1 { F. pL. E. par. r.

II.	1785	Stars.	M.	S.	D.	M.	Ob.	Description.			
454	Oct. 5	90 ( $\phi$ ) Aqua	f	3	11	n	1	17	1	F. S. almost stellar.	
455	}	17 Eridani	f	11	19	n	0	26	2	{ Two. The p. pB. cL. E. lbM. The f. eF. vS. E.	
456			f	11	46	n	0	25			
457		61 ( $\omega$ )	p	4	31	f	0	2	2	F. cL. lbM.	
458		620	f	8	52	f	0	46	1	pB. R. bM.	
459			f	9	14	f	1	4	1	F. R. lbM.	
460			f	12	7	n	1	6	1	pB. S. lE. mbM. N.	
461		8 III ( $\xi$ ) Pisciu	p	28	48	f	1	32	3	F. pL. iR. vgbM. $1\frac{1}{2}$ dia.	
462			p	27	52	f	1	32	2	pB. R. vgbM. $1\frac{1}{4}$ dia.	
463			p	26	40	f	1	15	3	F. S. iE. par. mbM.	
464		44 Eridani	p	9	2	n	0	0	1	F. vS. r.	
465		9 82 ( $\delta$ ) Ceti	f	7	12	f	0	34	3	F. pL. iR. lbM.	
466			f	7	4	f	0	49	3	pB. cL. iR. mbM.	
467		25 7 ( $\beta$ ) Piscium	p	4	23	n	1	22	1	pB. pL. iF.	
468		26	f	0	11	f	1	10	1	F. pL. iF. r.	
469		26 49 Aquarii	f	5	14	f	0	4	1	F. pS. lE. er. some of the st. visible.	
470	Nov. 22	67 Ceti	f	37	51	f	3	27	2	pB. S. stellar.	
471		23 34 Piscium	f	20	53	f	0	55	1	F. iF. lbM.	
472		27 18 Ceti	f	2	18	n	1	24	1	F. pS.	
473		47	f	6	3	n	0	54	1	F. S. iF. er. some of the st. visible.	
474		72 ( $\rho$ )	p	9	28	n	0	56	2	pB. pL. lE. lbM.	
475		83 ( $\epsilon$ )	f	24	23	f	0	3	1	pF. pL. iF. bM.	
476		28 58 Aquarii	f	2	43	n	0	31	1	F. pL. iR. lbM.	
477		70	p	2	28	f	0	27	1	pB. pL. iR. lbM.	
478		17 Ceti	p	10	10	n	0	53	1	pB. L. lE. lbM.	
479			p	5	13	n	1	35	1	pB. mE. mer. 2' l.	
480			p	2	34	n	0	34	1	F. pL. lE. lbM.	
481		53 ( $\chi$ )	p	0	24	n	0	23	1	pB. cL. R. $1\frac{1}{2}$ f. Sst.	
482	}	55 (1st?)	f	17	54	n	0	15	1	{ Four. The p. 2, both F. E. S. within 1' dist. par. The f. two, both pF. pS. E. about 2' dist. and nearly mer.	
483			f	17	56	n	0	11	1		
484			f	20	13	n	1	5	1		F. S. E.
485			f	37	18	f	0	7	1		F. cL. iF. lbM.
486			f	49	13	f	0	50	1	F. S. iF. bM.	
487			f	8	36	n	0	42	1	F. S. lE. contains 3 ft. uncon.	
488			f	8	10	n	0	54	1	pF. mE. r. 3' l. $1\frac{1}{2}$ b.	
489	Dec. 7	66 (4 $\sigma$ ) Canc	f	8	10	n	0	54	1	pF. mE. r. 3' l. $1\frac{1}{2}$ b.	
490		18 Leo. min.	p	13	13	f	0	30	1	pB. pL. iF. lbM.	
491			f	1	47	n	0	0	1	pB. pL. lE. near. par.	
492		37	f	13	7	n	0	49	1	F. S.	
493		46 Ursa	p	3	47	f	0	36	1	pB. pL. iR.	
494		28 3 Leonis	f	3	34	n	0	16	1	F. pL. E. iF.	
495		9 ( $\circ$ ) Virgin	f	11	52	f	1	5	1	F.	
496		31 (ft d)	p	14	27	n	1	25	1	pF. vS.	
497											

II.	1785	Stars.	M.	S.	D.M.	Ob	Description.			
498	Dec. 28	31 (1st <i>d</i> ) Vir	p	12 30	n	1 3	1 F. pL.			
499	—	—	p	10 55	n	1 18	1 F.			
500	—	—	p	7 43	n	1 24	1 vL. er. some ft. visible.			
501	30	52 ( $\tau$ ) Ceti	f	4 36	n	1 1	1 F. S. R. vSpBN.			
502	—	76 ( $\sigma$ ) —	f	29 37	n	0 30	1 F. eS. stellar. p. pBft.			
503	—	—	f	31 37	f	0 15	1 pB. S. iF. mbM.			
504	—	20 Eridani	p	30 24	n	1 44	1 pB. S. lE. mbM.			
505	31	9 Hydræ	f	34 16	f	0 15	1 pB. S. lE. fp nf. fmbM.			
506	—	—	f	7 32	f	0 37	1 pB. S. lE. lb ffM.			
507	—	4 ( $\nu$ ) Crater	f	13 25	f	0 3	1 F. S. E.			
508	—	30 ( $\eta$ ) —	f	4 26	f	0 41	1 pB. S. lE. bM.			
509	—	—	f	6 52	n	0 46	1 F. cL. iR. lbM.			
510	—	53 Virginis	f	2 58	f	0 25	1 F. lE. $1\frac{1}{2}$ l.			
511	—	—	f	3 21	f	0 12	2 pB. pL. R. bM.			
512	—	—	f	3 55	f	0 12	2 F. S.			
513	—	—	f	4 53	f	0 27	2 pB. pL. iF. mbM.			
1786										
514	Jan. 1	49 Eridani	p	0 34	f	1 9	1 F. pL. E. fp nf. 2' l. 1' b.			
515	—	—	f	2 57	f	1 33	1 F. or pB. S. bM.			
516	—	—	f	21 45	f	1 16	1 F. S. iR. lbM.			
517	—	29 ( $\gamma$ ) Virgin	f	19 8	n	1 22	2 pB. pL. R. bM.			
518	}	2	13	Canum	p	44 34	n 2 49	2	}	Two. The p. F. S. E. The f. F.
519										
520	—	27	7 ( $\eta$ ) Hydræ	f	24 25	n	0 7	2 F. S. lE. par. er.		
521	—	—	77 ( $\sigma$ ) Leonis	p	3 42	f	1 28	3 F. vS. iF. fmbM. er.		
522	—	30	47 Eridani	f	6 29	f	0 21	1 F. pS. iE. r. 1' fp. Sft.		
523	—	—	—	f	10 15	f	0 17	1 F. vS. iR. bM. almost stellar.		
524	Feb. 1	57 ( $\mu$ ) —	p	9 24	n	0 3	1 F. S. iF. lbM. p. 2 Sft.			
525	—	—	p	4 5	n	1 27	1 F. pL. lE.			
526	—	—	f	0 16	n	0 51	1 F. cS. R. lbM.			
527	—	—	f	7 30	n	0 12	2 pB. S.			
528	—	—	f	7 40	n	0 12	1 F. S. lbM.			
529	—	28 (A) Hydr	p	26 37	n	0 8	1 F. S.			
530	—	260 ( $\sigma$ ) Virg	p	52 32	n	0 19	1 F. S.			
531	—	—	p	47 19	f	1 12	2 pB. pL. E. b. f. M. 3' l.			
532	—	—	p	35 12	f	1 28	2 F. pL. lbM.			
533	—	64 —	f	26 8	f	1 17	2 F. pL. vlbM. 6 or 7' l. 4' b.			
534	—	—	f	34 2	f	0 15	2 pB. vL. glbM.			
535	—	24	10 ( $\nu$ ) —	f	43 43	f	0 39	1 F. mE. np ff. 2' l. $\frac{3}{4}$ b.		
536	—	—	f	48 21	f	0 21	1 pB. mE. mbM. 2' $\frac{1}{2}$ l. 1' b.			
537	—	92 —	p	46 53	n	0 43	1 F. pL. iR. er.			
538	—	108 —	p	1 8	n	0 59	1 pB. cL. iR.			
539	—	110 —	p	2 58	f	0 11	1 pB. cL. lE. gbM.			
540	—	—	f	1 11	f	0 53	1 pB. S. mbM.			
541	—	—	f	2 31	f	0 28	1 F.			

II.	1786	Stars.	M.	S.	D.M.	Ob	Description.							
542	Feb. 24	110 Virg	f	2 31	n 0 0	1	pB.							
543	—	—	f	4 14	f 0 34	1	F.							
544	—	—	f	4 52	n 0 27	2	pB. vS.							
545	—	—	f	6 51	f 1 39	4	pB. S. iE. lbM.							
546	} Mar. 3	6 (b) Leonis	p	6 16	n L 42	1	{ Two. Both F. S. The place in- accurate in RA.							
547		—	p	10 27	f 0 8	1		F. pL. mE. np ff. but near. par.						
548		14 Virginis	f	17 34	f 0 33	1		pB. vL. iF. lbM.						
549	—	26 (x) —	f	4 13	n 0 35	2	{ Two. Both F. S. lbM. cBft. be- tween, but 1 1/2 f. of them.							
550	} 4	14 (e) Crate	p	4 0	n 0 36	2								
551		—	21 (θ) —	p	2 24	f 0 2	1	F. pS. iR. f. vSft.						
552	—	—	f	11 21	f 1 9	2	pB. pL. iF. gbM. fp. is Sft.							
553	18	1 Cancri	f	4 36	f 0 4	2	pB. pL. er. vgmbM.							
554	19	26 Hydræ	f	7 26	n 0 21	2	pB. pL. iR. b. f. M.							
555	20	6 (3d b) Crat	p	76 10	f 1 11	3	pB. cL. iR. vgmbM.							
556	24	16 (ζ) Hydræ	f	3 21	n 0 22	1	F. mE. unequally B. 3'l. 1'b.							
557	25	21 (q) Virgin	f	10 43	f 0 38	1	F. E. mer. 3'l. f. cBft.							
558	—	49 (g) —	p	14 43	n 1 33	1	F. S.							
559	—	—	p	13 0	n 0 31	1	pF. pS. iR.							
560	—	—	p	3 39	n 0 24	1	pB. pL. R. vgmbM.							
561	27	16 (x) Crater	f	4 56	f 1 54	2	F. S. iR. bM. r.							
562	—	68 (i) Virgin	p	29 28	f 0 55	1	pB. iF. bM.							
563	28	19 Ursæ	p	3 1	n 0 23	1	pB. S. R. mbM.							
564	—	46 Leo. min.	p	5 3	n 0 28	1	pB. cL. iF. lbM.							
565	—	* 1102 (e) Hy	f	35 28	n 0 53	1	F. pS. E. * See note.							
566	—	—	f	37 17	n 0 51	1	pB. pL. iF. gbM.							
567	} Apr. 17	11 (s) Virgin	f	10 14	n 0 34	1	{ Four nebulae. They are scat- tered about. The place is that of the last.							
568								—	f	11 34	f 0 26	1	A nebula.	
569								2	—	f	10 18	f 0 26	1	A nebula, cloudy.
570								2	3 Serpentis	p	40 48	i 0 20	1	F. S. iE. r. p. 2 vcf.
571								—	—	p	36 3	n 0 33	1	pB. cL. iR. mbM.
572								—	—	p	21 26	f 0 54	1	F. S. iE. like 2 stellar: joined closely.
573								30	37 Virginis	p	11 22	n 0 4	1	F. S. making a triangle with 2 Bft.
574								—	—	p	2 29	n 0 20	1	F. S.
575								—	109 —	p	26 11	n 2 10	1	pB. cL. E.
576								—	—	p	16 35	n 1 24	1	{ Two. The f. pB. pL. R. gbM. The n.eF. cL. dist. 2'. The place is of 1.
577	—	—	p	8 33	n 0 25	1	F. mE. r. 2' 1. 1/4 b. f. f. 6m. 16'' in time.							
578	May 3	14 (1st A) Ser	f	17 48	f 1 2	2	pB. S. E. nearly par. bM.							
579	20	5 (g) Ophiuc	f	27 48	n 1 8	1	pB. cL. gbM. er. undoubtedly ft.							
580	27	3 Serpentis	p	5 43	f 1 52	1	F. S. iE. r.							
581	28	40 (e) Ophiuc	f	28 13	n 0 57	1	pB. S. iF.							

II.	1786	Stars		M. S.		D.M.	Ob	Description.
587	June 3	61 Ophiuchi	f	0 23	n	0 36	1	F. cL. iF.
588	Sept. 4	24 ( $\xi$ ) Ariet	p	39 40	f	0 17	2	F. S. iE. r. bM.
589	—	—	p	36 21	n	0 50	2	F. pL. E. b. f. M. 2' fp. cBft.
590	18	2 Piscium	f	2 2	n	0 48	1	F. S. bM.
591	—	88 ( $\gamma$ ) Pegasi	p	4 29	n	0 38	1	F. pL. iF. unequally B.
592	—	85 Ceti	p	3 19	n	0 5	1	pB. S. E. bM.
593	20	54 Eridani	p	61 14	n	0 43	1	pB. pS. R. resembling I. 107. but less.
594	—	—	p	55 40	n	0 10	1	pB. vS. R. bM.
595	23	66 Aquarii	p	41 2	f	0 1	2	F. cL. l and iE. nearly par. lbM.
596	30	51 Ceti	f	10 14	n	0 51	1	F. S. bM. 1' f Sft.
597	—	32 Eridani	p	8 30	f	1 10	2	F. S. E. iF. in a row with some ft.
598	Oct. 13	59 ( $\nu$ ) Aqua	f	13 11	f	1 39	1	pB. pL. iR, vgmbM.
599	17	77 Cygni	f	20 15	f	0 6	1	F. pS. E. er.
600	—	10 Androme	f	2 5	f	1 14	2	{ pB. mE. np ff. but near mer. lbM. r. 5' l. 1' $\frac{1}{2}$ b. also ob. 1784.
601	—	26 ( $\beta$ ) Persei	p	15 16	n	1 14	1	F. S. iF. r.
602	—	—	p	13 38	n	0 34	1	F. pS. iR. lbM.
603	—	—	f	11 27	n	0 35	1	pB. stellar. or pcft. with S. vF. chev.
604	18	59 Androme	p	2 10	f	0 17	1	pB. cL. iE. mbM.
605	—	—	p	0 54	n	0 9	1	pB. S. iF.
606	24	6 Lacertæ	p	17 44	n	2 18	3	F. S. er. or rather a patch of ft.
607	—	30 Persei	p	12 50	f	1 44	1	F. cL. E.
608	—	—	p	11 45	n	0 19	1	F. cL. er. some ft. visible.
609	26	65 ( $\iota$ ) Piscium	p	1 55	f	0 6	1	pB. S. iR. gbM.
610	—	90 ( $\nu$ ) —	f	24 26	n	1 31	1	F. S. bM. r.
611	—	—	f	27 38	n	0 41	1	F. S. iE.
612	—	10 ( $\alpha$ ) Triang	p	28 30	f	1 8	1	pB. pL. iE. nearly par. mbM.
613	—	—	p	4 46	f	0 47	1	F. S. iE. par. bM.
614	}	— 34 ( $\theta$ ) Gemin	p	5 37	f	0 25	1	{ Two. The f. F. S. R. bM. The n. F. cS. R. bM.
615								
616								
617	Nov. 13	6 ( $\beta$ ) Arietis	p	3 55	n	0 56	1	F. cL. vglbM.
618	—	—	p	3 23	n	1 45	1	vS. stellar.
619	—	52 —	p	5 39	f	0 3	1	pB. cL. pmE. mer. r. 1' f. ft.
620	Dec. 11	27 ( $\kappa$ ) Persei	p	5 48	n	1 31	2	F. S. iR. bM. L. stellar.
621	13	34 Ceti	p	23 45	f	0 34	1	F. E. np ff. lbM. 1' $\frac{1}{2}$ l.
622	20	26 —	f	9 8	f	0 22	1	F. R. bM. er.
623	21	2 ( $\epsilon$ ) Corvi	p	16 4	f	0 33	2	F. S. E. mer. or few deg. np ff. lb. f. M.
624	29	1 Sextantis	f	8 54	f	1 8	1	F. iE. nearly par. 1' $\frac{1}{2}$ l.
625	—	29 ( $\gamma$ ) Virgin	p	17 56	f	1 58	2	pB. mE. 20° fp nf. 2' l.
626	30	77 ( $\sigma$ ) Leonis	p	4 44	f	1 30	1	pB. S. iE. mbM.
1787								
627	Jan. 11	55 ( $\delta$ ) Gemi	f	54 51	f	0 26	3	F. S. iF. iE. fp nf.
628	14	6 Comæ	f	6 36	n	0 38	1	pB. cL. E.
629	—	—	f	13 46	f	0 49	1	F.



H.	1787	Stars.	M.	S.	D.M.	Ob	Description.
630	Jan. 14	6 Comæ	f	13 20	f	0 56	1 cL.
631	—	—	f	16 3	f	1 31	1 F.
632	—	29 —	p	8 57	n	1 12	1 F. pL. R. vgbM.
633	17	16 (1 ft p) Perſ	p	7 2	f	1 1	1 F. cL. lbM. 4' dia.
634	Feb. 13	33 (n) Cancr	p	12 7	n	0 34	1 F. S. bM.
635	22	21 (θ) Crater	p	13 5	n	1 9	1 F. pS. iR. vgbM.
636	—	65 Virgin	p	43 8	f	0 49	1 F. vL. bM.
637	March 11	44 (k) —	f	12 41	f	0 36	1 F. cL. iR. lbM. time inaccurate.
638	15	* 1 139(r) Ce	f	22 49	f	0 12	1 pB. S. 1E. fp nf. * See note.
639	17	32 Leo. min.	p	16 31	f	0 11	1 pB. cS. r.
640	—	—	p	16 11	f	0 18	1 F. vS. r. with 300 the fame.
641	—	38 —	f	2 41	f	0 36	2 F. vS.
642	—	6 Canum	p	15 18	n	0 30	2 pB. S. E.
643	—	10 —	p	0 37	f	2 11	1 F. pL. gbM. r.
644	—	—	f	2 55	f	1 1	1 pB. S. R. mbM. among scattered ft.
645	—	—	f	4 33	f	1 2	1 pB. S. R. mbM.
646	—	17 —	f	12 21	n	0 12	1 pB. L. iF. uneq. B. 3. or 4' dia.
647	—	12 (λ) Coronæ	f	33 4	n	1 27	1 F. S. iF.
648	18	53 (2d v) Boot	p	55 31	n	1 11	2 pB. pL. lbM.
649	—	—	p	54 11	f	0 13	2 F. S. E. nearly mer: r.
650	—	—	p	16 19	n	1 13	3 pB. E. BNM. and F. br. 2' l. $\frac{1}{4}$ b.
651	—	—	p	5 42	n	0 51	2 pB. pL. iE. er.
652	—	30 (g) Hercul	p	0 57	f	0 57	1 F. pL. r.
653	19	70 Virginis	p	4 21	n	0 11	1 pB. vS. mbM. juſt p. pcft.
654	—	9 Serpentis	f	7 56	f	0 28	1 F. E. np ff. $1\frac{1}{2}$ l.
655	—	—	f	15 44	n	0 16	1 F. E. mer. $1\frac{1}{2}$ l.
656	—	—	f	16 59	f	1 17	1 pB. E. np ff. bM. $1\frac{1}{2}$ l.
657	—	28 (β) —	f	8 2	f	0 52	1 F. iF. bM. $1\frac{1}{4}$ dia. between 2Bft.
658	20	44 Lyncis	p	47 39	f	0 23	1 pF. vS. mbM.
659	—	13 Canum	p	18 44	n	1 47	1 F. S. R. juſt np. V. 42.
660	April 9	8 —	f	7 58	f	0 5	1 pB. pL. R. mbM.
661	—	—	f	9 42	f	0 20	1 pB. vS. ſtellar. juſt p. Sft.
662	—	—	f	15 2	n	0 36	1 F. S. R. bM.
663	—	19 —	p	9 58	n	0 56	1 pB. vS. ſtellar. near and n. Sft.
664	—	—	p	3 47	n	3 13	2 pB. mE. ip nf. near. mer. 5' l. $\frac{3}{4}$ b.
665	—	20 —	f	2 52	n	2 31	1 pB. cS. E. with 300 ft. with burrs.
666	—	—	f	5 24	n	2 30	1 pB. S. iR. mbM.
667	—	—	f	7 35	n	2 42	1 pB. vS. 1E. bM.
668	—	—	f	27 51	n	0 51	1 F. E. par. miniature of I. 170.
669	—	—	f	33 20	n	0 41	1 pB. pL. vgbM.
670	—	—	f	35 10	n	2 37	1 pB. pL.
671	—	—	f	37 51	n	0 35	1 pB. pL. E.
672	—	—	f	43 59	n	0 17	1 pF. pS. bM.
673	—	—	f	66 36	n	1 0	1 F. pL. E. vlbM.
674	—	—	f	71 16	n	0 27	1 pB. E. nearly par. $1\frac{1}{2}$ l. $\frac{1}{2}$ b.

II.	1787	Stars.		M.	S.		D.M.	Ob	Description.	
675	April 9	20 Canum	i	80	7	n	0	51	1	F. vS.
676	—	—	f	98	12	n	1	42	1	pB. vS. stellar.
677	—	—	f	99	9	n	1	39	1	F. pS. lbM.
678	—	—	f	117	42	n	1	1	1	F. S. r. in a row with 3 ft.
679	} 11	79 (ζ) Virgin	p	4	17	f	1	1	2	{ Two. The p. F. pS. iF. The f. pB. pL. iF. bM.
680				4	7	f	1	4	2	
681	—	1 Serpentis	p	19	44	f	0	7	2	pB. pL. iF.
682	—	—	p	16	35	f	0	4	2	pB. cS. iE.
683	—	—	f	0	49	f	0	55	1	pB. pL. R. mbM. ff. cft.
684	—	4 —	p	6	6	n	0	7	1	{ Two. The 2d pB. S. iE. for the 1st see II. 545.
685	15	90 (ρ) Virgin	p	2	37	f	0	44	2	F. pL. iR. f. and par. with 2Fft.
686	—	—	p	0	37	n	0	4	2	pB. S. mbM.
687	—	102 (1 ft v) —	p	6	18	f	0	57	2	pB. cL. mE. 20° sp nf.
688	May 11	19 (λ) Bootis	p	30	37	n	0	7	2	F. mE. 15° sp nf. lbM. 4'1. $\frac{3}{4}$ b.
689	12	—	p	47	20	n	0	46	3	pB. pL. R. mbM.
690	—	22 (τ) Hercu	f	7	2	n	2	3	2	F. pL. iF. gbM.
691	15	85 (η) Urfæ	f	15	34	f	0	12	1	pB. pL. E. nearly par. mbM.
692	}	—	f	19	36	n	1	20	1	{ Two. The p. F. pS. R. vgbM. The f. F. vS. stellar. fmbM. dist. 2' $\frac{1}{2}$ .
693										
694	—	24 (g) Bootis	p	6	31	n	0	43	1	pF. pS. iE. mbM.
695	—	—	f	1	7	f	0	12	1	pB. cL. iR. vgmbM.
696	—	—	f	3	40	n	0	3	1	pB. S. E.
697	16	*C Canu. 6m	f	6	23	f	0	39	1	F. E. par. bM. 1' $\frac{1}{2}$ l. 1' b. *See note.
698	—	—	f	10	0	f	0	58	1	F. S. R. vsmBM.
699	—	—	f	13	19	n	0	20	3	F. pL. R. lbM. 1' $\frac{1}{2}$ dia.
700	—	27 (γ) Bootis	f	5	15	n	0	9	1	pF. S. iE.
701	—	25 Herculis	f	17	43	f	0	40	1	pB. pS. E. sp nf. vgmbM.
702	Sept. 11	68 (2dg) Aqu	f	4	23	f	1	1	1	{ pF. pL. E. np ff. but near. par. mbM. 1' $\frac{1}{2}$ l.
703	—	*A Ceti 7m	f	4	47	n	1	7	1	F. cL. E. * See note.
704	16	47 Cassiop	f	61	37	n	3	48	2	F. pL. mE. np ff. mbM.
705	Nov. 3	25 Cephei	f	21	6	f	1	35	1	pB. S. iR. er. almost equally B.
706	—	1 (e) Cassiop	f	6	26	n	2	5	2	{ pBM. 2cft. involved in nebulo- sity. 2' l. 1' $\frac{1}{2}$ b.
707	30	19 (ξ) —	p	2	50	f	2	12	1	pB. vL. iR. vgmbM. r. 5 or 6 dia.
708	1788 Jan. 14	37 Lyncis	f	3	50	f	1	15	1	pB. S. stellar.
709	—	56 Urfæ	p	6	51	f	1	57	2	pB. S. iE. mer. bM.
710	—	27 (γ) Bootis	p	43	42	n	1	51	1	F. S.
711	—	—	p	42	47	n	1	50	1	pB. cL. iF.
712	—	—	p	41	48	n	1	25	1	F. S. R. bM.
713	—	—	p	39	13	n	2	12	2	pB. pL.
714	}	—	p	39	5	n	2	9	2	{ Two. Both pB. S. R. 2' dist. in the same mer.
715										

II.	1788	Stars.		M.	S.		D.M.	Ob	Description.
716	Jan. 14	27 ( $\gamma$ ) Bootis	p	36	48	n	2 19	2	pB. L. iR. FN. mbM. 4 or 5' dia.
717	15	15 Leo. min.	f	0	58	f	1 58	1	F. pL. iF. lbM.
718	—	45 ( $\omega$ ) Urfæ	p	2	24	n	0 32	2	{ pB. S. IE. the np. corner of a S. trapezium.
719	Feb. 3	32 Lyncis	p	20	34	f	0 16	1	F. pL. iR. bM.
720	—	34 ( $\mu$ ) Urfæ	p	2	13	n	1 29	1	F. vS.
721	—	— — —	p	1	57	n	1 26	1	F. vS. stellar.
722	—	— — —	p	1	43	n	1 27	1	F. vS. stellar.
723	—	13 Canum	p	73	0	f	0 22	1	pB. S. IE.
724	—	— — —	p	65	22	f	0 44	1	F. vS.
725	—	— — —	p	61	59	f	0 19	1	pB. E. spnf. but nearer mer. mbM. 2' 1
726	5	80 ( $\pi$ ) Gemi	f	22	56	n	0 38	1	pF. pL. iR. lbM. r. f. 2 ft. par.
727	—	59 (2 $\sigma$ ) Canc	p	13	13	n	1 47	1	pF. pL. iR. r.
728	—	60 Urfæ	p	25	2	n	1 32	2	pB. pL. vgmbM.
729	—	— — —	p	20	38	f	1 4	2	F. cL. IE. par. lbM.
730	—	— — —	p	5	27	n	0 14	1	pB. bM. r. 4' l. 3' b.
731	—	— — —	p	0	54	n	1 5	2	pB. S. E. sp nf.
732	—	— — —	f	0	47	f	0 19	1	{ F. S. almost. betw. 2 sp. ft. chev. touches them.
733	6	59 — — —	f	20	28	n	0 21	1	pB. mE. mer. pBSN. & vF. br. 4' l. $\frac{3}{4}$ ' b.
734	9	20 Lyncis	p	14	20	n	0 28	1	F. pL. iF. mbM. ff. a triangle of S. ft.
735	—	— — —	p	12	44	f	1 37	1	F. stellar.
736	—	— — —	p	11	9	f	0 3	1	pF. vS. lbM. r.
737	—	63 ( $\chi$ ) Urfæ	p	4	55	f	0 4	2	pF. pS. iR. lbM.
738	—	— — —	f	2	30	n	0 57	1	pB. pL. R. mbM.
739	—	— — —	f	2	50	n	0 56	1	F. vS.
740	—	— — —	f	5	48	n	0 56	1	pF. pS. stellar.
741	—	— — —	f	17	7	n	0 53	1	pF. S. R. gbM.
742	—	3 Canum	p	1	50	f	1 35	2	F. S. E.
743	—	— — —	f	5	22	f	0 10	1	F. S.
744	—	— — —	f	21	50	n	1 27	2	pF. S. er.
745	April 1	Neb. II. 728.	p	35	22	f	0 57	2	pF. pS. E. f. & lp. ft. among ft. not con.
746	8	54 Virginis	p	0	24	f	0 43	1	pB. S. pBN.
747	10	60 Urfæ	f	31	58	f	0 24	1	pB. E. 15 or 20° np ff. 3' l.
748	—	— — —	f	38	3	n	0 16	1	pB. pL. E. sp. and in a line with 2 ft.
749	—	— — —	f	47	57	f	1 10	2	pB. pL. iF.
750	27	19 ( $\lambda$ ) Bootis	p	109	46	f	1 1	1	pF. pL. E. sp nf.
751	—	37 ( $\xi$ ) — — —	f	16	12	n	0 25	2	{ Two. The p. cF. cS. The f. pF. pL.
752	—	— — —	f	16	20	n	0 24	2	{ Both IE. np ff. but nearer par.
753	28	27 ( $\beta$ ) Hercul	f	2	50	f	1 42	1	pF. pS. vIE. mbM.
754	29	27 ( $\gamma$ ) Bootis	p	11	15	n	1 27	1	pB. pL. R. FN.
755	May 1	23 ( $\theta$ ) — — —	f	44	59	n	0 31	1	pB. pL. IE.
756	5	Neb. II. 757.	p	11	47	f	3 7	2	pB. pL. iF. r.
757	—	12 ( $\iota$ ) Draco	p	16	38	f	1 56	3	pB. S. iR. or IE. mbM.
758	—	Neb. II. 757.	f	5	28	f	1 31	1	pF. pS. iR.

II.	1788	Stars.		M.	S.		D.M.	Ob	Description.
759	May 5	Neb. II. 757.	f	6	6	f	0 42	1	pB. FNM. 8 or 10' l. 2' b.
760	—	—	f	6	29	f	1 37	1	pF. pS. R.
761	—	—	f	24	8	f	0 33	1	pF. pS. iF.
762	—	—	f	24	37	f	0 25	1	pF. pL. E.
763	25	12 (i) Draco	p	13	7	n	0 54	2	pB. mE. nearly mer. 2' l. $\frac{1}{2}$ ' b.
764	—	—	f	13	58	n	0 20	1	pB. S. iR. one p. suspected vF. lE.
765	—	—	f	14	36	f	0 58	1	pF. cS.
766	—	—	f	15	0	n	0 18	1	pB. cL. iE. r.
767	June 6	31 (1st $\Psi$ )	p	31	23	n	0 15	1	pB. pL. R. vgmbM.
768	Nov. 4	14 Camelop	p	42	52	n	1 57	1	pB. S. lE. BN. just f. pB. ft.

## Third class. Very faint nebulae.

III.	1785	Stars.		M.	S.		D.M.	Ob	Description.
377	April 26	92 Leonis	f	3	6	f	1 24	2	{ Two. The n.F.S.lBM. The f.vF. vS. dist. 5' sp. the place of n.
378									
379	—	1 Comæ	p	10	26	f	0 6	3	vF. vS. lE. er. or S. patch of ft.
380	—	—	p	8	56	f	1 7	2	F. S.
381	—	—	p	7	56	f	1 12	1	vF. R.
382	—	2 —	f	1	58	f	0 49	2	{ Three. The place is of the last or most n. which is vF. S. The other two are sp. eF. vS.
383									
384									
385	27	93 Leonis	p	2	54	f	0 35	1	vF. vS. r.
386	—	—	p	3	10	f	0 27	1	vF. vS. r.
387	—	—	p	2	10	f	0 27	1	vF. vS. r.
388	—	—	p	1	48	n	0 11	2	vF. pL. iR. lBM. r. 7' nf. cBft.
389	—	—	f	4	49	n	0 25	1	vF. vS.
390	—	5 Comæ	p	8	56	f	1 47	1	Suspected.
391	—	—	p	7	54	f	0 9	2	{ Six nebulae. The places belong to the three first which are vF. vS. The other three are 10 or 12' more south, but there was not time to take their places. more suspected.
392									
393									
394									
395									
396									
397	—	—	f	3	36	n	0 5	2	vF. vL. iR. bM. 6' l. 5' b.
398	—	26 —	f	8	12	f	1 33	1	vF. vS. r.
399	28	61 Urfæ	f	31	26	n	1 57	2	vF. pL. lE. r.
400	May 1	—	f	25	34	n	2 38	4	vF. vS. stellar. 2' $\frac{1}{2}$ n. Sft.
401	—	14 Canum	f	2	36	f	0 36	1	vF. stellar. with 300 the same.
402	—	—	f	19	23	n	0 35	1	{ Two. Both vF. cS. The place is that of the p. The 2d, 3'nf.
403									
404									
405	—	—	f	20	44	n	0 7	1	{ Two. Both vF. pS. The place is of the p. The 2d, 5 or 6'nf.
406									
406	—	—	f	25	14	f	0 59	1	vF. vS. lE.

III.	1785	Stars.	M.	S.	D.M.	Ob	Description.	
407	} May 1	49 ( $\delta$ ) Bootis	P	102	40	n	I 37	} Two. Both vF. vS. A star between them about half way.
408				102	22	n	I 39	
409	—	14 Canum	f	30	38	f	O 17	vF. pL. R. lbM.
410	—	—	f	35	6	n	I 1	vF. S. lE. er.
411	—	—	f	54	30	f	I 8	eF. vS.
412	—	—	f	54	45	n	O 25	vF. vS.
413	—	—	f	58	30	f	O 53	vF.
414	—	51 ( $\mu$ ) Bootis	p	70	6	f	O 55	vF. mE.
415	—	—	p	65	4	f	I 57	eF. pL.
416	} —	—	p	64	2	f	I 57	} Two. Both vF. S. dist. 6 or 7'. The place is that of the ff.
417								
418	—	—	p	62	52	f	O 6	eF. stellar.
419	—	—	p	61	4	f	O 42	vF. vS. E. er.
420	—	—	p	54	54	f	O 50	vF. S.
421	—	—	p	49	26	f	O 40	vF. vS.
422	} 2	49 ( $\delta$ ) —	p	86	2	n	O 24	} Two. Both eF. stellar. dist. 4 or 5'. nearly mer. The n. faintest.
423								
424	3	—	p	147	32	n	O 11	vF. stellar. or little larger.
425	—	—	p	101	48	n	I 39	vF. vS. in the field with III. 407. 408.
426	Aug. 30	17 ( $\iota$ ) Piscium	p	8	48	f	I 42	eF. pL. iR.
427	—	19 —	f	0	14	n	O 19	vF. S. lE. nearly mer.
428	Sept. 10	30 —	f	14	30	f	O 19	vF. S. iF. lbM.
429	—	41 Ceti	p	26	42	n	O 35	vF. pS. E.
430	—	—	p	26	54	n	O 44	vF. vS.
431	—	Neb. I. 100.	f	0	22	n	O 0	} The 2d of two. eF. S. 5 or 6' dist. from I. 100.
432	—	41 Ceti	f	15	36	n	O 22	eF.
433	—	67 —	p	15	39	n	O 59	vF. vS.
434	—	—	f	18	40	f	O 42	vF. cL. iF. lbM. 4 or 5' l. 2 or 3' b.
435	26	59 ( $p$ ) Pegasi	f	8	42	f	O 20	vF. vS.
436	—	32 (2dc) Pifc	f	1	20	f	I 1	vF. pL. lbM.
437	27	26 —	p	7	39	f	O 13	eF. vS. er. confirmed by 240.
438	28	93 (2 $\Psi$ ) Aqua	f	9	22	f	O 15	eF. S. stellar. p. $1\frac{1}{2}$ . pBft.
439	Oct. 1	20 Ceti	p	0	42	f	I 3	vF. S. iE.
440	—	38 —	f	1	5	n	O 8	vF. vL. requires great attention.
441	—	43 —	f	5	8	f	I 29	vF. vS. iE.
442	—	—	f	5	23	f	I 26	vF. vS. iE.
443	5	17 Eridani	p	17	51	f	O 13	vF. vS. confirmed by 240.
444	—	—	p	9	23	n	O 25	eF. vS.
445	—	—	p	5	37	f	O 41	vF. pS. E.
446	—	—	f	3	4	f	O 3	vF. S. between some Sft.
447	—	20 ( $\tau$ ) Orion	f	10	23	n	I 32	vF. cL. iR. near a hook of vSft.
448	—	—	f	34	45	f	O 26	vF. S. R. r. lbM.
449	6	1 (1st $\tau$ ) Erid	f	4	8	n	I 34	vF. pL. broadly E. lbM.
450	—	—	f	6	30	n	I 56	vF. S. lE.

III.	1785	Stars.	M.	S.	D.M.	Ob	Description.
451	Oct. 6	20 Eridani	f	2 30	f 0 59	I	vF. S. R.
452	8	52 ( $\pi$ ) Aqua	p	30 46	n I 39	I	vF. pL. R. r.
453	—	10 Orionis	f	5 7	f 0 4	I	vF. vS. confirmed 240.
454	9	60 Ceti	p	27 18	n 0 27	I	eF. pL. 240. left doubtful.
455	—	82 ( $\delta$ ) —	f	4 11	n I 2	2	vF. vL. lbM. er. 6 or 7' dia.
456	25	28 ( $\omega$ ) Piscium	f	13 6	f 0 28	I	vF. pS. iF.
457	—	78 ( $\nu$ ) Ceti	p	20 29	n 0 20	I	vF. cL. vlbM. m. p. Bft. and joining.
458	26	49 Aquarii	p	2 52	n 0 6	I	vF. S. er. time inaccurate.
459	—	56 (1st $\nu$ ) Ceti	p	7 44	f I 17	I	vF. vS. er.
460	—	— — —	p	2 55	f I 16	I	vF. vS.
461	27	18 ( $\epsilon$ ) Pis. Au.	f	9 20	n I 56	I	vF. cL. lE. glbM. 4 or 5' l.
462	Nov. 7	82 ( $\delta$ ) Ceti	f	8 1	f 0 36	I	vF. S.
463	22	25 —	p	12 56	f 0 23	2	vF. pL. iR. r.
464	—	67 —	p	20 11	n 0 59	I	eF. S. found in gaging.
465	23	46 ( $\xi$ ) Pegasi	f	11 21	n 0 54	I	eF. S. iF. 240 the same.
466	—	82 —	f	5 54	f 0 15	I	vF. S. R. lbM.
467	27	18 Ceti	p	11 15	n 0 12	I	eF. vS. 240 left some doubt.
468	—	72 ( $\epsilon$ ) —	p	27 13	n 0 43	I	vF. E. nearly mer. lbM. 1' $\frac{1}{2}$ l. 1' b.
469	—	83 ( $\epsilon$ ) —	f	19 27	f 0 30	I	vF. stellar. 240 left some doubt.
470	28	91 Aquarii	p	1 53	f 0 7	I	eF. vS. 240 left doubtful.
471	—	53 ( $\chi$ ) Ceti	p	13 54	n 0 40	I	A few Sft. mixed with nebosity.
472	—	55 (1st $\zeta$ ) —	f	41 48	f 0 18	I	vF. pL. vlbM. near scattered ft.
473	29	87 ( $u$ ) Pegasi	p	44 53	f I 26	I	eF. cL. some doubt. p. a row of ft.
474	—	23 (2d $\theta$ ) Arie	f	7 29	n 0 50	I	eF. vS. iR. confir. 240.
475	—	34 ( $\mu$ ) —	p	1 44	f 0 44	I	vF. S. confir. 240.
476	Dec. 5	34 ( $\zeta$ ) Andro	p	11 14	f 0 23	I	vF. vS. stellar. sp. pBft.
477	—	36 —	p	2 25	n 0 44	I	vF. S. R. just p. vFft.
478	7	20 Leo. min.	f	1 20	n 0 47	I	eF. S. left doubtful.
479	26	2 ( $\epsilon$ ) Can. min	f	26 18	n 0 25	I	suspected. eF. vS. lE.
480	28	9 ( $\sigma$ ) Virgin	f	12 46	f 2 5	I	vF. L. seen by looking at II. 137.
481	—	31 (1st $d$ ) —	p	17 49	n I 44	I	vF.
482	—	— — —	p	15 22	n I 39	I	eF.
483	—	— — —	p	12 49	n I 24	I	vF.
484	—	— — —	f	11 8	n I 34	I	vF.
485	30	46 Ceti	p	40 9	f I 4	2	vF. S. iF. r.
486	—	76 ( $\sigma$ ) —	p	12 32	f 0 52	I	vF. vS. iF. better with 240.
487	—	20 Eridani	p	3 52	n 2 14	I	vF. S. E.
488	31	9 Hydræ	f	38 13	f 0 26	I	vF. cL. glbM. 3' l. 2' b. p. pBft.
489	—	53 Virginis	p	18 36	f 0 47	I	vF. S. lbM.
490	1786 Jan. 1	45 Eridani	p	11 41	f 0 42	I	vF. vS. lE. better with 240.
491	—	13 ( $n$ ) Virgin	p	16 10	n 0 35	2	vF. S. R. bM.
492	—	15 ( $n$ ) —	f	7 0	f 0 15	2	vF. cL. mE. r.
493	—	29 ( $\gamma$ ) —	p	6 35	n I 12	2	eF. S. iF.
494	—	— — —	f	1 24	n 0 48	2	vF. pS. E.

III.	1786	Stars.	M.	S.	D.M.	Ob	Description.
495	Jan. 2	61 Urfæ	f	58 0	f 0 46	1	eF. S. iF. r.
496	—	—	f	70 52	f 0 3	1	eF. vS. pmE.
497	27	36 Sextantis	f	6 47	n 1 20	2	cF. S. R. vlbM.
498	—	58 (d) Leonis	f	0 43	n 0 1	1	vF. mE.
499	30	39 (A) Erida	p	6 26	n 1 25	1	vF. S. E. er.
500	—	69 (λ) —	p	3 50	f 0 24	1	cF. S. iF. bM.
501	Feb. 1	57 (μ) —	f	4 13	n 0 30	1	vF. vS.
502	—	—	f	6 2	n 0 39	1	vF. S.
503	—	—	f	14 49	f 0 1	1	vF. pL. sp. 2pBft. equil. triang.
504	2	60 (σ) Virg	p	38 27	n 0 34	2	vF. pS.
505	—	64 —	f	16 1	f 0 39	2	vF. vS. R.
506	—	—	f	32 47	n 0 7	1	vF. E. 2' l.
507	4	82 —	p	9 23	f 0 4	1	vF. vS. er. 240 rather confir.
508	—	19 Libræ	p	18 52	f 0 27	1	vF. cL. iE. nearly mer.
509	22	5 (β) Virgin	f	49 54	f 0 35	1	vF. vS.
510	24	55 Orionis	f	1 13	n 0 7	1	eF. E. er. probably a patch of ft.
511	—	110 Virginis	f	3 5	f 0 25	1	vF. R. precedes I. 128.7' $\frac{1}{2}$ . and is 5' n
512	March 3	17 (β) Cancr	p	14 9	n 0 9	1	vF. S. R. mbM. 240 ditto.
513	—	6 (h) Leonis	f	2 1	n 0 25	1	eF. vS. stellar. 240 verif.
514	—	26 (χ) Virgin	f	10 4	f 1 8	2	eF. S. mE.
515	—	—	f	12 19	f 0 26	1	vF. S. E.
516	—	—	f	14 18	f 0 41	1	vF. S.
517	—	—	f	14 43	f 0 48	1	vF. S.
518	19	41 (λ) Hydræ	p	0 28	i 0 5	1	vF. S. R. in the field with λ
519	24	1 Sextantis	f	1 47	n 0 7	1	{ vF. pL. vgvlbM. betw. 2 groups of ft. np. ff.
520	25	27 Hydræ	f	3 9	f 0 51	1	vF. S. E.
521	—	—	f	22 39	f 0 45	1	cF. pS. lE.
522	—	14 (ε) Crater	p	34 1	f 2 2	2	cF. pL. iR. lb. near M.
523	—	21 (q) Virgin	f	13 23	f 0 38	1	vF. E. sp nf. 4' l. 3' b.
524	—	—	f	15 14	f 1 59	2	cF. mE. r. 4' l. $\frac{3}{4}$ ' b.
525	—	49 (g) —	p	14 19	n 1 14	1	vF. vS.
526	—	—	p	13 15	n 0 8	1	eF. eS. some little doubt.
527	27	8 Sextantis	p	10 33	f 0 31	3	vF. S. iR. vgbM.
528	—	—	p	9 10	f 1 32	1	vF. S. E. nearly mer.
529	—	16 (κ) Crater	p	13 0	f 1 46	1	eF. S.
530	—	—	p	3 32	f 1 30	1	vF. stellar.
531	—	—	p	2 47	f 1 32	1	cF. stellar. vlbM.
532	—	—	f	1 7	f 0 51	1	vF. lE. vlb. about M.
533	—	24 (ι) —	f	28 31	f 0 59	1	vF. S. iF. time a little inacc.
534	—	—	f	33 51	f 0 48	1	vF. pL. of uneq. light.
535	—	—	f	40 50	n 0 58	2	vF. pS. iF.
536	—	68 (ι) Virgin	p	36 17	f 0 33	1	cF. stellar.
537	—	—	p	34 23	f 0 39	1	vF. vS. iF.
538	—	—	p	31 24	n 0 8	2	eF. S. er.

III.	1786	Stars.		M.	S.		D.M.	Ob	Description.
539	Mar. 27	68 ( $\iota$ ) Virgin	p	5	57	f	I 21	I	vF. vS.
540	28	19 Ursæ	p	15	27	n	I 3	2	vF. S. E. $20^\circ$ np ff. contains 2vFft.
541	—	8 Leo. min.	f	9	41	n	O 50	3	cF. S. iR. gbM. r. $1\frac{1}{2}$ dia.
542	—	21 — — —	p	7	55	n	O 8	3	cF. vL. iF. $5'1.4'$ b. sp. a double ft.
543	April 17	11 ( $\zeta$ ) Virgin	f	37	39	f	I 31	I	eF. pL.
544	—	— — — —	f	43	12	f	I 23	I	vF. vS.
545	—	— — — —	f	62	44	f	I 9	I	eF. cS. er.
546	} 29	64 — — —	f	36	17	n	I 4	I	} Two. Both vF. vS. r. the place betw. them. sp nf. but near. mer.
547									
548	30	43 ( $\delta$ ) — — —	p	0	31	f	O 30	I	vF. cS. with 240 lE. near vSft.
549	—	84 ( $\epsilon$ ) — — —	f	9	1	f	I 13	I	eF. vS. stellar. confir. 240.
550	—	109 — — —	p	5	32	n	I 35	I	vF. S. p. and in a line with 2Bft.
551	} May 1	31 Bootis	p	23	9	f	O 42	I	} Two. Both eF. vS. The place is that of the f. dist. 3 or 4'
552									
553	3	50 ( $\sigma$ ) Serpen	p	12	7	f	O 15	I	cF. iF, r. $5'1.3'$ b.
554	27	3 — — —	p	21	20	f	I 19	I	vF. S. E. np ff. but nearly mer.
555	June 22	101 Hercul	p	2	55	f	I 28	I	cF. S. lE. iF. r.
556	Sept. 4	71 ( $\epsilon$ ) Piscium	f	22	10	n	I 24	I	vF. mE. $75^\circ$ sp nf. $1\frac{1}{2}$ l.
557	18	85 Ceti	p	6	18	n	O 52	I	vF. vS. lE. r. 240 the same.
558	20	97 Aquarii	p	14	9	f	O 33	I	eF. cL. iR. 5 or 6' dia.
559	—	54 Eridani	p	65	18	f	O 59	I	3vSft. in a line with vF. nebulosity.
560	21	45 Androm	f	16	14	f	O 32	I	vF. S. E. among ft.
561	—	58 — — —	p	17	32	f	I 34	I	vF. stellar.
562	} —	— — — —	p	15	22	f	I 48	I	} Four. stellar. unequal. Three in a row, and the fourth making a rectangle with them. That at the angle is much larger.
563									
564									
565									
566	—	— — — —	p	5	4	n	O 14	2	vF. pL. iR.
567	—	— — — —	f	2	29	f	O 12	I	vF. S. lE.
568	30	17 Eridani	p	8	17	n	2 17	I	eF. S. iF. among 3 or 4 ft.
569	—	— — — —	f	9	13	n	O 27	I	eF. lE. er.
570	Oct. 17	26 ( $\beta$ ) Persei	p	43	39	n	O 51	I	eF. vS. lE.
571	—	— — — —	p	42	9	n	O 44	I	eF. stellar. not verified.
572	} —	— — — —	p	32	26	f	O 11	I	} Two. Both vF. vS. er. dist. 4' the place between them.
573									
574	} —	— — — —	f	13	6	n	O 27	I	} Two. Both vF. stellar. vlbM. but the f. is the brightest and largest.
575									
576	18	12 Androm	p	24	27	f	I 47	I	vF. S. iR. stellar.
577	—	53 ( $\tau$ ) — — —	p	18	55	f	O 8	I	vF. pL. lE. lbM.
578	—	28 ( $\omega$ ) Persei	p	2	50	f	I 16	I	vF. vS.
579	24	17 ( $\nu$ ) Andro	p	3	21	n	I 3	I	vF. vS. just f. pBft.
580	—	30 Persei	p	20	43	f	I 3	I	suspected. r. some ft. visible.
581	25	40 Arietis	p	8	24	f	O 18	I	vF. E. iF. time inaccurate.
582	—	— — — —	p	1	17	f	2 7	I	vF. S. iF.
583	26	10 ( $\alpha$ ) Triang	p	18	21	f	O 29	I	vF. vS, E. or 3Fft. with vF. Nebul.



III.	1786	Stars.		M. S.		D.M.	Obj	Description.
584	Oct. 26	35 Arietis	p	0 41	n	0 50	1	vF. S. bM.
585	Nov. 26	48 (v) Eridani	p	3 33	f	1 2	1	suspected; hazy weather.
586	—	— — —	p	3 6	f	0 56	2	{ eF. S. E. nearly par. another suspec. 3' ff. stellar.
587	28	42 (ξ) —	f	2 34	n	0 9	1	vF. S. bM. betw. 2 ft.
588	—	— — —	f	7 35	f	1 57	1	vF. S.
589	—	— — —	f	10 10	f	1 22	1	vF. cL. iE. nearly par. bM.
590	Dec. 14	8 Leporis	f	9 18	f	0 6	1	eF. stellar. a little doubtful.
591	15	13 (ζ) Eridani	p	4 35	f	0 6	1	eF. stellar. about 1' nf. II. 286.
592	}	20 Neb. II. 447.	p	0 0	f	0 2	1	{ Two. The p. vF. vS. The next eF. eS. and left doubtful.
593								
594	—	26 Ceti	f	18 21	f	0 23	1	vF. mE. bM. 3½ l. 1½ b.
595	21	29 —	p	28 42	n	1 17	1	vF. S. some ft. in it.
596	—	44 Hydræ	p	34 21	n	0 50	2	vF. S. lbM. ff. a trapezium of S. ft.
597	24	— — —	p	59 13	n	2 44	1	vF. S. R. vglbM.
598	30	59 (c) Leonis	f	2 40	f	1 19	1	eF. S. iE. not verified.
1787								
599	Jan. 11	55 (δ) Gemini	f	68 4	f	0 12	1	eF. pL. r.
600	14	30 (n) Leonis	p	11 47	f	0 20	1	vF. S. iR.
601	—	— — —	p	11 4	n	0 3	1	vF. cS. iE. er.
602	—	29 Comæ	p	12 7	n	0 6	1	vF. cL. vgbM. f. cBft.
603	—	— — —	p	6 16	n	0 11	1	vF. E. np ff. 2½ l.
604	17	58 Androm	f	2 45	f	0 21	1	vF. stellar. confir. 240.
605	Feb. 10	9 (1st μ) Canc	p	3 15	n	0 46	1	vF. S. iF.
606	13	10 (2d μ) —	f	11 31	f	1 1	2	vF. S. stellar.
607	—	33 (n) —	p	12 33	n	0 38	1	vF. vS.
608	22	69 (v) —	f	2 5	n	0 33	1	eF. S. R. vlbM.
609	—	21 (θ) Crater	f	2 28	n	0 28	1	vF. vS. R. with 240 gbM.
610	—	65 Virginis	p	33 51	f	0 12	1	cF. pL. E.
611	—	— — —	p	32 29	n	0 50	1	vF. S. no time to verify.
612	March 11	87 (e) Leonis	f	23 23	f	0 57	1	vF. cS. E.
613	—	44 (k) Virgin	p	1 42	n	0 8	1	vF. E. er.
614	—	— — —	f	0 57	f	0 49	1	cF. S. iR.
615	17	38 Leo. min.	p	1 27	f	0 27	2	cF. S. er.
616	—	6 Canum	p	34 40	f	1 1	2	vF. cL. iF. 4' dia. 5' f. ft. 6 m.
617	—	— — —	p	27 3	f	1 13	2	eF. pL. iR. 1' dia. or more.
618	—	12 —	p	3 8	f	1 31	1	eF. vS.
619	—	17 —	f	11 29	n	0 2	1	vF. S. E. nearly mer.
620	—	— — —	f	26 36	f	0 14	1	cF. E. nearly par. r. ¾ l.
621	—	— — —	f	38 29	f	0 46	1	vF. S. iR. conf. 300.
622	—	12 (λ) Coronæ	f	7 17	f	0 37	1	vF. S. R. discov. in gaging.
623	—	— — —	f	24 7	f	0 19	1	vF. vS. n. 2 ft. 300 confir.
624	—	— — —	f	27 24	f	0 9	1	vF. S. bM. discov. with 300.
625	18	10 (n) Ursæ	p	2 41	f	2 27	1	vF. vS. 300.
626	—	— — —	f	7 14	f	0 4	2	vF. S. iF. lbM. r.

III.	1787	Stars.		M.	S.		D.M.	Ob	Description.	
627	March 18	43 Lyncis	p	17	50	f	1	2	vF. vS. stellar. 300.	
628	—	—	p	16	48	f	0	9	cF. cS.	
629	}	—	p	15	24	f	0	8	1	{ Two. Both vF. vS. dist. 3'. nearly mer. 300.
630										
631	—	34 ( $\mu$ ) Urfæ	f	3	39	f	1	55	2	vF. S. R. 300.
632	—	47 —	p	2	8	n	0	32	2	cF. vS. lE. mer. gmbM.
633	—	20 Canum	f	1	58	f	0	13	1	vF. S. lbM.
634	—	54 ( $\phi$ ) Bootis	p	1	24	f	0	36	1	vF. vS. conf. 300 sp. 2 vBst.
635	}	—	f	6	42	n	0	46	1	{ Two. The nf. vF. vS. verif. 300. The sp. discov. with 300 eF. S. iF.
636										
637	—	30 ( $g$ ) Hercul	p	24	18	f	1	5	1	vF. eS. 300. shewed 2 vSft. with nebu.
638	—	—	p	3	33	f	0	59	1	vF. vS.
639	—	—	p	2	53	f	1	24	1	eF. eS.
640	—	—	f	0	54	f	1	5	1	vF. vS.
641	—	—	f	1	10	f	1	16	1	vF. vS.
642	19	70 Virginis	f	1	37	f	0	26	1	vF. S. iF. time l. inaccurate.
643	—	—	f	3	43	n	0	5	1	vF. S. lE. just ff. ft.
644	—	5 ( $\nu$ ) Bootis	f	25	41	f	0	44	1	vF. vS. E. confir. 300.
645	—	30 ( $\zeta$ ) —	p	10	16	n	0	17	1	eF. vS. lbM. betw. 2 vFft. 300.
646	—	28 ( $\beta$ ) Serpen	f	11	15	n	0	24	1	vF. S. lE.
647	20	44 Lyncis	p	33	11	f	2	16	1	vF. vS. verif. 300.
648	—	13 Canum	p	36	41	n	0	46	1	vF. E. par. 1' l.
649	—	—	f	12	29	n	1	0	1	vF. S. lE.
650	—	—	f	19	54	n	2	18	1	eF. vS.
651	—	—	f	27	11	n	1	11	1	vF. S.
652	—	—	f	29	47	n	0	17	1	eF. vS.
653	—	—	f	62	59	n	1	34	1	vF. pS. E. mer. 300.
654	April 9	19 —	p	7	36	n	0	51	1	vF. vS. lbM.
655	—	—	p	*	26	n	2	57	1	vF. pS. lbM. *forgot, but is 5, 6, or 7'.
656	—	20 —	f	15	21	n	1	15	1	vF. vS. lbM.
657	}	—	f	83	33	n	1	57	1	{ Two. Both vF. vS. E. in differ. di- rections. 2 or 3' dist. par. each f. Sft.
658										
659	—	—	f	117	16	n	0	18	1	vF. vS. r.
660	—	—	f	119	18	n	1	16	1	eF. cS.
661	—	—	f	125	25	n	0	44	1	eF. S.
662	11	29 ( $\gamma$ ) Virgin	f	2	8	n	0	54	1	vF. pL.
663	—	—	f	3	33	n	0	55	1	vF. S. iF.
664	—	—	f	6	6	f	0	14	1	vF. S.
665	15	90 ( $p$ ) —	p	1	48	n	0	23	2	cF. cL. R. vlbM. r. 5' dia.
666	—	102 (1stv) —	p	19	18	f	0	55	1	eF. vS.
667	—	—	p	18	53	f	0	33	2	eF. vS. verif. 300. 2d obf. vF. S.
668	—	105 ( $\phi$ ) —	p	3	29	f	0	58	1	cF. S. r.
669	May 7	61 —	p	5	42	n	1	30	1	vF.
670	—	—	p	2	45	n	1	46	1	vF.
671	—	8 Libræ	p	9	38	f	1	28	1	cF. S. R. sp and joining 2 Sft.

III.	1787	Stars.		M.	S.	D.M.	Ob	Description.
672	May 12	19 ( $\lambda$ ) Bootis	p	48	58	n	0 41 3	cF. vS. stellar. 300.
673	—	—	p	38	30	n	2 21 1	cF. S. R. or lE.
674	—	—	p	5	52	n	2 31 1	cF. cS. iR.
675	—	38(2d b) —	p	11	18	n	0 35 1	vF. pS. iF. sp. 2 S. unequal ft.
676	15	24 (g) —	p	16	34	n	0 35 2	cF. cS. lE nearly par.
677	—	—	p	2	56	f	1 15 1	vF. pS. lE.
678	}	*A Bootis 7m	f	0	3	n	0 3 1	{ Two. The p. vF. vS. The f. eF. eS. * See note.
679			f	0	48	n	0 1 1	
680	—	42 Herculis	p	13	17	n	1 0 2	vF. S. R. lbM. er. near some Sft.
681	16	*C Can <sup>m</sup> 6m.	f	0	44	f	0 15 1	cF. vS. lE. * See note.
682	—	—	f	7	35	f	0 7 1	eF. cS. E. sp. Sft.
683	—	—	f	12	35	f	0 25 1	cF. pL. iF.
684	—	—	f	13	49	n	0 34 1	vF. vS. R.
685	—	27 ( $\gamma$ ) Bootis	p	19	48	n	1 6 2	vF. cS. R. fbM.
686	—	—	f	8	42	n	0 18 1	eF. cS. lbM.
687	—	—	f	13	27	n	0 23 1	cF. pS. another suspec. 2' n. 300.
688	—	16( $\tau$ ) Coronæ	f	7	34	f	0 48 1	vF. cS. iR.
689	—	67 ( $\pi$ ) Hercu	p	20	32	f	0 11 1	eF. cL. iE. nearly par.
690	19	10 Libræ	p	4	6	f	0 43 1	vF. cS. iF. lbM.
691	—	—	f	6	51	f	0 59 1	cF. fmbM. stellar.
692	Aug. 12	33 ( $\iota$ ) Aquarii	p	5	23	n	0 38 1	eF. E. np ff. 2' l. 1' b.
693	Sept. 11	41 —	p	11	30	n	0 36 1	eF. vS. 360 confirmed it.
694	Oct. 11	50 (f) Cassio	f	90	22	n	0 30 1	vF. vS. iR. bM.
695	Nov. 3	10 Camelop	p	155	0	n	0 53 1	eF. pL. iF.
696	5	17 ( $\xi$ ) Cephei	p	16	35	f	0 47 2	vF. S. R. lbM. r. 1' dia.
1788								
697	Jan. 14	67 Urfæ	f	11	9	n	0 40 3	vF. E. np ff. 5' l. 1' b.
698	—	27 ( $\gamma$ ) Bootis	p	39	53	n	1 29 2	vF. S.
699	—	—	p	38	20	n	2 8 2	vF. S. iF.
700	Feb. 3	45 ( $\omega$ ) Urfæ	p	14	53	f	1 33 1	cF. L. iE. mb. f. M. 4' l. 2 $\frac{1}{2}$ b.
701	—	—	p	6	6	f	0 1 1	vF. vS. iF.
702	—	13 Canum	p	42	13	f	0 55 1	vF. vS.
703	5	71 ( $\theta$ ) Gemin.	p	10	3	f	0 46 1	vF. vS. perhaps a patch of ft.
704	—	60 Urfæ	p	81	11	f	0 22 1	eF. vS. perhaps a patch of Sft.
705	—	—	p	39	57	f	0 43 1	vF.
706	—	—	p	23	49	n	0 38 2	vF. vS. lE. f. cBft.
707	—	63 ( $\chi$ ) —	f	11	2	n	0 34 2	vF. vS. another fusp. ff. eF. eS.
708	6	59 —	f	30	14	f	0 31 1	vF. vS. in a line with 2 ft. nf sp.
709	March 9	21 Lyncis	f	34	50	n	1 41 1	vF. R. vgbM. 2 $\frac{1}{2}$ dia.
710	—	9 ( $\iota$ ) Urfæ	p	45	51	n	0 49 1	vF. iF. 2 $\frac{1}{2}$ l. 1 $\frac{3}{4}$ b.
711	—	—	p	41	10	n	1 49 1	eF. E. sp nf. 3 $\frac{1}{2}$ l. 2 $\frac{1}{2}$ b.
712	—	—	p	4	49	n	1 6 1	eF. cS. r. p. some Fft.
713	—	—	f	25	7	n	1 15 1	cF. cS. lE.
714	—	—	f	24	58	n	1 11 1	cF. cS. lE.
715	—	63 ( $\chi$ ) —	f	3	26	n	0 39 1	eF. pS.

III.	1787	Stars.		M. S.		D.M.	Ob	Description.
716	March 9	63 ( $\chi$ ) Urfæ	f	5 2	n 2	26	I	vF. vS.
717	—	3 Canum	p	14 1	n 0	37	I	cF. mE. nearly mer. 5' l.
718	—	—	p	4 6	f 0	51	I	vF. vS.
719	}	—	p	2 47	f 1	31	I	} Two. Both vF. vS. dist. 1' in the same meridian.
720		—						
721	—	—	f	32 1	f 1	21	I	vF. S.
722	11	49 ( $g$ ) Virgin	p	18 9	f 0	21	I	eF. S.
723	April 1	Neb. II. 728.	p	0 25	f 0	2	I	eF. vS.
724	8	61 Virginis	f	1 43	f 2	23	I	cF. vS. iF.
725	10	60 Urfæ	f	39 40	f 1	14	2	eF. cL. iR. lbM. 3' dia.
726	—	—	f	42 45	f 0	34	2	vF. pS. R.
727	12	35 ( $\sigma$ ) Hercul	f	16 11	n 0	14	I	cF. S. E. par.
728	13	42 —	f	20 46	n 0	54	I	vF. cS. iR.
729	27	19 ( $\lambda$ ) Bootis	p	113 28	f 0	3	I	vF. S.
730	28	27 ( $\beta$ ) Hercul	f	4 6	n 0	2	I	eF. vS. E.
731	29	27 ( $\gamma$ ) Bootis	p	15 47	n 1	16	I	vF. vS.
732	—	—	p	15 33	n 1	22	I	vF. vS. lE.
733	—	—	p	9 25	n 2	4	I	vF. vS.
734	—	—	p	8 52	n 2	8	I	cF. pS.
735	—	22 ( $\tau$ ) Hercul	f	30 17	f 1	2	I	eF. pS. with 300 iF.
736	30	21 (1 ft.) Libr	f	7 7	n 1	59	I	vF. pL. E. mer. lbM. 300.
737	May 1	23 ( $\theta$ ) Bootis	f	49 59	f 1	46	I	vF. vS. stellar.
738	25	12 ( $i$ ) Draco	f	17 8	n 0	44	I	vF. vS.
739	June 2	14 ( $n$ ) —	p	32 30	n 0	57	I	vF. R. vgbM. er. 3' dia.
740	3	15 (A) —	p	10 14	f 3	21	I	cF. pL. iE.
741	6	31 (1 ft. $\Psi$ ) —	p	5 13	f 0	5	I	eF. stellar. with 300 lE. par.
742	July 8	*B Draco 7m.	f	4 25	f 0	27	I	vF. stellar. verif. 300. * See note.
743	30	19 Aquilæ	f	9 24	n 0	26	I	cF. iR. r. 3 or 4' dia.
744	Aug. 2	51 —	p	8 8	n 0	29	I	vF. pL. R. vgmbM.
745	Nov. 1	27 ( $\delta$ ) Cephei	f	26 10	f 1	24	I	vF. pL. iF. er.
746	—	36 Camelop.	f	64 5	f 0	38	I	vF. S. R. lbM.
747	Dec. 3	*22 Cam Hev	p	37 1	f 0	8	I	{ cF. pL. iF. mbM. er. some ft. visible. * See note.

## Fourth class. Planetary nebulae.

Stars with burs, with milky chevelure, with short rays, remarkable shapes, &amp;c.

IV.	1785	Stars.		M. S.		D.M.	Ob	Description.
30	May 1	14 Canum	p	6 48		55	2	Two ft. dist. 3' connected with a vF. narrow nebulosity.
31	Oct. 3	50 Aquarii	f	7 55		37	1	F. S. stellar. with pL. chev.
32	5	62 ( $b$ ) Eridani	f	0 35	n 0	21	2	vB. vS. mbM. like a ft. affected with irregular burs.
33	—	49 ( $d$ ) Orion	p	2 33	n 0	28	4	A ft. with m. chev. or vBN. with m. nebulosity.

IV.	1785	Stars.		M.	S.		D.M.	Ob	Description.	
34	Dec. 28	40(2dφ) Orio	f	5	41	f	0	12	2	cB. S. nearly R. like a ft. with L. dia. with 240 like an ill defined planetary neb.
35	31	9 Hydrae	p	8	19	f	0	14	1	A S ft. with a brush sp. FS. it resembles fig. 7. Phil. Transf. Vol. LXXIV. Tab 17.
36	1786 Jan. 1	60 Orionis	p	11	38	f	0	20	3	A ft. affected with vF. extensive m. chev. The ft. not quite central.
37	Feb. 15	28 (ω) Draco	f	20	33	f	2	12	1	A planetary neb. vB. has a disk of about 35'' dia. but very ill defined edge. With long attention a vB. well defined R. center becomes visible.
38	24	55 Orionis	f	18	3	n	1	17	2	A cft. affected with vF. m. chev.
39	March 19	2 Navis	p	3	32	f	0	5	1	pB. R. r. within the 46th of the <i>Connoiss. des Temps</i> almost of an equal light throughout 2' dia. no connection with the cluster, which is free from nebulosity.
40	27	68 (ι) Virgin	p	30	45	f	0	18	1	A pBft. with a seeming brush to it np. may be a vS neb. close to it.
41	May 26	14 Sagittarii	p	11	58	f	1	15	1	A double ft. with extensive nebulosity of different intensity. About the double ft. is a black opening resembling the neb. in Orion in miniature.
42	Sept. 30	51 Ceti	f	7	26	n	0	27	1	A ft. about 8 or 9 m. with vF. branches. each branch 1' l.
43	Oct. 17	26 (β) Persei	p	2	48	n	1	54	2	A pBft. with 2 F. branches.
44	Nov. 28	5 Monocero	p	7	16	f	0	2	1	A ft. involved in m. chev.
45	1787 Jan. 17	55 (δ) Gemin	f	9	6	f	1	1	2	A ft. 9 m. with a pB. m. nebulosity. equally dispersed all around. A very remarkable phenomenon.
46	Feb. 22	99 (ι) Virgin	p	4	38	n	0	57	1	pB. almost cB. vS. stellar. like a star with burs.
47	March 11	44 (k) —	f	1	48	f	0	46	1	pB. stellar. resembles a ft. with a bur all around.
48	18	19 Leo. min	f	6	32	f	0	17	1	A vFft affected with vF. nebulosity. E. sp nf. 1' l. 3co.
49	April 15	102 (1ftv) Vir	p	6	9	f	0	52	2	pB. stellar. like a ft. with a S. bur all around.
50	May 12	77 (z) Hercul	p	40	13	f	0	28	1	vB. R. 4' dia. almost equally B. with a F. r. margin.

IV.	1787	Stars.		M.	S.		D.M.	Ob	Description.	
51	Aug. 8	61 (g) Sagitt	p	13	56	n	1	23	2	A cB. S. beautiful planetary nebula; but c. hazy on the edges, of a uniform light; 10 or 15'' dia. perfectly R. I shewed it to M. DE LA LANDE.
52	Nov. 3	4 (d) Cassio	p	4	0	f	1	6	2	A ft. 9 m. with vF. nebulosity of S. extent about it.
53	—	10 Camelop	p	55	42	n	0	11	2	A pB. planetary nebula. near 1' dia. R. of uniform light and pretty well defined. 2 obf. with 360 magnified in proportion; but still pretty abruptly defined, and a little elliptical.
	1788									
54	Jan. 14	67 Urfæ	f	7	32	f	0	30	1	cB. S. N. with F. chev.
55	Feb. 6	34 Lyncis	p	28	4	n	0	2	2	pB. R. almost of an even light throughout, approaching to planetary, but ill defined and a little fainter on the edges $\frac{3}{4}$ or 1' dia. p. 1' pc ft.
56	—	59 Urfæ	f	25	11	n	0	56	1	cB. iR. cBNM. with extensive chev. 5' dia.
57	June 11	35 ( $\sigma$ ) Hercul	f	34	27	f	0	18	2	AvS.F. ft. involved in eF. nebulosity.
58	Nov. 25	24 Cephei	f	116	28	n	0	2	1	A ft. 9 m. surrounded with vF. m. nebulosity. The ft. is either double, or not R. Less than 1' dia.

## Fifth class. Very large nebulae.

V.	1785	Stars.		M.	S.		D.M.	Ob	Description.	
25	Nov. 27	18 Ceti	f	1	30	n	1	2	1	Four or five pL. ft. forming a trapezium of about 5' dia. The inclosed space is filled up with faintly terminated m. nebulosity. The ft. seem to have no connexion with the nebulosity.
26	Dec. 7	18 Leo. min.	p	8	7	n	1	1	2	cB. mE. par. 8' l. 3' b.
27	26	15 Monocero	p	0	12	f	0	6	2	Some pBft. 7 or 8' sp. 15th Monce. are involved in eF. m. nebulosity which loses itself imperceptibly.
	1786									
28	Jan. 1	48 ( $\sigma$ ) Orion	f	2	46	n	0	44	2	Remarkable m. nebulosity, divided in 3 or 4 large patches, including a dark space; cannot

V.	1786	Stars.		M.	S.		D.M.	Ob	Description.	
29		261 Ursa	f	45	38	f	0	40	1	take up less than $\frac{1}{2}$ degree, but I suppose it to be much more extensive.
30	18	42 } c Orioni 45 }	p	0	0	n	0	0	2	eF. vL. vlbM. r. 10' l. 8 or 9' b. The 1st and 2d c Orionis, and the stars about them, are involved in eF. unequally B. m. nebulosity.
31	31	44 (i) —	p	0	0	n	0	0	2	Orionis with its neighbouring st. are involved in eF. m. nebulosity to a great extent.
32	Feb. 1	28 (n) —	p	17	26	f	i	4	2	cB. vL. m. diffused and vanishing. near and ff. Bst.
33	—	— — —	f	1	26	f	0	7	1	Diffused eF. m. nebulosity. The means of verifying this phenomenon are difficult.
34	—	46 (e) Orionis	p	0	0	n	0	0	1	I am pretty certain e Orionis is involved in unequally diffused m. nebulosity.
35	—	36 (v) — 56 —	f p	3 2	39 16	f n	0 0	40 28	4	Diffused m. nebulosity, extending over no less than 10 degrees of PD. and many degrees of RA. It is of very different brightness, and in general extremely F. and difficult to be perceived. Most probably the nebulosities of the 28th, 30, 31, 33, 34, and 38th of this class are connected together, and form an immense stratum of far distant stars, to which must also belong the nebula in Orion.
36	Oct. 17	35 (v) Andro	p	9	8	f	0	20	2	vF. vL. E. nearly mer. or a little from np ff. about 20' l.
37		2457 Cygni	f	5	1	f	1	1	1	vL. diffused nebulosity. bM. 7 or 8' l. 6' b. and losing itself vg. and imperceptibly.
38	Dec. 20	19 (beta) Orion	f	11 11	9 35	n f	1 0	19 52	1	Strongly suspected nebulosity of v. great extent. Not less than 2° 11' of PD. and 26'' of RA. in time.
39		2111 (beta) Crater	p	8	15	f	0	17	2	vF. mE. nearly par. or about 10° sp nf. vgbM. 8' l. 3' b.

V.	1786	Stars.		M.	S.		D.M.	Ob	Description.
40	Dec. 21	11 ( $\beta$ ) Crater	p	7	49	f	0 26	2	vF. mE. $15^\circ$ sp nf. vlbM. about 7' l. 4' b.
	1787								
41	March 17	6 Canum	p	8	27	f	1 12	1	vB. E. $60^\circ$ sp nf. 20' l. 2' b.
42	20	13 —	p	18	39	n	1 48	1	vB. mE. sp nf. but nearly par. mbM. 16' l.
	1788								
43	March 9	3 —	p	0	38	f	1 41	3	v brilliant. BN. with Fm. bran. np ff. 15' l. and to the ff. running into vF. nebulosity extending a great way. the N. is not R.
44	Nov. 1	36 Camelop	f	84	33	n	0 23	2	cB. R. vgbM. BN. 6 or 7' dia. with a F. branch extending a great way to the np. side; not less than $\frac{1}{2}$ degree. and to the n. or nf. the nebulosity diffused over a space not less than a whole degree.

Sixth class. Very compressed and rich clusters of stars.

Additional }  
abbreviations. }

Cl. Cluster.  
fc. scattered.

com. compressed.  
co. coarsely.

VI	1785	Stars.		M.	S.		D.M.	Ob	Description.
20	Oct. 27	18 ( $\epsilon$ ) Pif. Aufst	f	133	24	n	0 23	2	cB. iR. 8 or 9' dia. a great many of the st. visible, so that there can remain no doubt but that it is a Cl. of vS. stars.
21	Dec. 7	25 Gemino	f	2	15	f	1 15	1	A v. rich and v. com. Cl. st. of. about 5' dia. some of the largest st. are in a row.
	1786								
22	Feb. 1	31 Monocero	p	30	4	n	1 20	4	A beautiful Cl. of much com. st. confid. rich. 10 or 12' dia. C. H. discovered it in 1783.
23	June 27	46 ( $\nu$ ) Sagitt	p	49	15	f	0 42	1	A beautiful Cl. of vS. st. of various sizes. 15' dia. very rich.
24	Oct. 17	58 ( $\nu$ ) Cygni	f	15	56	n	1 18	2	A v. com. and v. rich Cl. of eSst. about 6' l. 4' b. nearly par.
25	Dec. 11	27 ( $\ast$ ) Persei	f	5	55	n	2 25	2	A beautiful com. and rich Cl. of S. and L. st. 7 or 8' dia. the L. st. arranged in lines like interwoven letters.



VI.	1786	Stars.		M.	S.		D.M.	Ob	Description.
26	Dec. 11	53 ( <i>d</i> ) Persei	f	13	34	f	1 13	1	A vF. and v. com. Cl. of eS. ft. near 4' dia.
27	27	22 Monocero	p	20	9	n	0 51	1	A v. beautiful Cl. of much com. S. and L. ft. above 20' dia.
	1787								
28	Jan. 11	75 ( <i>l</i> ) Orionis	f	21	25	n	1 2	1	A Cl. of e. com. and eS. ft. c. rich iF. the f. and most com. part R.
29	Oct. 14	3 Lacertæ	p	7	52	n	2 7	1	A com. Cl. of eS. ft.
30	18	7 ( <i>q</i> ) Cassiop	f	3	10	f	0 46	3	A beautiful Cl. of v. com. Sft. v. rich. C. H. discovered it 1783.
31	Nov. 3	37 ( <i>δ</i> ) —	f	19	48	n	1 2	1	A beautiful Cl. of pL. ft. near 15" dia. conf. rich.
	1788								
32	Sept. 21	80 ( <i>ι</i> ft π) Cyg	p	11	26	n	0 28	1	A beautiful Cl. of p: com. ft. 8 or 9' dia. nearly R. c. rich.
33	Nov. 1	7 ( <i>χ</i> ) Persei	f	1	7	f	0 22	1	A v. beautiful and brilliant Cl. of L. ft. v. rich. the M. contains a vacancy.
34	—	—	f	4	0	f	0 23	1	A v. beautiful, brilliant Cl. of L. ft. iR. v. rich. near $\frac{1}{2}$ degree in dia.
35	26	15 ( <i>κ</i> ) Cassiop	p	1	22	f	1 26	1	A S. Cl. of vF. and e. com. ft. about 1' dia. The next step to an er. neb.

Seventh class. Pretty much compressed clusters of large or small stars.

VII.	1785	Stars.		M.	S.		D.M.	Ob	Description.
18	July 17	12 Vulpeculæ	p	7	56	n	0 44	1	An E. Cl. of i. sc. ft. of various sizes. c. rich.
19	30	21 Aquilæ	p	5	49	n	1 55	1	A p. com. Cl. of p. sc. ft. of var. sizes, magnitudes, and colours. iF. and unequally com. 12 or 15' dia.
20	Nov. 1	7 Monocero	f	1	3	n	0 35	3	A beautiful Cl. of p. com. and equally sc. ft. 10' or 12' dia.
21	Dec. 26	109 ( <i>n</i> ) Tauri	p	14	59	n	1 37	1	A Cl. of p. com. ft. with many eS. ft. mixed with them.
22	28	13 Monocero	f	2	48	n	0 21	1	A S. Cl. of p. com. vS. ft.
23	30	31 ( <i>n</i> ) Canis	f	32	6	f	0 39	1	A com. Cl. of pL. ft. c. rich.
	1786								
24	Jan. 1	60 Orionis	p	5	9	f	0 9	2	A Cl. of p. com. pS. sc. ft. with many eS. fuspec. betw. them 7 or 8' dia.
25	27	8 Monocero	p	11	46	n	0 49	1	A Cl. of p. com. ft. of several sizes 4 or 5' dia. with extensively straggling ones.

VII.	1786	Stars.		M. S.		D.M.	Ob	Description.
26	Jan. 30	6 Monocero	f	8 59	n	1 7	1	A Cl. of eS. and pm. com. ft. with a few L. but not rich. in the shape of a hook.
27	Feb. 24	11 ———	f	42 13	f	1 21	2	An i. Cl. of eS. ft. c. com. 9 or 10' l. 4 or 5' b. with an extending bran. towards sp. C. H. discov. 1783.
28	Mar. 19	2 Navis	p	8 23	n	0 47	1	A Cl. of pS. ft. p. rich. 15' dia.
29	April 30	5 (ε) Scorpii	p	7 14	n	0 38	1	A Cl. of vS. ft. p. rich 6' l. 4' b. in the form of a parallelogram.
30	May 26	14 Sagittarii	p	1 35	n	0 9	1	A Cl. of pS. fc. ft. above 15' dia.
31	—	—	f	1 29	f	0 25	1	A Cl. of vS. and p. com. ft. c. rich. 2 or 3' dia.
32	Sept. 21	58 Androm	p	10 49	f	0 8	4	A vL. co. fc. Cl. of vL. ft. iR v. rich. takes up $\frac{1}{2}$ degree like a nebulous ft. to the naked eye.
33	Oct. 18	11 (μ) Aurigæ	f	6 32	n	0 54	1	A Cl. of p. com. pS. Sft. c. rich. contains 1 L. the rest are all of a size.
34	Dec. 11	13 (α) ———	f	9 7	n	0 32	1	A Cl. of vF. and vSft. p. com. but not rich. iF. 3' dia.
35	24	70 (ξ) Orionis	f	15 53	f	1 29	1	A Cl. of S. pm. com. ft. with suspected m. nebulosity.
36	26	18 Monocero	p	3 48	n	1 0	1	A Cl. of v. fc. ft. c. rich. and of great extent.
37	27	77 Orionis	f	12 24	n	0 55	1	A Cl. of v. com. eSft. c. rich. 3 or 4' dia. most com. M.
38	—	22 Monocero	p	7 39	n	1 31	2	A beautiful Cl. of vSft. of several sizes. c. com. and rich M. 10 or 12' dia.
	1787							
39	Jan. 17	21 (σ) Aurigæ	f	3 25	f	2 6	1	A p. com. Cl. of Sft. 4' dia.
40	Oct. 14	3 Lacertæ	p	38 31	n	1 35	1	A Cl. of Sft. of several sizes. 3 or 4' dia. p. rich. like a forming one.
41	—	—	f	5 8	n	0 2	2	A S Cl. of ft. p. com. e. rich in vS. ft. The com. part 4 or 5' dia.
42	18	24 (η) Cassio	f	29 41	n	0 26	2	A brilliant Cl. of L. and vS. ft. c. rich.
43	Nov. 3	1 (ε) ———	p	11 41	n	1 25	1	A S. Cl. of vSft. c. com. and p. rich.
44	—	—	f	4 34	n	1 8	2	A Cl. of p com. pLft. c. rich. The ft. arranged chiefly in lines from sp. nf.
45	—	37 (δ) ———	p	9 29	f	1 28	2	A S. p. com. Cl. of ft. not rich. iF. like a forming one.
46	—	—	f	17 23	n	1 44	2	A S. Cl. of pL. ft. c. rich.
47	—	10 Camelop	p	55 40	n	1 37	2	A Cl. of ft. p. rich and c. com. 1E. 3 or 4' dia. iF.

VII.	1787	Stars.		M.	S.		D.M.	Ob	Description.
48	Nov. 9	32 Cassiop	f	17	1	f	1 40	1	A com. Cl. of some pL. and many vS. ft. iR. 6 or 7' dia.
49	—	45 (ε) —	p	11	8	n	0 20	1	A Cl. of some cL. ft. and many eS. so as hardly to be seen. The Lft. arranged in circular order 3 or 4' dia.
50	1788 Sept. 27	81 (2d π) Cyg	p	22	13	f	1 14	1	A few Sft. with suspected nebulo- fity. with 300 many vS. ft. inter- mixed with the former, so as to make a Cl.
51	Oct. 19	71 (g) —	p	5	49	f	0 9	1	A p. com. Cl. of pS. ft. c. rich iR. 5 or 6' dia.
52	—	— — —	p	0	42	n	0 34	1	An extensive Cl. of Lft. c. rich above 20' dia.
53	—	73 (ρ) —	f	30	41	n	0 48	2	A L. Cl. of p. com. cLft. above 15' dia. c. rich.
54	Nov. 1	36 Camelop	f	29	1	n	0 16	1	A vF. patch. or S. Cl. of eSft.
55	23	32 (ι) Cephei	f	57	34	n	1 47	3	A Cl. of cS. ft. iF. p. rich and com. contains a vacancy M.

Eighth class. Coarsely scattered clusters of stars.

VIII.	1785	Stars.		M.	S.		D.M.	Ob	Description.
41	Dec. 7	98 (k) Tauri	f	12	11	f	0 54	1	A co. Cl. of ft. or projecting point of the m. way.
42	—	125 —	p	1	22	f	0 4	2	A Cl. of co. fc. ft. above 15' dia. The ft. nearly of a size and equally fc.
43	26	109 (n) —	p	15	30	n	1 29	1	A Cl. of v. co. fc. Lft. join. to VII. 21.
44	28	5 (n) Can. min	f	0	38	f	1 54	1	A Cl. of v. co. fc. Lft. form a cross. not rich.
45	31	6 Navis	p	32	48	f	0 1	1	A co. fc. Cl. of ft. not rich.
46	—	— — —	p	10	18	n	0 49	1	A vL. but co. fc. Cl. of ft.
47	—	— — —	p	10	27	n	0 39	1	A Cl. of fc. ft. or the m. way crowd- ed with ft. of equal size and colour.
48	1786 Jan. 1	78 Orionis	f	10	59	f	1 9	1	A Cl. of v. fc. ft. of various sizes. above $\frac{1}{2}$ degree of extent.
49	3	*B Gemi. 6m	p	33	23	n	0 35	1	A Cl. of co. fc. Lft. not rich. *See note
50	27	8 Monocero	f	10	58	n	0 49	2	A Cl. of ft. arranged in a broad row. 25' l. 6 or 8' b. not v. com. but p. rich.
51	Feb. 23	11 —	f	25	25	f	0 1	1	A Cl. of v. fc. ft.
52	Mar. 19	2 Navis	p	12	16	n	1 32	1	A Cl. of vL. co. fc. ft. not rich.
53	June 27	46 (ν) Sagitta	p	82	10	f	1 4	1	A Cl. of fc. Sft. 8' dia. not v. rich.

VIII.	1786	Stars.		M. S.		D.M.	Ob	Description.
54	June 27	46 ( $\nu$ ) Sagitta	p	71 19	f	0 25	1	A co. fc. Cl. of cLft. The place is that of a S. triangle.
55	— — — —	— — — —	p	64 17	f	0 23	1	A co. fc. Cl. of Lft.
56	Oct. 17	37 ( $\gamma$ ) Cygni	f	0 53	n	0 32	1	A S. Cl. of co. fc. ft. of various sizes. E. like a forming one.
57	—	58 ( $\nu$ ) —	f	8 47	n	0 20	1	A Cl. of co. fc. pS. ft. of several sizes. not rich.
58	24	57 —	f	3 19	n	0 16	2	A Cl. of pL. fc. ft. not v. rich.
59	—	59 Persei	f	7 59	n	0 21	1	A Cl. of co. fc. pL. ft. not v. rich.
60	Nov. 26	19 Monocero	p	5 3	f	0 23	1	A Cl. of pL. fc. ft. not v. rich. may be a projecting point of the m. way.
1787								
61	Jan. 17	21 ( $\sigma$ ) Aurigæ	p	16 38	f	0 30	1	A Cl. of co. fc. Lft. iF. not rich. like a forming one.
62	Sept. 19	35 ( $\gamma$ ) Cephei	p	4 43	f	4 50	2	A Cl. of co. fc. Lft. not rich. but the ft. are brilliant. one 7 m.
63	Oct. 16	21 ( $\xi$ ) —	f	1 21	f	0 56	1	A S. Cl. of pL. ft.
64	Nov. 3	27 ( $\gamma$ ) Cassiop	f	11 12	n	0 53	2	A forming cluster of p. com. ft. C. H. disc. 1783.
65	—	37 ( $\delta$ ) —	f	17 56	n	0 29	2	A S. Cl. of Sft. not v. rich. C. H. 1783.
66	—	45 ( $\epsilon$ ) —	f	47 9	f	1 58	2	A Cl. of co. fc. cLft. 8 or 10' dia. one 7 m. near M.
67	9	17 ( $\xi$ ) Cephei	p	10 0	f	2 0	1	A Cl. of co. fc. L. and S. ft. 7' dia. like a forming one.
68	12	41 Aurigæ	p	8 57	n	1 9	1	A S. Cl. of fc. ft. not rich one. 7 m. towards the n. but this does not seem connected with the Cl.
69	Dec. 3	18 Androm	p	8 59	f	1 20	1	A Cl. of co. fc. pL. ft. one 8 m. in the ff. part.
1788								
70	Feb. 3	41 ( $\nu$ ) Persei	f	46 17	n	1 28	1	A Cl. of co. fc. Lft. p. rich above 20' dia.
71	March 4	58 Aurigæ	p	1 22	f	0 44	1	A Cl. of co. fc. pL. ft. p. rich the place is that of a double ft. of the 3d class
72	July 30	62 Serpentis	p	27 26	n	0 6	3	A Cl. of co. fc. Lft. C. H. 1783.
73	—	59 ( $\xi$ ) Aquilæ	p	4 2	f	0 34	1	A Cl. of co. fc. ft. with one pBft. M.
74	Sept. 21	80 (1st $\pi$ ) Cyg	p	34 12	f	0 12	1	A Cl. of co. fc. Lft. not rich 6' dia.
75	26	3 Lacertæ	p	7 29	f	2 21	2	A Cl. of co. fc. Lft. lE. sp nf. 16' l.
76	27	59 (1st $f$ ) Cyg	p	4 1	f	0 7	1	A ft. 6 m. surrounded by many cft. forming a brilliant fc. Cl. the Lft. not M. but f.
77	Nov. 1	27 ( $\delta$ ) Cephei	f	17 23	f	0 22	2	A Cl. of co. fc. ft. 8' dia. C. H. 1787.
78	26	15 ( $\nu$ ) Cassio	f	10 56	f	1 8	2	A Cl. of v. co. fc. Lft. take up 15 or 20'. C. H. disc. 1784.

Notes to some nebulæ and clusters of stars.

- I. 138. The number refers to DE LA CAILLE's southern catalogue in the Cœlum Australe Stelliferum.
- I. 190. A star of the sixth magnitude, not contained in any catalogue. I have called it C Canum Venaticorum. It follows FL. 17. Can. Ven.  $37' 34''$  in time, and is  $0^{\circ} 2'$  more south than that star.
- II. 566. See the note to I. 138.
638. See the note to I. 138.
697. See the note to I. 190.
703. A star of the 7th magnitude, not contained in any catalogue. I called it A Ceti. Not having settled its place, I can only give it in a coarse way. RA. about 0 h.  $31' 37''$ , PD. about  $94^{\circ} 22'$ .
- III. 678. A star of the 7th magnitude, not contained in any catalogue. I have called it A Bootis. It follows FL. 39 Bootis  $6' 56''$  in time, and is  $0^{\circ} 55'$  more north.
681. See the note to I. 190.
742. A star of the 7th magnitude, not contained in any catalogue. I have called it B Draconis. Its place very coarsely is RA. 18 h.  $47'$ . PD.  $41^{\circ} \frac{3}{4}$ .
747. See Mr. WOLLASTON's general catalogue. Zone  $20^{\circ}$ .
- VIII. 49. A star of the 6th magnitude, not contained in any catalogue. I have called it B Geminorum. Not having settled its place, I can only give it in a coarse way, RA. about 6 h.  $52' 4''$ . PD. about  $55^{\circ} 17'$ .

P. S. The planet Saturn has a *sixth satellite* revolving round it in about 32 hours, 48 minutes. Its orbit lies exactly in the plane of the Ring, and within that of the first satellite. An account of its discovery with the forty-foot reflector, and a more accurate determination of its revolution and distance from the planet will be presented to the Royal Society at their next Meetings.

WILLIAM HERSCHEL.



XXI. *An Attempt to explain a Difficulty in the Theory of Vision, depending on the different Refrangibility of Light. By the Rev. Nevil Maskelyne, D. D. F. R. S. and Astronomer Royal.*

Read June 18, 1789.

THE ideas of sight are so striking and beautiful, that we are apt to consider them as perfectly distinct. The celebrated EULER, taking this for granted, has supposed, in the Memoirs of the Royal Academy of Sciences at Berlin for 1747, that the several humors of the human eye were contrived in such a manner as to prevent the latitude of focus arising from the different refrangibility of light, and considers this as a new reason for admiring the structure of the eye; for that a single transparent medium, of a proper figure, would have been sufficient to represent images of outward objects in an imperfect manner; but, to make the organ of sight absolutely complete, it was necessary it should be composed of several transparent mediums, properly figured, and fitted together agreeable to the rules of the sublimest geometry, in order to obviate the effect of the different refrangibility of light in disturbing the distinctness of the image; and hence he concludes, that it is possible to dispose four refracting surfaces in such a manner as to bring all sorts of rays to one focus, at whatever distance the object be placed. He then assumes a certain hypothesis of refraction of the differently refran-

refrangible rays, and builds thereon an ingenious theory of an achromatic object-glass, composed of two meniscus glasses with water between them, with the help of an analytical calculation, simple and elegant, as his usually are.

He has not, however, demonstrated the necessary existence of his hypothesis, his arguments for which are more metaphysical than geometrical; and, as it was founded on no experiment, so those made since have shewn its fallacy, and that it does not obtain in nature. Moreover, which is rather extraordinary, it does not account, according to his own ideas, for the very phænomenon which first suggested it to him, namely, the great distinctness of the human vision, as was observed to me, many years ago, by the late Mr. JOHN DOLLOND, F. R. S. to whom we are so much obliged for the invention of the achromatic telescope; for the refractions at the several humors of the eye being all made one way, the colours produced by the first refraction will be increased at the two subsequent ones instead of being corrected, whether we make use of NEWTON'S or EULER'S law of refraction of the differently refrangible rays.

Thus EULER produced an hypothetical principle, neither fit for rendering a telescope achromatic, nor to account for the distinctness of the human vision; and the difficulty of reconciling that distinctness with the principle of the different refrangibility of light discovered by Sir ISAAC NEWTON remains in its full force.

In order to go to the bottom of this difficulty, as the best probable means of obviating it, I have calculated the refractions of the mean, most, and least refrangible rays at the several humors of the eye, and thence inferred the diffusion of the rays, proceeding from a point in an object, at their falling  
upon

upon the retina, and the external angle which such coloured image of a point upon the retina corresponds to.

I took the dimensions of the eye from M. PETIT, as related by Dr. JURIN; and, the specific gravities of the aqueous and vitreous humors having been found to be nearly the same with that of water, and the refraction of the vitreous humor of an ox's eye having been found by Mr. HAWKSBEЕ to be the same as that of water, and the ratio of refraction out of air into the crystalline humor of an ox's eye having been found by the same accurate experimenter to be as 1 to ,68327, I took the refraction of the mean refrangible rays out of air into the aqueous or vitreous humor, the same as into water, as 1 to ,74853, or 1,33595 to 1; and out of air into the crystalline humor as 1 to ,68327, or 1,46355 to 1. Hence I find, according to Sir ISAAC NEWTON'S two theorems, related at Part II. of Book I. of Optics, p. 113. that the ratio of refraction of the most, mean, and least refrangible rays at the cornea should be as 1 to ,74512, ,74853 and ,75197; at the fore-surface of the crystalline as 1 to ,91173, ,91282, and ,91392; and at the hinder surface of the crystalline as 1 to 1,09681, 1,09550, and 1,09420.

Now, taking with Dr. JURIN 15 inches for the distance at which the generality of eyes in their mean state see with most distinctness, I find the rays from a point of an object so situate will be collected into three several foci, *viz.* the most, mean, and least refrangible rays at the respective distances behind the crystalline ,5930, ,6034, and ,6141 of an inch, the focus of the most refrangible rays being ,0211 inch short of the focus of the least refrangible ones.

Moreover, assuming the diameter of the pencil of rays at the cornea, proceeding from the object at 15 inches distance,



to be  $\frac{1}{5}$ th of an inch in a strong light, which is a large allowance for it, the semi-angle of the pencil of mean refrangible rays at their concurrence upon the retina will be  $7^{\circ} 12'$ , whose tangent to the radius unity, or ,1264 multiplied into ,0211 inch, the interval of the foci of the extreme refrangible rays, gives ,002667 inch for the diffusion of the different coloured rays, or the diameter of the indistinct circle upon the retina. Now, I find, that the diameter of the image of an object upon the retina is to the object as ,6055 inch to the distance of the object from the center of curvature of the cornea; or the size of the image is the same as would be formed by a very thin convex lens, whose focal distance is ,6055 inch, and consequently a line in an object which subtends an angle of  $1'$  at the center of the cornea will be represented on the retina by a line of  $\frac{1}{5678}$ th inch. Hence the diameter of the indistinct circle on the retina before found, ,002667 will answer to an external angle of  $,002667 \times 5678' = 15' 8''$ , or every point in an object should appear to subtend an angle of about  $15'$ , on account of the different refrangibility of the rays of light.

I shall now endeavour to shew that this angle of ocular aberration is compatible with the distinctness of our vision. This aberration is of the same kind as that which we experience in the common refracting telescope. Now, by computation from the tabular apertures and magnifying powers of such telescopes, it is certain that they admit of an angular indistinctness at the eye of no less than  $57'$ ; therefore the ocular aberration is near four times less than in a common refracting telescope, and consequently the real indistinctness, being as the square of the angular aberration, will be 14 or 15 times less in the eye than in a common refracting telescope, which may be easily allowed to be imperceptible.

Moreover,

Moreover, Sir ISAAC NEWTON has observed, with respect to the like difficulty of accounting for the distinctness with which refracting telescopes represent objects, that the erring rays are not scattered uniformly over the circle of dissipation in the focus of the object-glass, but collected infinitely more densely in the center than in any other part of the circle, and in the way from the center to the circumference grow continually rarer and rarer, so as at the circumference to become infinitely rare; and by reason of their rarity are not strong enough to be visible, unless in the center and very near it.

He farther observes, that the most luminous of the prismatic colours are the yellow and orange, which affect the sense more strongly than all the rest together; and next to these in strength are the red and green; and that the blue, indigo, and violet, compared with these, are much darker and fainter, and compared with the other stronger colours, little to be regarded; and that therefore the images of the objects are to be placed not in the focus of the mean refrangible rays, which are in the confine of green and blue, but in the middle of the orange and yellow, there where the colour is most luminous, that which is in the brightest yellow, that yellow which inclines more to orange than to green.

From all these considerations, and by an elaborate calculation, he infers, that though the whole breadth of the image of a lucid point be  $\frac{1}{55}$ th of the diameter of the aperture of the object-glass, yet the sensible image of the same is scarce broader than a circle whose diameter is  $\frac{1}{250}$ th part of the diameter of the aperture of the object-glass of a good telescope; and hence he accounts for the apparent diameters of the fixed stars as observed with telescopes by astronomers, although in reality they are but points.

The like reasoning is applicable to the circle of dissipation on the retina of the human eye; and therefore we may lessen the angular aberration, before computed at  $15'$ , in the ratio of 250 to 55, which will reduce it to  $3' 18''$ .

This reduced angle of aberration may perhaps be double the apparent diameter of the brightest fixed stars to an eye disposed for seeing most distinctly by parallel rays; or, if short-sighted, assisted by a proper concave lens; which may be thought a sufficient approximation in an explication grounded on a dissipation of rays, to which a precise limit cannot be assigned, on account of the continual increase of density from the circumference to the center. Certainly some such angle of aberration is necessary to account for the stars appearing under any sensible angle to such an eye; and if we were, without reason, to suppose the images on the retina to be perfect, we should be put to a much greater difficulty to account for the fixed stars appearing otherwise than as points, than we have now been to account for the actual distinctness of our sight.

The less apparent diameter of the smaller fixed stars agrees also with this theory; for the less luminous the circle of dissipation is, the nearer we must look towards its center to find rays sufficiently dense to move the sense. From Sir ISAAC NEWTON's geometrical account of the relative density of the rays in the circle of dissipation, given in his system of the world, it may be inferred, that the apparent diameters of the fixed stars, as depending on this cause, are nearly as their whole quantity of light.

In farther elucidation of this subject let me add my own experiment. When I look at the brighter fixed stars, at considerable elevations, through a concave glass fitted, as I am short-sighted, to shew them with most distinctness, they appear

to me without scintillation, and as a small round circle of fire of a sensible magnitude. If I look at them without the concave glass, or with one not suited to my eye, they appear to cast out rays of a determinate figure, not exactly the same in both eyes, somewhat like branches of trees (which doubtless arise from something in the construction of the eye) and to scintillate a little, if the air be not very clear. To see day objects with most distinctness, I require a less concave lens by one degree than for seeing the stars best by night, the cause of which seems to be, that the bottom of the eye being illuminated by the day objects, and thereby rendered a light ground, obscures the fainter colours blue indigo and violet in the circle of dissipation, and therefore the best image of the object will be found in the focus of the bright yellow rays, and not in that of the mean refrangible ones, or the dark green, agreeable to NEWTON'S remark, and consequently nearer the retina of a short-sighted person; but the parts of the retina surrounding the circle of dissipation of a star being in the dark, the fainter colours, blue, indigo, and violet, will have some share in forming the image, and consequently the focus will be shorter.

The apparent diameter of the stars here accounted for is different from that explained by Dr. JURIN, in his Essay on distinct and indistinct vision, arising from the natural constitution of the generality of eyes to see objects most distinct at moderate distances, and few being capable of altering their conformation enough to see distant objects, and among them the celestial ones, with equal distinctness. But the cause of error, which I have pointed out, will affect all eyes, even those which are adapted to distant objects.


If this attempt to shew the compatibility of the actual distinctness of our sight with the different refrangibility of  
light

light shall be admitted as just and convincing, we shall have fresh reason to admire the wisdom of the Creator in so adapting the aperture of the pupil and the different refrangibility of light to each other, as to render the picture of objects upon the retina relatively, though not absolutely, perfect, and fitted for every useful purpose; “where,” to borrow the words of our religious and oratorical philosopher DERHAM, “all the glories of the heavens and earth are brought and exquisitely pictured.”

Nor does it appear, that any material advantage would have been obtained, if the image of objects on the retina had been made absolutely perfect, unless the acuteness of the optic nerve should have been increased at the same time; as the *minimum visibile* depends no less on that circumstance than the other. But that the sensibility of the optic nerve could not have been much increased beyond what it is, without great inconvenience to us, may be easily conceived, if we only consider the forcible impression made on our eyes by a bright sky, or even the day objects illuminated by a strong sun. Hence we may conclude, that such an alteration would have rendered our sight painful instead of pleasant, and noxious instead of useful. We might indeed have been enabled to see more in the starry heavens with the naked eye, but it must have been at the expence of our daily labours and occupations, the immediate and necessary employment of man.

I shall only mention farther, and obviate an objection to the diffusion of the rays upon the retina by the different refrangibility of light. It may be said, that the ocular aberration, being a separate cause from any effect of the telescope, should subsist equally when we observe a star through a telescope as when we look at it with the naked eye; and that therefore the

fixed stars could not appear so small as they have been found to do through the best telescopes, and particularly by Dr. HERSCHEL with his excellent ones. To this I answer, that the ocular aberration, which is proportional to the diameter of the pupil when we use the naked eye, is proportional to the diameter of the pencil of rays at the eye when we look through a telescope, which being many times less than that of the pupil itself, the ocular aberration will be diminished in proportion, and become insensible.



XXII. *Experiments and Observations on Electricity.* By Mr. William Nicholson; communicated by Sir Joseph Banks, Bart. P. R. S.

Read June 25, 1789.

SECT. I. *On the Excitation of Electricity.*

1. **A** GLASS cylinder was mounted, and a cushion applied with a silk flap, proceeding from the edge of the cushion over its surface, and thence half round the cylinder. The cylinder was then excited by applying an amalgamated leather in the usual manner. The electricity was received by a conductor, and passed off in sparks to LANE'S electrometer. By the frequency of these sparks, or by the number of turns required to cause spontaneous explosion of a jar, the strength of the excitation was ascertained.

2. The cushion was withdrawn about one inch from the cylinder, and the excitation performed by the silk only. A stream of fire was seen between the cushion and the silk; and much fewer sparks passed between the balls of the electrometer.

3. A roll of dry silk was interposed, to prevent the stream from passing between the cushion and the silk. Very few sparks then appeared at the electrometer.

4. A metallic rod, not insulated, was then interposed, instead of the roll of silk, so as not to touch any part of the apparatus.

apparatus. A dense stream of electricity appeared between the rod and the silk, and the conductor gave very many sparks.

5. The knob of a jar being substituted in the place of the metallic rod, it became charged negatively.

6. The silk alone, with a piece of tin-foil applied behind it, afforded much electricity, though less than when the cushion was applied with a light pressure. The hand, being applied to the silk as a cushion, produced a degree of excitation seldom equalled by any other cushion.

7. The edge of the hand answered as well as the palm.

8. When the excitation by a cushion was weak, a line of light appeared at the anterior part of the cushion, and the silk was strongly disposed to receive electricity from any uninfulated conductor. These appearances did not obtain when the excitation was by any means made very strong.

9. A thick silk, or two or more folds of silk, excited worse than a single very thin flap. I use the silk which the milleners call *Perfian*.

10. When the silk was separated from the cylinder, sparks passed between them; the silk was found to be in a weak negative, and the cylinder in a positive state.

The foregoing experiments shew that the office of the silk is not merely to prevent the return of electricity from the cylinder to the cushion, but that it is the chief agent in the excitation; while the cushion serves only to supply the electricity, and perhaps increase the pressure at the entering part. There likewise seems to be little reason to doubt but that the disposition of the electricity to escape from the surface of the cylinder is not prevented by the interposition of the silk, but by a compensation after the manner of a charge; the silk being then as strongly negative as the cylinder is positive: and, lastly, that the line of light between



between the silk and cushion in weak excitations does not consist of returning electricity, but of electricity which passes to the cylinder, in consequence of its not having been sufficiently supplied, during its contact with the rubbing surface.

11. When the excitation was very strong in a cylinder newly mounted, flashes of light were seen to fly across its inside, from the receiving surface to the surface in contact with the cushion, as indicated by the brush figure. These made the cylinder ring as if struck with a bundle of small twigs. They seem to have arisen from part of the electricity of the cylinder taking the form of a charge. This appearance was observed in a nine-inch and a twelve-inch cylinder, and the property went off in a few weeks. Whence it appears to have been chiefly occasioned by the rarity of the internal air produced by handling, and probably restored by gradual leaking of the cement.

12. With a view to determine what happens in the inside of the cylinder, recourse was had to a plate machine. One cushion was applied with its silken flap. The plate was nine inches in diameter and two-tenths of an inch thick. During the excitation, the surface opposite the cushion strongly attracted electricity, which it gave out when it arrived opposite the extremity of the flap. So that a continual stream of electricity passed through an insulated metallic bow terminating in balls, which were opposed, the one to the surface opposite the extremity of the silk, and the other opposite the cushion; the former ball shewing positive, and the latter negative signs. The knobs of two jars being substituted in the place of these balls, the jar, applied to the surface opposed to the cushion, was charged negatively, and the other positively. This disposition of the back surface seemed, by a few trials, to be weaker than  
stronger.

stronger the action of the cushion, as judged by the electricity on the cushion side.

Hence it follows, that the internal surface of a cylinder is so far from being disposed to give out electricity during the friction by which the external surface acquires it, that it even greedily attracts it.

13. A plate of glass was applied to the revolving plate, and thrust under the cushion in such a manner as to supply the place of the silk flap. It rendered the electricity stronger, and appears to be an improvement of the plate machine; to be admitted if there were not essential objections against the machine itself.

14. Two cushions were then applied on the opposite surfaces with their silk flaps, so as to clasp the plate between them. The electricity was received from both by applying the finger and thumb to the opposite surfaces of the plate. When the finger was advanced a little towards its correspondent cushion, so that its distance was less than between the thumb and its cushion, the finger received strong electricity, and the thumb none; and, contrariwise, if the thumb were advanced beyond the finger, it received all the electricity, and none passed to the finger. This electricity was not stronger than was produced by the good action of one cushion applied singly.

15. The cushion in experiment 12. gave most electricity when the back surface was supplied, provided that surface was suffered to retain its electricity till the rubbed surface had given out its electricity.

From the two last paragraphs it appears, that no advantage is gained by rubbing both surfaces; but that a well managed friction on one surface will accumulate as much electricity as the present methods of excitation seem capable of collecting; but

but that when the excitation is weak, on account of the electric matter not passing with sufficient facility to the rubbed surface, the friction enables the opposite surface to attract or receive it, and if it be supplied, both surfaces will pass off in the positive state; and either surface will give out more electricity than is really induced upon it, because the electricity of the opposite surface forms a charge. It may be necessary to observe, that I am speaking of the facts or effects produced by friction; but how the rubbing surfaces act upon each other to produce them, whether by attraction, or otherwise, I do not here enquire.

It will hereafter be seen, that plate machines do not collect more electricity than cylinders (in the hands of the electrical operators of this metropolis) do with half the rubbed surface; which is a corroboration of the inference here made.

16. When a cylinder is weakly excited, the appearances mentioned (par. 8.) are more evident, the more rapid the turning. In this case, the avidity of the surface of the cylinder beneath the silk is partly supplied from the edge of the silk which throws back a broad cascade of fire, sometimes to the distance of above twelve inches. From these causes it is that there is a determinate velocity of turning required to produce the maximum of intensity in the conductor. The stronger the excitation the quicker may be the velocity; but it rarely exceeds five feet of the glass to pass the cushion in a second.

17. If a piece of silk be applied to a cylinder, by drawing down the ends, so that it may touch half the circumference, and the cylinder be then turned and excited by applying the amalgamed leather, it will become very greedy of electricity during the time it passes under the silk. And if the entering surface of the glass be supplied with electricity, it will give it out at the other extremity of contact; that is to say, if insu-

lated conductors be applied at the touching ends of the silk, the one will give, and the other receive, electricity until the intensities of their opposite states are as high as the power of the apparatus can bring them; and these states will be instantly reversed by turning the cylinder in the opposite direction.

As this discovery promises to be of the greatest use in electrical experiments, because it affords the means of producing either the plus or minus states in one and the same conductor, and of instantly repeating experiments with either power, and without any change of position or adjustment of the apparatus, it evidently deserved the most minute examination.

18. There was little hope (par. 6.) that cushions could be dispensed with. They were therefore added; and it was then seen, that the electrified conductors were supplied by the difference between the action of the cushion which had the advantage of the silk and that which had not; so that the naked face of the cylinder was always in a strong electric state. Methods were used for taking off the pressure of the receiving cushion; but the extremity of the silk, by the construction, not being immediately under that cushion, gave out large flashes of electricity with the power that was used. Neither did it appear practicable to present a row of points or other apparatus to intercept the electricity which flew round the cylinder; because such an addition would have materially diminished the intensity of the conductor, which in the usual way was such as to flash into the air from rounded extremities of four inches diameter, and made an inch and half ball become luminous and blow like a point. But the greatest inconvenience was, that the two states with the backward and forward turn were seldom equal; because the disposition of the amal-

gam

gam on the silk, produced by applying the leather to the cylinder in one direction of turning, was the reverse of what must take place when the contrary operation was performed.

Notwithstanding all this, as the intensity with the two cushions was such as most operators would have called strong, the method may be of use, and I still mean to make more experiments when I get possession of a very large machine which is now in hand.

19. The more immediate advantage of this discovery is, that it suggested the idea of two fixed cushions with a moveable silk flap and rubber. Upon this principle, which is so simple and obvious, that it is wonderful it should have been so long overlooked, I have constructed a machine with one conductor, in which the two opposite and equal states are produced by the simple process of loosening the leather rubber, and letting it pass round with the cylinder (to which it adheres) until it arrives at the opposite side, where it is again fastened. A wish to avoid prolixity prevents my describing the mechanism by which it is let go, and fastened in an instant, at the same time that the cushion is made either to press or is withdrawn, as occasion requires.

20. Although the foregoing series of experiments naturally lead us to consider the silk as the chief agent in excitation; yet as this business was originally performed by a cushion only, it becomes an object of enquiry to determine what happens in this case.

21. The great BECCARIA \* inferred, that in a simple cushion, the line of fire, which is seen at the extremity of contact from which the surface of the glass recedes, consists of returning electricity; and Dr. NOOTH grounded his happy

\* Philosophical Transactions, Vol. LVI. p. 117.

invention of the silk flap upon the same supposition. The former asserts, that the lines of light both at the entering and departing parts of the surface are absolutely similar; and thence infers, that the cushion receives on the one side, as it certainly does on the other. I find, however, that the fact is directly contrary to this assertion; and that the opposite inference ought to be made, as far as this indication can be reckoned conclusive: for the entering surface exhibits many luminous perpendiculars to the cushion, and the departing surface exhibits a neat uniform line of light. This circumstance, together with the consideration that the line of light behind the silk in par. 8. could not consist of returning electricity, shewed the necessity of farther examination. I therefore applied the edge of the hand as a rubber, and by occasionally bringing forward the palm, I varied the quantity of electricity which passed near the departing surface. When this was the greatest, the sparks at the electrometer were the most numerous. But, as the experiment was liable to the objection that the rubbing surface was variable, I pasted a piece of leather upon a thin flat piece of wood, then amalgamed its whole surface, and cut its extremity off in a neat right line close to the wood. This being applied by the constant action of a spring against the cylinder, produced a weak excitation, and the line where the contact of the cylinder and leather ceased (as abruptly as possible) exhibited a very narrow fringe of light. Another piece of wood was prepared of the same width as the rubber, but one quarter of an inch thick, with its edges rounded, and its whole surface covered with tin-foil. This was laid on the back of the rubber, and was there held by a small spring, in such a manner as that it could be slid onward, so as occasionally to project beyond the rubber, and cover the departing and excited surface

of

of the cylinder, without touching it. The sparks at the electrometer were four times as numerous when this metallic piece was thus projected; but no electricity was observed to pass between it and the cylinder. The metallic piece was then held in the hand to regulate its distance from the glass; and it was found, that the sparks at the electrometer increased in number as it was brought nearer, until light appeared between the metal and the cylinder, at which time they became fewer the nearer it was brought, and at last ceased when it was in contact.

The following conclusions appear to be deducible from these experiments. 1. The line of light on a cylinder departing from a simple cushion consists of returning electricity; 2. the projecting part of the cushion compensates the electricity upon the cylinder, and by diminishing its intensity prevents its striking back in such large quantities as it would otherwise do; 3. that if there were no such compensation, very little of the excited electricity would be carried off; and, 4. that the compensation is diminished, or the intensity increased, in an higher ratio than that of the distance of the compensating substance; because if it were not, the electricity which has been carried off from an indefinitely small distance, would never fly back from a greater distance and form the edge of light.

22. I hope the considerable intensity I shall speak of will be an apology for describing the manner in which I produce it. I wish the theory of this very obscure process were better known; but no conjecture of mine is worth mentioning. The method is as follows:

Clean the cylinder, and wipe the silk.

Grease the cylinder by turning it against a greased leather till it is uniformly obscured. I use the tallow of a candle.

Turn

Turn the cylinder till the silk flap has wiped off so much of the grease as to render it semi-transparent.

Put some amalgam on a piece of leather, and spread it well so that it may be uniformly bright. Apply this against the turning cylinder. The friction will immediately increase, and the leather must not be removed until it ceases to become greater.

Remove the leather, and the action of the machine will be very strong.

My rubber, as before observed, consists of the silk flap pasted to a leather, and the cushion is pressed against the silk by a slender spiral spring in the middle of its back. The cushion is loosely retained in a groove, and rests against the spring only, in such a manner that by a sort of libration upon it as a fulcrum, it adapts itself to all the irregularities of the cylinder, and never fails to touch in its whole length. There is no adjustment to vary the pressure, because the pressure cannot be too small when the excitation is properly made. Indeed, the actual withdrawing of the cushion to the distance of one-tenth of an inch from the silk, as in par. 2. will not materially affect a good excitation.

The amalgam is that of Dr. HIGGINS, composed of zinc and mercury. If a little mercury be added to melted zinc, it renders it easily pulverable, and more mercury may be added to the powder to make a very soft amalgam. It is apt to crystallize by repose, which seems in some measure to be prevented by triturating it with a small proportion of grease: and it is always of advantage to triturate it before using.

A very strong excitation may be produced by applying the amalgamed leather to a clean cylinder with a clean silk. But



it soon goes off, and is not so strong as the foregoing, which lasts several days.

23. To give some distinctive criterions by which other electricians may determine whether the intensity they produce exceeds or falls short of that which this method affords, I shall mention a few facts.

With a cylinder 7 inches diameter and cushion 8 inches long, three brushes at a time constantly flew out of a three-inch ball in a succession too quick to be counted, and a ball of  $1\frac{1}{2}$  inch diameter was rendered luminous, and produced a strong wind like a point. A nine-inch cylinder with an eight-inch cushion occasioned frequent flashes from the round end of a conductor 4 inches diameter: with a ball of  $2\frac{1}{2}$  inches diameter the flashes ceased now and then, and it began to appear luminous: a ball of  $1\frac{1}{2}$  inch diameter first gave the usual flashes; then, by quicker turning, it became luminous with a bright speck moving about on its surface, while a constant stream of air rushed from it; and, lastly, when the intensity was greatest, brushes, of a different kind from the former, appeared. These were less luminous, but better defined in the branches; many started out at once with a hoarse sound. They were reddish at the stem, sooner divided, and were greenish at the point next the ball, which was brass. A ball of  $\frac{4}{10}$  inch diameter was surrounded by a steady faint light, enveloping its exterior hemisphere, and sometimes a flash struck out at top. When the excitation was strongest a few flashes struck out sideways. The horizontal diameter of the light was longest, and might measure one inch, the stem of the ball being vertical.

This last phenomenon is similar to a natural event related by M. LOAMMI BALDWIN\*, who raised an electrical kite in

\* Memoirs of the American Academy, Vol. I. p. 257.

July, 1771, during the approach of a severe thunder-storm, and observed himself to be surrounded by a rare medium of fire, which, as the cloud rose nearer the zenith, and the kite rose higher, continued to extend itself with some gentle faint flashes. Mr. BALDWIN felt no other effect than a general weakness in his joints and limbs, and a kind of listless feeling; all which he observes might possibly be the effect of surprize, though it was sufficient to discourage him from persisting in any farther attempt at that time. He therefore drew in his kite, and retired to a shop till the storm was over, and then went to his house, where he found his parents and friends much more surprized than he had been himself; who, after expressing their astonishment, informed him, that he appeared to them (during the time he was raising the kite) to be in the midst of a large bright flame of fire, attended with flashings; and that they expected every moment to see him fall a sacrifice to the flame. The same was observed by some of his neighbours, who lived near the place where he stood.

This fact is similar to another observed by M. DE SAUSSURE on the Alps, and both are referable to my luminous ball with the second kind of brush. The cloud must have been negative.

With a 12-inch cylinder, and rubber of  $7\frac{1}{2}$  inches, a five-inch ball gave frequent flashes, upwards of 14 inches long, and sometimes a six-inch ball would flash. I do not mention the long spark, because I was not provided with a favourable apparatus for the two larger cylinders. The 7-inch cylinder affords a spark of  $10\frac{3}{4}$  inches at best. The 9-inch cylinder, not having its conductor insulated on a support sufficiently high, afforded flashes to the table which was 14 inches distant. And the 12-inch cylinder, being mounted only as a model or trial for constructing a larger apparatus, is defective in several respects

respects which I have not thought fit to alter. When the five-inch ball gives flashes, the cylinder is enveloped on all sides with fire which rushes from the receiving part of the conductor. I never use points, but in a simple machine bring the conductor almost in contact with the cylinder. In this apparatus that cushion to which the rubber is not applied serves that purpose.

24. These marks exhibit the intensity as deduced from simple electrifying. I will now mention the rate of charging, which was nearly the same in all the three cylinders.

A large jar of 350 square inches, or near  $2\frac{1}{2}$  square feet, with an uncoated varnished rim, of more than four inches in height, was made to explode spontaneously over the rim. The jar, when broken, proved to be 0.082 inches thick on an average; and the number of square feet of the surface of the cylinder which was rubbed, to produce the charge of one foot, was, when least, 18.03, and when most, with good excitation, 19.34. The great machine at Harlem charges \* a single jar of one foot square by the friction of 66.6 square feet, and charges its battery of 225 square feet at the rate of 94.8 square feet rubbed for each foot. The intensity of electricity on the surface of the glass is therefore considerably less than one-fourth of that here spoken of; but if we take the most favourable number 66.6 at the commencement of turning, and halve it on account of the unavoidable imperfection of a plate machine (as shewn in par. 14.), it will be found, that the management applied to that machine would cause a cylinder to charge one square foot by the friction of  $33\frac{1}{3}$  square feet. It must be observed, however, that M. VAN MARUM's own machine, con-

\* To explode from the central wire, which, from some trials, I find to require less force than from coating to coating at equal distances.

fisting of two plates, 33 inches diameter, has only half the intensity, though he reckons it a very good one. This machine is about equal in absolute power to my 9-inch cylinder, with its short rubber; but it is near thirty times as dear in price. In all these deductions I omit the computations, for the sake of brevity, and because they are easily made. The *data* are found in the description of the Teylerian machine, and its continuation published at Harlem in the years 1785 and 1787.

I shall here take the liberty of observing, that the action of the cylinder, by a simple cushion or the hand, which excited the astonishment of all Europe, in the memory of our contemporaries, was first improved by the addition of a leathern flap; then by moistening the rubber; afterwards by applying the amalgam; and, lastly, by the addition of a silk flap. Now, I find, by experiment, that we at present obtain upwards of forty times the intensity which the bare hand produces; and consequently that, since eighteen times our present intensity will equal the utmost we can now condense on strong glass even in the form of a charge; we have a less step to take before we arrive at that amazing power, than our immediate predecessors have already made.

My 9-inch cylinder, when broken, proved to be  $\frac{1}{21}$  of an inch thick.

## SECT. II. *Upon the luminous Appearances of Electricity and the Action of Points.*

25. Some of the luminous appearances, with balls in the positive state, have been slightly noticed as criterions of intensity. I shall here add, that the escape of negative electricity from a ball is attended with the appearance of strait sharp sparks with a hoarse

hoarse or chirping noise. When the ball was less than two inches in diameter, it was usually covered with short flames of this kind, which were very numerous.

26. When two equal balls were presented to each other, and one of them was rendered strongly positive, while the other remained in connection with the earth, the positive brush or ramified spark was seen to pass from the electrified ball: when the other ball was electrified negatively, and the ball, which before had been positive, was connected with the ground, the electricity (passing the same way according to FRANKLIN) exhibited the negative flame, or dense straight and more luminous spark, from the negative ball; and when the one ball was electrified *plus* and the other *minus*, the signs of both electricities appeared. If the interval was not too great, the long zig-zag spark of the *plus* ball struck to the strait flame of the *minus* ball, usually at the distance of about one-third of the length of the latter from its point, rendering the other two-thirds very bright. Sometimes, however, the positive spark struck the ball at a distance from the negative flame. These effects are represented in Plate IV. fig. 1, 2 and 3.

27. Two conductors of three-quarters of an inch diameter, with spherical ends of the same diameter, were laid parallel to each other, at the distance of about two inches, in such a manner as that the ends pointed in opposite directions, and were six or eight inches asunder. These, which may be distinguished by the letters P and M, were successively electrified as the balls were in the last paragraph. When one conductor P was positive, fig. 5. it exhibited the spark of that electricity at its extremity, and struck the side of the other conductor M. When the last-mentioned conductor M was electrified negatively, fig. 4. the former being in its turn connected with the earth,

the sparks ceased to strike as before, and the extremity of the electrified conductor M exhibited negative signs, and struck the side of the other conductor. And when one conductor was electrified *plus* and the other *minus*, fig. 6. both signs appeared at the same time, and continual streams of electricity passed between the extremities of each conductor to the side of the other conductor opposed to it. In each of these three cases, the current of electricity, on the hypothesis of a single fluid, passed the same way.

28. In drawing the long spark from a ball of four inches diameter, I found it of some consequence that the stem should not be too short, because the vicinity of the large prime conductor altered the disposition of the electricity to escape; I therefore made a set of experiments, the result of which shewed, that the disposition of balls to receive or emit electricity is greatest when they stand remote from other surfaces in the same state; and that between this greatest disposition in any ball, whatever may be its diameter, every possible less degree may be obtained by withdrawing the ball towards the broader or less convex surface out of which its stem projects, until at length the ball, being wholly depressed beneath that surface, loses the disposition entirely. From these experiments it follows, that a variety of balls is unnecessary in electricity; because any small ball, if near the prime conductor, will be equivalent to a larger ball whose stem is longer.

29. From comparing some experiments, made by myself many years ago, with the present set, I considered a point as a ball of an indefinitely small diameter, and constructed an instrument consisting of a brass ball of six inches diameter, through the axis of which a stem, carrying a fine point, was screwed. When this stem is fixed in the prime conductor, if  
the

the ball be moved on its axis in either direction, it causes the fine point either to protrude through a small hole in its external surface, or to withdraw itself; because by this means the ball runs along the stem. The disposition of the point to transmit electricity may thus be made equal to that of any ball whatever, from the minutest size to the diameter of six inches. See fig. 7. let. A.

30. The action of pointed bodies has been a subject of discussion ever since it was first discovered, and is not yet well explained. To those who ascribe this effect to the figure of electric atmospheres, and their disposition to fly off, it may be answered, that they ought first to prove their existence, and then shew why the cause which accumulated them does not prevent their escape; not to mention the difficulty of explaining the nature of negative atmospheres. If these be supposed to consist of electrified air, it will not be easy to shew why a current of air passing near a prime conductor does not destroy its effects. The opinion, supported by the celebrated VOLTA and others, that a point is the coating to an infinitely small plate of air, does not appear better founded: for such a plate must be broken through at a greater distance only because higher charged; whence it would follow, that points should not act but at high intensities. I must likewise take notice, as a proof that the charge has little to do here, that if a ball be presented to the prime conductor, at the same time that a point proceeds from the opposite side of the ball, the electricity will pass by the point, though it is obliged to go round the ball for that purpose; but it can hardly be doubted, that whatever charge obtains in this case is on the surface of the ball next the conductor, and not on the remote side to which the electricity directs its course.

31. ACHARD'S experiments with a number of pointed cones, screwed in a plate of metal, and likewise the pointed apparatus described (par. 29.) shew that the effect of points depends on the remoteness of their extremities from the other parts of the conductor. This leads to the following general law.

*In any electrified conductor the transition or escape of electricity will be made chiefly from that part of the surface which is the most remote from the natural state.*

Thus in the apparatus of the ball and stem, the point having a communication with the rest of the whole conductor, constantly possesses the same intensity; but the influence of the surrounding surface of the ball diminishes its capacity. This diminution is less the farther the ball is withdrawn, and consequently the point will really possess more electricity, and be more disposed to give it out when it is prominent than when depressed. The same explanation serves for negative electricity.

32. The effect of a positive surface appears to extend farther than that of a negative: for the point acts like a ball when considerably more prominent if it be positive than it will if negative. This property was used by me some years ago for the construction of an instrument to distinguish the two electricities\*.

For the sake of conciseness I pass over many facts which have presented themselves in the course of my experiments on the two electricities, and content myself with observing, that there is scarcely any experiment made with the positive power which will not afford a result worthy of notice, if repeated with the negative.

\* Introduction to Natural Philosophy, Vol. II. p. 320.



33. When we consider that our machines can cause a ball of an inch and half diameter to act like a point, and that our apparatus makes a point act like a ball; if at the same time we remark the small elevation of our conductors for lightning above the extended surface of the ground, and the small size of the balls proposed by some to be used as terminations; the dispute, which was so much agitated respecting them, will perhaps be found to relate to a very minute circumstance, among the many which govern the great operations of nature. It does not seem probable, that any conductor would act silently if the main course of the electricity of a negative cloud were to pass through it, and many would probably receive the stroke from a positive cloud. It does not, however, follow from this, that they might not conduct it with safety.

### SECT. III. *Of compensated Electricity.*

34. It is unnecessary to insist upon what is called the equilibrium of an electrical charge, because Dr. FRANKLIN has admirably explained it according to his hypothesis. But there is another important particular, which has been almost entirely overlooked, namely, the uncompensated electricity which is as essential to the charge as that which is in equilibrium. Whenever a jar is charged, the greatest part of the electricity becomes latent on account of the compensation; but there is a certain proportion which remains on the insulated side, and exerts its force to prevent the electricity from returning to the outer surface. In moderate intensities, this will explode, and carry the charge with it, to distances which are in proportion to the quantity of the charge itself; but in greater intensities the distances greatly exceed that proportion. With glasses of different thicknesses, this intensity,

as measured by the explosive spark, is as the thickness, when the charges are equal, as Mr. CAVENDISH has determined, and I find likewise by experiments with thin substances; but when the thicknesses are greater, it increases in a higher proportion, as is found by the explosion which takes place between the electrophore and its plate, as well as by other experiments.

35. This uncompensated part of the charge (which is commonly in proportion to the quantity of latent or compensated electricity, or to the distance at which it exerts its action) was found to be greatly increased when a series of jars were made to charge each other. If a jar be insulated and made to explode by LANE'S electrometer at a determinate number of turns; and another jar be then connected with its external coating so as to become charged by that means, the explosion, from the outside of the last to the inside of the first, will take place at the electrometer (unaltered) with much fewer turns. Or if the electrometer be altered till the explosion takes place at the original number, the distance will be much greater than before. Hence we see, that the intensity of the uncompensated part must be greater when there is a greater charge to be maintained, whether it be on one surface only, or on two surfaces successively connected. I have not yet made the experiments necessary to ascertain the law of this last action.

36. It is evident, that the breaking of jars is not effected by any attraction between the electricities which form the charge, but by this necessary surplus: for thicker glasses require much less electricity to produce an intensity which breaks them than thinner do; and I found a piece of Muscovy talc, one hundredth of an inch thick, to bear a charge consisting of ten times the quantity of electricity which was sufficient to have charged an equal surface of common glass so as to break it.

But

But the intensity of the very dense charge on the talc was so low as to afford an explosion of no more than about one-tenth of an inch, while that of the glass jar it was compared with exploded through about five inches.

The perforation of glass by the long spark, or by the spark through oil or cement, seems to depend on the very great intensity of the electricity which has not time to diffuse itself, but charges a minute part of the surface very high.

37. Muscovy talc \* being a very perfect non-conductor, and capable of being divided into plates of less thickness than one two-hundredth part of an inch, I made many experiments with it, which are too numerous to enter into this Paper. In consequence of its great capacity it gives very strong shocks. Contrary to the assertion of BECCARIA, I found that its laminæ are naturally in strong opposite states of electricity, and flash to each other when torn asunder in the dark. A large piece being split in two, the parts were found to be in opposite states. The greatest care was taken in these experiments to avoid friction, and to use such pieces as had never been excited, nor brought near the machine.

38. The most plausible objection against the probability of danger from the returning stroke of the Earl of STANHOPE is, that the quantity of electricity in an animal is too small to produce any mischievous effect. This the noble author has answered by remarking, that the quantity has not been shewn to be small †. My experiments with talc shewing that it naturally possesses much electricity, led me to investigate the quantity which a man may contain. I melted sealing wax upon BENNET's electrometer with a burning glass, and found it pro-

\* I am not certain whether HENLEY or BECCARIA first used this substance; but little attention was paid to it by either. † Phil. Trans. Vol. LXXVII. p. 143.

duced no electricity either in heating or cooling. I also placed a piece of red-hot glass upon the same instrument, and it cooled without affording electric signs. These experiments shewed, that the natural quantity of electricity is the same in these bodies, whether they be in the conducting or non-conducting state; and consequently, if it can be proved, that an electric contains a large quantity of electricity, the inference may be fairly extended to non-electrics. And it will not be disputed, upon any hypothesis, but that a non-conductor, or its coating, contains as much of what we call electricity as can be driven out of it in the act of charging. Two square inches of talc, of the thickness of 0,011 inch, were repeatedly charged and made to explode over the uncoated part, by each turn of a seven-inch cylinder. The intensity of the excitation was such, that a conductor, of three feet long, and seven inches diameter, gave a dense spark of 9 inches long at each turn. Now, in round numbers 45 such plates of talc, laid upon each other, would have formed a solid inch of matter; and from this, if fitted up as a BECCARIA'S battery, we could with our machine drive out electricity enough simply to charge a conductor 45 times as long (neglecting the ends); that is to say, we find that one solid inch of talc contains electricity enough to charge a conductor of 7 inches diameter, and 135 feet long, so high as to give a nine-inch spark at least; but how much more it contains we know not.

If it be here objected, that the talc does nothing more than separate the coatings, we may make use of gold leaf for our coating; which substance being (as I find by weight and measurement) no more than  $\frac{1}{282000}$  of an inch thick, would increase the result near three thousand times.

Without

Without referring to the intense electricity of a cloud, or the bulk of a man, it may be observed, that such a spark would be very painful. But to pursue our computation. The cylinder charged a square foot of glass, of about 0.08 thick, in 15 turns so as to explode over a rim above four inches high. Fifteen of the pieces of talc would therefore possess as much electricity as makes the charge of a jar of one foot square, and the 45 pieces or solid inch would contain enough to charge three square feet. If we suppose the bulk of a man to be only 3 solid feet or 5184 solid inches, the natural electricity of this mass, as deduced from the foregoing facts, will be equal to the charge of a battery of upwards of 15,000 square feet.

I beg leave to observe, in concluding this Paper, that I have been very careful in repeating the experiments with many precautions which the experienced in this branch of natural philosophy will perceive the necessity of; though, in order to keep this communication within proper limits, I have here avoided a minute description of them. With the same view I have likewise forborn to speak either of theory, or of a number of other experimental researches I have made during the course of this enquiry. Dates are entirely omitted from a conviction that the priority of accidental discovery is not worth contending for, and that no disputes ever arise about that general tenor of conduct in the cultivation of science, upon which the rational part of mankind ground their approbation or censure.

New North-street,

May 14, 1789.

P. S. Since the above was written, the *Journal de Physique* for April, 1789, has arrived. It contains an excellent Paper of M. VAN MARUM, giving an account of some very considerable amendments of the rubber, and of the manner of applying it to plate machines. The chief improvement consists in fixing the silk to the posterior part of the rubber, so that it covers the whole face, and has the amalgam applied upon it. I cannot, however, avoid expressing my surprize, that this improvement, which has been in common use in England for upwards of twelve years past, should now be offered as a discovery by so experienced a philosopher. With his new rubbers M. VAN MARUM excites his plates of 33 inches diameter so strongly as to produce nearly two-thirds of the former effect of the Teylerian machine, though the rubbed surfaces of these machines are now as 691 to 2409. This power would charge the single jar by the friction of 28.6 square feet, or the battery by rubbing 36.2 feet, instead of the numbers 66.6 and 94.8, as given in par. 24. This is a vast acquisition of intensity; but still little more than half that of the surface of a cylinder, as mentioned in the same paragraph. But if par. 14. be admitted to prove that plate machines gain nothing by the friction of the back surfaces, it will follow, that M. VAN MARUM'S management, if applied to a cylinder, would do better than mine.



Fig. 1



Fig. 2

Fig. 3



Fig. 5





Fig. 4

Fig. 6




Fig. 7





XXIII. *Experiments on the Transmission of the Vapour of Acids through an hot earthen Tube, and further Observations relating to Phlogiston.* By the Rev. Joseph Priestley, LL.D. F. R. S.

Read July 2, 1789.

**I**N my late experiments on the *phlogistication of spirit of nitre by heat* it appeared, that when pure air was expelled from what is called dephlogisticated spirit of nitre, the remainder was left phlogisticated. This I find abundantly confirmed by repeating the experiments in a different manner, and on a larger scale; and I have applied the same process to other acids and liquors of a different kind. From these it will appear, that oil of vitriol and spirit of nitre, in their most dephlogisticated state, consist of a proper saturation of the acids with phlogiston, so that what we have called the *phlogistication* of them, ought rather to have been called their *super-phlogistication*.

I began with treating a quantity of oil of vitriol as I had done the spirit of nitre, *viz.* exposing it to heat in a glass tube, hermetically sealed, and nearly exhausted; and the result was similar to that of the experiment with the nitrous acid, with respect to the expulsion of air from it, though the phlogistication not appearing by any change of *colour*, I did not in this method ascertain that circumstance. The particulars were as follows.

After the acid had been made to boil some time, a dense white vapour appeared in quick motion at a distance above the acid,

acid, and though, on withdrawing the fire, that vapour disappeared, it instantly re-appeared on renewing the heat. When the tube was cool, I opened it under water, and a quantity of air rushed out, though the acid had been made to boil violently while it was closing, so that there could not have been much air in the tube. This air, which must therefore have been generated in the tube, was a little worse than common air, being of the standard of 1.12 when the latter was 1.04. I repeated the experiment several times, and always with the same result.

That this air should be worse than common air, I cannot well explain. But in my former experiments it appeared that vitriolic acid air injures common air; and that in proportion as pure air is expelled from this acid, the remainder becomes phlogisticated, or charged with vitriolic acid air, clearly appeared in the following experiment.

Making a quantity of oil of vitriol boil in a glass retort, and making the vapour pass through a red-hot earthen tube, glazed inside and out, and filled with pieces of broken tubes, I collected the liquor that distilled over, and found it to be the same thing with water impregnated with vitriolic acid air. The smell of it was exceedingly pungent, and it was evident, that more of this air had escaped than could be retained by that quantity of water. The oil of vitriol used in this process was 1 oz. 9 dw. 18 gr. and the liquor collected was 6 dw. 12 gr. When I collected the air that was produced in this manner, which I did not do at this time, it appeared to be very pure, about the standard of 0.3 with two equal measures of nitrous air.

At another time, expending 1 oz. 11 dw. 18 gr. of oil of vitriol, of the specific gravity of 1856 (that of water being 1000),

1000), I collected 19 dw. 6 gr. of the volatile acid, of the specific gravity of 1340, and 130 oz. measures of dephlogisticated air of the purest kind, *viz.* of the standard of 0.15.

It is easy in this manner to collect a great quantity of dephlogisticated air; but the principal objection to the process is, that after using a few times, the earthen tubes become tender, and too easily break, especially in heating or cooling. It is also difficult to lute the retort containing the acid and the earthen tube. The air produced in this manner is filled with the densest white cloud imaginable.

Going through the same process with spirit of nitre, the result was in all respects similar, but much more striking, the production of both dephlogisticated air and phlogisticated acid vapour being prodigiously quicker, and more abundant. Expending 5 oz. 8 dw. 6 gr. of spirit of nitre, I collected 600 oz. measures of very pure dephlogisticated air, being of the standard of 0.2. I also collected 1 oz. 7 dw. 14 gr. of a greenish acid of nitre, which emitted copious red fumes. All the apparatus beyond the hot tube was filled with the densest red vapour, and the water of the trough in which the air was received was so much impregnated with it, that the smell was very strong; and it spontaneously yielded nitrous air several days, just as water does when impregnated with nitrous vapour. Perceiving the emission of air from the water, after it had stood some time, I filled a jar containing 30 oz. measures with it, and without any heat it yielded two oz. measures of the strongest nitrous air.

Taking the specific gravity of the acid before and after this distillation, the former was to the latter as 1471 to 1182. When the weight of the air produced in this experiment, and that of the liquor distilled, is compared with that of the acid

before

before distillation, it will appear, that there must have been a great loss of acid vapour, which was either retained in the water of the trough, or escaped through it.

I do not see that these experiments can be explained, but on the supposition that the most dephlogisticated oil of vitriol and spirit of nitre are, in a proper sense, saturated with phlogiston; and that when part of the acidifying principle is expelled in the form of the air, the remainder is supersaturated with it.

To try whether the acid, thus supersaturated with phlogiston, was convertible into pure air by this process, I heated the liquor collected after the distillation of the oil of vitriol, that is, water impregnated with vitriolic acid air, and made the vapour pass through the hot tube, but no air came from it; and when collected a second time, it was not at all different from what it had been before. The specific gravity was also the same.

It is evident, however, though this process does not shew it, that the volatile vitriolic acid contains the proper element of dephlogisticated air; since by melting iron in vitriolic acid air, a quantity of fixed air (which is composed of inflammable and dephlogisticated air) is produced. Melting iron in 9 oz. measures of vitriolic acid air, it was reduced to 0.3 oz. measures, and of this 0.17 oz. measures was fixed air. I repeated the experiment with the same result, and putting the residuums together found the air to be inflammable.

But the result was something different when I sent through the hot tube the liquor that I had collected in the process with spirit of nitre. No air, however, was produced at the first, nothing appearing besides a *red vapour* that was wholly absorbed by water, or escaped through it into the atmosphere; but towards the end of the process I collected 10 oz. measures of dephlo-

dephlogisticated air. The quantity of the liquor expended was about 2 oz. measures. It may, however, be presumed, that this small quantity of air came from some of the acid which escaped the action of the fire in the former process. Indeed its coming at the last only may be considered as a proof of this, as all the more volatile acid, which came over first, yielded no air.

I submitted a quantity of *spirit of salt* to both these processes, *viz.* exposing it to a boiling heat in glass tubes, hermetically sealed, and making the vapour pass through a red hot earthen tube, but no air was produced in either case. In the former case, the water rushed into, and completely filled, the tube, when it was opened under water; and in the other process the liquor distilled was precisely of the same specific gravity, and, no doubt, in all other respects, the same as before distillation; but the acid that remained in the retort was of less specific gravity, in consequence of the acid vapour being expelled by the heat in the form of marine acid air, which appeared not to be affected by a red heat.

Though, in the process with spirit of salt, the result be different from that of those with oil of vitriol and spirit of nitre, yet there is an analogy among all these three acids in this respect, *viz.* that the marine and both the volatile acids of vitriol and nitre are made by impregnating water with the acid vapour, so that in its usual state it may be said to be phlogisticated as well as these.

It was evident that the water in the worm-tub was much more heated by the distillation of the spirit of salt than by that of the oil of vitriol, and especially that of the spirit of nitre; so that much of the heat by which it had been raised in vapour must, in the latter case, have been *latent* in the air that was

formed; whereas, in the other case, it was communicated to the water in the worm-tub.

In one of the processes with boiling spirit of salt, in a glass tube, hermetically sealed, I had the same white vapour dancing in the middle of the tube as in the experiment with the oil of vitriol; but this tube burst, and I never had the same appearance again, though I repeated the experiment several times for the sake of it.

The vapour of dephlogisticated marine acid, which M. BERTHOLLET discovered, and with which water may be impregnated as with fixed air, being made to pass through the hot earthen tube, became dephlogisticated air as in the following experiment.

Having poured a quantity of spirit of salt upon some manganese in a glass retort, I heated it as in the preceding experiments with a proper apparatus both for receiving the distilled liquor, and the air. I found seven-tenths of the air was fixed air, and the remainder very pure dephlogisticated. The quantity I could not measure on account of one of the junctures in the apparatus giving way; but I do not imagine that quite so much pure air could be got in this method as from the manganese itself in a direct process. The liquor received in this distillation resembled strong spirit of salt in which manganese had been put.

This process immediately succeeding that in which the glass tube, joining the earthen tube and worm-tub, was left full of black matter by the distillation of the alkaline liquor (which will be mentioned hereafter), the blackness presently vanished, and the tube became transparent as before. On this account, however, it is possible that I might receive less pure air than I should otherwise have done.

Distilled

Distilled vinegar submitted to this process yielded air two-thirds of which was fixed air, and the rest inflammable: expending 2 oz. 19 dw. 0 gr. of the acid, I got 1 oz. 19 dw. 0 gr. of a liquor which had a more pungent smell than it had before distillation. It had also some black matter in it, and some of the same remained at the bottom of the retort when the liquor was evaporated to dryness. The air I received was 90 oz. measures.

Alkaline air is converted into inflammable air in this process as well as by the electric spark, but by no means, I think, in so great a degree. I put 2 oz. 10 dw. 0 gr. of water pretty strongly impregnated with alkaline air into the retort, and heating it, sent the vapour through the hot tube; when I collected 2 oz. 3 dw. 0 gr. of liquor, which had a disagreeable empyreumatic smell, as well as that of a volatile alkali, and it was quite opaque with a *black matter*, which subsided to the bottom of the vessel. Also the tube through which the air and vapour had been conveyed was left quite black, as mentioned above. One of the junctures of the apparatus not having been air-tight, I did not collect all the air, but it came only at the beginning of the process, and before the tube became black, or any liquor was distilled, and it was all strongly inflammable.

I shall now recite a few experiments of a different kind from those that have been mentioned above, and more immediately relating to the doctrine of phlogiston.

It is said, by those who do not admit the doctrine of phlogiston, that the metals are simple substances, which, having a strong affinity to dephlogisticated air, imbibe it when they become calces, without parting with any thing. But that something is really parted with in the calcination (as they will call

it) of iron in dephlogisticated air, appears to me to be very evident, as well as in the process with steam.

That fixed air is found in the vessel in which iron is melted in dephlogisticated air, I observed before; but I never took much care to ascertain the quantity of it. This I have lately done in many instances, and in all of them find it to be much more considerable than can be accounted for, by supposing it to come from plumbago in the very small quantity of iron that I melted; so that it must necessarily have been formed by the phlogiston from the iron, and the pure air in the vessel, at the same time that the iron became finery cinder by imbibing water from the air; and I have shewn, that by far the greatest part of the weight of this air is water. The experiments were made with a very good burning lens, of sixteen inches diameter, with which Mr. PARKER has generously furnished me; and by means of it I can now make these experiments, which require a great degree of heat, with much more ease and certainty than I could do before.

In  $6\frac{1}{2}$  oz. measures of dephlogisticated air I melted turnings of malleable iron till there remained only  $1\frac{1}{3}$  oz. measure, and of this  $\frac{2}{3}$  oz. measure was fixed air. In 6 oz. measures of dephlogisticated air, of the standard of 0.2, I melted iron till it was reduced to two-thirds of an ounce-measure, of which one-half was fixed air, and the remainder completely phlogisticated. Again, I melted iron in  $7\frac{1}{2}$  oz. measures of dephlogisticated air, of the same purity with that in the last experiment, when it was reduced to  $1\frac{1}{3}$  oz. measure, and of this four-fifths was fixed air, and the remainder phlogisticated. In this case I carefully weighed the finery cinder that was formed in the process, and found it to be nine grains, so that the iron that had been melted (being about two-thirds of this weight) had



had been about six grains. I repeated the experiment with the same result.

When the dephlogisticated air is more impure, the quantity of fixed air will always be less in proportion. Thus, having melted iron in seven ounce measures of dephlogisticated air of the standard of 0.65, it was reduced to 1.6 oz. m.; and of this only one-third of an ounce measure was fixed air. This, however, is much more than can come from the plumbago in the iron; but as the production of this fixed air is by many ascribed to this plumbago, it may be worth while to shew by computation that it is impossible that it should have this origin. Both the quantity of plumbago in iron, and the quantity of fixed air in plumbago, are much too small for the purpose.

From half an ounce of the purest plumbago, I first got, in a coated glass retort, 13 ounce-measures of air, of which only three ounce measures were fixed air, the rest being inflammable; then putting it into an earthen tube, I kept it some hours in as great a heat as I could produce, and got 22 oz. m. more; and of this also only three were fixed, and the rest inflammable, and the last portion was wholly so.

But instead of supposing the fixed air that I got to be that which was expelled from the plumbago in the iron, I will suppose that even the whole of this plumbago afforded only one of the elements of the fixed air, *viz.* phlogiston, or that which the French chemists call *carbone*; and that this principle, by its union with the dephlogisticated air in the vessel, forms the fixed air, yet on this most unfavourable and improbable supposition the quantity will be found to be insufficient.

If 100 gr. of iron contain, according to M. BERGMAN, 0.12 gr. of plumbago, 7 gr. (which is the most that in any of the preceding processes I converted into finery cinder) would

contain only 0.0084 gr. of plumbago; and if we suppose with Mr. KIRWAN, that an hundred cubic inches of fixed air contains 8.14 gr. of phlogiston, the fixed air produced in one of the above-mentioned processes (*viz.* four-fifths of an ounce-measure) would contain .032 gr. of phlogiston, which is above three times more than the plumbago in the iron could furnish. It is evident, therefore, that the quantity of fixed air that I found must have been formed by phlogiston from the iron uniting with the dephlogisticated air in the vessel.

If, as I have inferred, from burning charcoal of copper in dephlogisticated air (see Experiment, Vol. VI. p. 272.) fixed air consists of 3.45 parts of dephlogisticated air and 1.5 of phlogiston, it will be found, that four-fifths of an ounce measure of fixed air will contain 0.21 gr. of phlogiston, which is much more than on the supposition of Mr. KIRWAN.

Another argument against the antiphlogistic doctrine may be drawn from an experiment which I made upon Prussian blue; if the small quantity of fixed air, that may be expelled from it by heat, be compared with the much greater quantity which is produced when heated in dephlogisticated air.

Prussian blue is generally said to be a calx of iron super-saturated with phlogiston, though of late it has been said by some that it has acquired something that is of the nature of an *acid*. From my experiments upon it, with a burning lens in dephlogisticated air, I should infer, that the former hypothesis is true, except that the substance contains some fixed air, which is no doubt an acid; for much of the dephlogisticated air disappears, just as in the preceding similar process with iron.

I threw the focus of the burning lens upon 2 dw. 5 gr. of Prussian blue in a vessel of dephlogisticated air, of the standard of 0.53, till all the colour was discharged. Being then weighed,

weighed, it was 1 dw. 2 gr. In this process  $7\frac{1}{4}$  oz. of fixed air had been produced, and what remained of the air was of the standard of 0.94. Heating the brown powder to which the Prussian blue was reduced in this experiment in inflammable air, it imbibed  $8\frac{1}{2}$  oz. m. of it, and became of a black colour; but it was neither attracted by the magnet, nor was it soluble in oil of vitriol and water, as I had expected it would have been.

Again, I heated Prussian blue in dephlogisticated air, of the standard of 0.2, without producing any sensible increase of its bulk, when I found three ounce measures of it to be fixed air, and the standard of the residuum, with two measures of nitrous air, was 1.35. The substance had lost eleven grains, the greatest part of which was evidently water.

To determine what quantity of fixed air Prussian blue would yield by mere heat, I put half an ounce of it into an earthen tube, and got from it 56 oz. m. of air, of which 16 oz. m. were fixed air, in the proportion of one-third in the first portion, and one-fourth in the last. The remainder was inflammable. There remained 5 dw. 20 gr. of a black powder, with a very little of it (probably the surface) brown.

Comparing these experiments, it will appear, that the fixed air procured by means of Prussian blue and dephlogisticated air must have been formed by phlogiston from the Prussian blue and the dephlogisticated air in the vessel: for if 240 gr. of this substance yield 16 oz. measures of fixed air, ten grains of it (which is more than was used in the experiment) would have yielded only 0.6 oz. m. Nor is it possible to account for the disappearing of so much dephlogisticated air, but upon the supposition of its being employed in forming this fixed air.



XXIV. *On the Production of nitrous Acid and nitrous Air.* By  
*the Rev. Isaac Milner, B. D. F. R. S. and President of*  
*Queen's College, Cambridge.*

Read July 2, 1789.

1. **I**T has been known for some time, that a relation subsists between nitrous acid and volatile alkali. The latter has frequently been produced by help of the former; but I do not recollect that, in any instance, the volatile alkali has been proved to contribute to the formation of nitrous acid or nitrous air. Some cases, however, have occurred to me where this evidently happens; and they appear so new and extraordinary, that I cannot but think they deserve the attention of philosophical chemists. The history of the experiments I allude to is as follows.

2. As soon as I had heard of the production of inflammable air by the transmission of steam through red-hot iron tubes, I had the curiosity to try whether some other substances in the form of air or vapour might not, by a similar process, undergo material alterations. In particular, the nitrous acid seemed well to deserve a trial, both on account of the obscurity and difficulties attending the theory of its production, and also of its important and extensive usefulness in chemistry.

In the relation of my experiments on this head, it will be unnecessary to mention the exact *quantity* of acid or of air expended or generated, though I noted those quantities pretty accurately

accurately at the time; for the main point I have in view in this description, is to ascertain the *nature* of the *changes* which took place; and these do not depend upon the quantities of aerial fluids, but upon their properties. Besides, whoever shall repeat these experiments will find the relative *quantities* to vary very much, according to the manner of operating; and therefore, for the sake of brevity, I omit to mention them entirely.

3. I began with boiling a little strong nitrous acid in a small retort, the neck of which was closely luted to one end of a gun-barrel. The other end of it was immersed sometimes in water, and sometimes in quicksilver, and eighteen or twenty inches of the middle part was surrounded with burning charcoal in a proper furnace. In this manner the vapour and fumes of the boiling acid were transmitted through the red-hot tube, and the produce received at the end in the usual manner.

When the acid was made to boil violently, there passed over a considerable quantity of undecomposed red nitrous vapour, together with a mixture of nitrous and phlogisticated airs.

When the process was conducted more moderately, there was less nitrous vapour; and in the mixture of airs which was received in the glass vessels, there was a much greater proportion of phlogisticated air.

4. In order to increase the surface of the red-hot iron, and effect a more complete decomposition of the nitrous vapour, the gun-barrel was crammed full of iron filings. The experiments were repeated with great caution, and almost the whole of the produce was found to be phlogisticated air. It is however proper to mention, that, notwithstanding every possible care, still there will generally be in some degree an admixture

of nitrous air, and frequently of dephlogisticated nitrous air. But I am satisfied that if the iron tube were sufficiently long, so that a very large portion of it might be heated red-hot, all the air received in this manner from any quantity of nitrous acid slowly boiled would be found of that species called phlogisticated air.

5. These experiments seem altogether analogous to those of Dr. PRIESTLEY, in which nitrous air, by exposure to iron, is converted first into dephlogisticated nitrous air, and afterwards into phlogisticated air. The only difference seems to be, that in my experiments the effect is brought about suddenly; whereas in the method of exposition to iron much time is required. And farther, in my method of operating, it is very difficult to conduct the process so as to insure the production of that singular species of air called dephlogisticated nitrous air. If the acid boil very quick, the product is nearly all nitrous vapour and nitrous air. If it boil very slow, and a sufficient quantity of the iron tube be well heated, then the decomposition is almost complete, and little is received but phlogisticated air. In both cases, the progress of the conversion of nitrous acid to the state of phlogisticated air seems to be the same. First, nitrous air is formed, then dephlogisticated nitrous air, and lastly phlogisticated air. This, I say, seems to me to be the natural order of the conversion, though I do not deny, that in the rapid manner of operating with the red-hot iron tube some particles of nitrous acid or vapour may probably be *instantly* changed into phlogisticated air. And even allowing this to be the case, the fact may easily be explained, by supposing the successive approaches to phlogisticated air to be made in too small spaces of time to be observed; nor does it in the least invalidate the general conclusion, that nitrous  
air

air is nearer the state of phlogisticated air than nitrous acid or nitrous vapour; and that dephlogisticated nitrous air is still nearer. It is very difficult to decide with certainty what the changes are which the particles of the acid undergo in their passage through different parts of the hot tube.

From what has been said, the most common process will probably appear to be, that a particle of the acid in the form of vapour first generates nitrous air; that the parts of this are applied to fresh surfaces of hot iron, and suddenly changed into dephlogisticated nitrous air; which, lastly, is applied to still fresh surfaces of the tube or fragments of iron, and so converted into phlogisticated air. When these successive contacts with fresh surfaces of hot iron are not sufficiently numerous or exact, it is not unnatural to conclude, that some portion of air may escape not perfectly decomposed.

6. These considerations induced me to alter the process a little. Instead of boiling the acid in the retort, I put some thin pieces of copper into a phial, poured nitrous acid upon them, and forced the nitrous air, as it was generated, to pass through the red-hot tube. The event answered my expectation; the decomposition was effected in this way easier than in the former.

But before I made this experiment, I examined what would be the effect of mere heat upon nitrous air, as I had already learned from the experiments of others, that nitrous acid, forced in the form of steam through red-hot tubes of clay or glass, underwent the most important alterations.

What might be the effect of long continued exposure to a red heat I cannot say; but I was soon convinced, that nitrous air might be forced through a red-hot glass tube, without suffering any material change.

7. Lastly, I determined to try the effect of the gun-barrel upon dephlogisticated nitrous air, as from all that I had seen it seemed reasonable to expect, that this species of air would be the easiest reduced to the state of phlogisticated air. For this purpose, I diluted a saturated solution of copper in the nitrous acid, and put pieces of iron wire into it, and as the neck of the retort which contained the solution was luted to one end of the gun-barrel, the dephlogisticated nitrous air was exposed in its passage to the action of the red-hot tube, and also to the surfaces of the red-hot iron turnings which it contained. In this case, when the process is conducted with proper care, all the air which is received at the other end of the tube will be found phlogisticated.

8. When the air received at the end of the gun-barrel was in the last mentioned state, *viz.* perfectly phlogisticated, I have frequently observed a white fume issuing along with the air, and sometimes ascending through the water or mercury into the glass receivers. Upon examining this white fume, I soon perceived by the smell that it contained volatile alkali. I was much struck with the observation, and immediately recollected Dr. PRIESTLEY's relation of a similar production by exposing nitrous air to pieces of iron.

9. Most of the experiments hitherto related were made in the summer of 1786; in general they agree with those of Dr. PRIESTLEY; the changes and productions are much the same, and the only new circumstance is, as was observed at art. 5. The same effects are brought about *instantly* by the action of red-hot iron, which require much time by the method of simple exposure to cold iron.

For which reason, though it gave me much pleasure at the time to see such curious transmutations brought about in a few minutes,



minutes, yet it scarcely appeared worth while to trouble the Royal Society with a detail of the experiments; and I only presume to do it now, because the conjectures which I then formed have been sufficiently verified by future experiments.

The conjectures were as follow:

10. Almost immediately upon seeing the volatile alkali produced by means of nitrous acid and metals, I conceived the possibility of inverting the order of the process, and of producing nitrous acid or nitrous air by the decomposition of volatile alkali. I knew of no experiments wherein this had been done, or any thing like it; yet as volatile alkali was beyond all dispute produced in the method just described, and as the iron turnings and inside of the gun-barrel were left after the operation in a state of calcination, it seemed not unnatural to suppose, that by forcing volatile alkali through the red-hot calces of some of the metals, nitrous acid or nitrous air might be produced. Some of my friends, to whom I mentioned the idea, considered it as a random conjecture. However, I made a memorandum of it as a thing that deserved to be tried, though in fact I neglected for near two years actually to make the trial. It was some time in the month of March, 1788, that the calx of manganese on account of its very great infusibility, and its yielding abundance of dephlogisticated air, occurred to me as a very proper substance for the purpose. I immediately crammed a gun-barrel full of powdered manganese; and to one end of the tube I applied a small retort, containing the caustic volatile alkali. As soon as the manganese was heated red-hot, a lighted candle was placed under the retort, and the vapour of the boiling volatile alkali forced through the gun-barrel. Symptoms of nitrous fumes and of nitrous air soon discovered themselves,  
and

and by a little perseverance I was enabled to collect considerable quantities of air, which on trial proved highly nitrous. I have since frequently repeated this experiment, and have always in some degree succeeded. Much depends on the *kind* of manganese employed, much on the heat of the furnace, and much on the patience of the operator; as these are varied, there will be great variations of the products. A minute detail of all the particulars of my experiments seems unnecessary; but it may be proper to give a general account of the principal facts, and of the methods which were used to avoid erroneous conclusions.

11. In general I made use of clean gun-barrels with which no previous experiments had been made. The manganese was used in rough powder; for when it is too finely powdered, the tube is choaked, and the air cannot pass.

In some experiments I applied the vapour of the volatile alkali directly to the hot manganese. In others I suffered the manganese to remain a considerable time in a red heat before I made the volatile alkali, contained in the retort at the end of the tube, to boil; and by this means I informed myself of the nature of the airs which the manganese yielded *per se*.

In neither case could I ever perceive the least appearance of nitrous acid or nitrous air till the volatile alkali was used. Manganese, *per se*, gives airs of different kinds (but chiefly fixed and dephlogisticated airs) as soon as ever it is subjected to a considerable heat; but nothing nitrous comes from it, either on the first application of heat, or after it has been continued a long time; and I examined this point with great diligence. But soon after the volatile alkali begins to be applied, the jars in which the air is received will frequently turn slightly

slightly red, and this redness will increase on admitting atmospheric air.

The caustic alkali should be strong, and as far as I have observed the longer the process is continued, the stronger will be the nitrous air produced. At least this evidently appeared to be the case in several instances, where the operation was continued for a long time.

In most instances, on the very commencement of this process, a small jar of the air thus collected discovers by the *smell* a nitrous impregnation. But it sometimes happens, that several jars of air may be collected, and the admission of atmospheric air to them will not produce a sensibly red colour.

Here, however, there exists a cause of deception against which the operator ought to be on his guard, lest he should conclude that no nitrous air is formed, when in reality there is a considerable quantity. The volatile alkali, notwithstanding every precaution, will frequently pass over in great quantities undecomposed. If the receivers are filled with water, a great part of this will indeed be presently absorbed; but still some portions of it will mix with the nitrous air formed by the process. Upon admitting the atmospheric air, the nitrous air is decomposed, and the red nitrous fumes instantly combine with the volatile alkali. The receivers are presently filled with white clouds of nitrous ammoniac; and in this manner a wrong conclusion may easily be drawn, from the want of the orange colour of the nitrous fumes. A considerable quantity of nitrous air may have been formed, and yet no orange colour appear, owing to this circumstance; and therefore it is easy to understand how a small quantity of nitrous air may be most effectually disguised by the same cause.

12. These observations are made principally for the sake of those who may wish to repeat these experiments. The main point to be established, is the actual formation of nitrous air by this method. And this truth I consider as proved beyond all controversy; for by continuing the process patiently, and applying repeatedly fresh portions of strong volatile alkali to the same manganese, kept constantly hot in the gun-barrel, I have often collected large jars of air, which was proved to be highly nitrous by mixture with atmospherical or with dephlogisticated air.

13. It is not easy to say, whether in this process dephlogisticated nitrous air, or even nitrous acid itself, be not sometimes immediately formed by the action of the volatile alkali on the manganese. Traces of the former, in some instances, seem to discover themselves; but I do not speak decidedly on this head. As to the latter, it is very certain, that fumes of the nitrous acid often circulate in the jars that receive the air. But possibly these fumes may arise from the decomposition of nitrous air, by means of the superfluous dephlogisticated air of the manganese.

14. The steam of boiling water was applied to red hot manganese in a similar way; not the least nitrous appearance; but the fixed and dephlogisticated airs were generated much more plentifully than when the manganese was urged by mere heat. When these airs had been collected in large quantities, the volatile alkali was applied as before to the residuum of the manganese, and nitrous air soon appeared.

15. As manganese is known to produce a very extraordinary change upon spirit of salt in a moderate heat, it seemed not improbable, that a still greater change might take place by working in this method. Accordingly I forced the vapour of  
boiling

boiling spirit of salt to pass through red-hot manganese. This experiment did not answer my expectation; the product was a mixture of fixed and inflammable air. But it deserves to be noticed, that even in this case, after the effect of the spirit of salt had been tried for a long time, a production of nitrous air upon the application of volatile alkali to the same manganese soon took place.

16. As there are many other substances besides the calx of manganese, which are known, *per se*, to afford dephlogisticated air, or a mixture of this with fixed air, it was natural to conclude from analogy, that such substances upon the application of volatile alkali would not fail to afford nitrous air.

It is best, however, in these matters to trust as little as possible to conjectures, and to bring every opinion to the test of experiment. Manganese is so singular a substance, that it is perhaps hardly safe, from what happens in making trials with it, to infer in any instance of another calx of a metal a similarity of effect. Red lead, however, is known to agree in such a variety of chemical effects with manganese, that I find it difficult to persuade myself that the volatile alkali properly applied to it would not yield nitrous acid or nitrous air; yet I have hitherto in vain attempted to bring this about. The red lead, indeed, melts during the process, flows into the cooler parts of the tube, and often choaks the passage of the air; but in some trials a great deal of air has been collected before that happened, and without any symptom of a nitrous mixture. It seems difficult to explain the reason of the failure; perhaps with a better adapted apparatus, and more perseverance, either the production in question may be obtained, or the cause of the failure discovered.

17. With calcined green vitriol I had much better success. The salt was calcined to whiteness, and put into a gun-barrel; and, after several trials of forcing the volatile alkali through the hot tube, I procured by the operation some ounces of strong nitrous air.

So extraordinary an effect would no doubt have proved highly grateful to the ancient chemists, and have been by them denominated a transmutation.

In the course of my enquiries, I considered this experiment as important, because it proved; that the same combinations might take place when substances were made use of different from manganese.

18. As calcined green vitriol, *per se*, in a strong heat yields dephlogisticated air, I had now no doubt but that any substance which had this property might, by similar treatment, be made to afford nitrous air.

But in this supposition I was entirely mistaken. The volatile alkali was applied to some calcined alum at the moment when it was yielding in a strong heat plenty of dephlogisticated air. The product was an astonishing quantity of inflammable air, mixed with hepatic air and actual sulphur. The residuum of the alum had a strong hepatic smell, and contained particles of perfectly formed sulphur.

Most of these experiments, if not all, were repeated in earthen tubes instead of gun-barrels, and with the same success.

19. It now only remains, that I should briefly propose what occurs to me as the probable theory and explanation of the facts related.

The ingredients which enter into the composition of nitrous acid seem to be the two principles or elements of the atmosphere,

sphere, *viz.* phlogisticated and dephlogisticated air. That this is the case, there seems little reason to doubt. Both the composition and decomposition of nitrous acid renders the supposition probable. For,

1. Nitrous air and dephlogisticated air by mixture produce nitrous acid; and nitrous acid, by mere heat, is converted into a mixture of phlogisticated and dephlogisticated airs.

2. Nitrous air, by the methods already related, is changed into phlogisticated air, and these methods seem to consist in abstracting from the nitrous air a quantity of dephlogisticated air.

3. When nitrous acid and nitre are produced in a natural way, the process is not well understood; but the presence of the atmosphere is known to be necessary.

4. Mr. CAVENDISH's experiment is decisive on this point. The union of the two airs in question is effected by means of the electrical spark, and nitrous acid is the product.

In the next place we are to consider, that volatile alkali contains phlogisticated air; for,

1. Volatile alkali, by mere heat, or by the electrical spark, is changed into a mixture of phlogisticated and inflammable air; and,

2. The residuum of volatile alkaline air, after the calces of lead have been revived in it, is phlogisticated air.

Therefore, when volatile alkali, in the form of fume or air, is applied to red-hot manganese, or calcined green vitriol (substances which are then yielding dephlogisticated air), with these facts in view, it seems not difficult to conceive, that one of the ingredients of the alkali, *viz.* phlogisticated air, should combine with dephlogisticated air, and form nitrous acid or nitrous air. If nitrous acid be formed, it will indeed in that

heat, as has been observed, be instantly decomposed; but if the effect of the union be nitrous air, that will sustain the heat without decomposition. How it happens, that nitrous air should be formed, and not nitrous acid, or what the reason is, that nitrous air can sustain a red heat without decomposition, when nitrous acid cannot, I am unable to say; and it is better to acknowledge our ignorance than advance groundless conjectures. So much, I think, may be pronounced as certain, *viz.* that nitrous air contains less dephlogisticated air than nitrous acid; because it requires the addition of dephlogisticated air to become nitrous acid.


And, lastly, if I mistake not, the experiment with the calcined alum proves, that, in order to produce nitrous air, it is not sufficient merely to apply volatile alkaline air to a substance which is actually yielding dephlogisticated air.

Perhaps the presence of another substance is required, which has a strong attraction for phlogiston. Perhaps, in the experiments with the calces of manganese and of iron, the inflammable principle of the volatile alkali combines with the calces of the metals, and the phlogisticated air, the other component part, unites with the dephlogisticated air; and if so, it seems not improbable to suppose, that when alum is made use of, the inflammable principle of the volatile alkali having little or no attraction for clay, the basis of the alum, should combine with its acid and form sulphur. If this reasoning be true, then it follows, that the vitriolic acid has a stronger affinity to the inflammable principle than it has to phlogisticated air; and the process with the green vitriol and manganese is to be explained by the operation of a double affinity: the inflammable principle of the volatile alkali joins with the calx of iron, the  
basis



basis of the vitriol, or with the manganese, and the phlogisticated air with the dephlogisticated air produced by the acid in the red heat.

Those who chuse to reject the doctrine of phlogiston must make the necessary alteration in these expressions; but the reasoning will be much the same.



## P R E S E N T S

MADE TO THE

## R O Y A L S O C I E T Y

From November 1788 to June 1789;

W I T H

## The N A M E S of the D O N O R S.

Presents.

Donors.

1788.

Nov. 6. Transactions of the Royal Irish Academy,  
1787. Dublin. 4°Transactions of the Society for Encourage-  
ment of Arts, Manufactures, and Com-  
merce. Vol. VI. London, 1788. 8°Commentationes Societatis Regiæ Scientiarum  
Gottingensis ad a. 1785 et 86. Vol. VIII.  
Gottingæ, 1787. 4°The Life of Capt. James Cook, by Andrew  
Kippis. London, 1788. 4°Georgii Forster de Plantis esculentis infula-  
rum Oceani Australis Commentatio. Bero-  
lini, 1786. 8°Florulæ infularum australium Prodromus,  
Auctore G. Forster. Gottingæ, 1786. 8°A new experimental Enquiry into the Nature  
and Qualities of the Cheltenham Water.  
Bath, 1788. 8°Fragments of English and Irish History in the  
ninth and tenth Centuries, translated  
from the Icelandic, by Grimr Johnson  
Thorkelin. London, 1788. 4°The Royal Irish Aca-  
demy.The Society for Encou-  
ragement of Arts,  
Manufactures, and  
Commerce.The Royal Society of  
Sciences at Gottin-  
gen.The Rev. Andrew Kip-  
pis, D. D. F. R. S.George Forster, M. D.  
F. R. S.Anth. Fothergill, M. D.  
F. R. S.

Mr. John Nichols.

Nov.

Presents.

Donors.

1788.

- Nov. 6. North View of the Church of Woodbridge, in the County of Suffolk, drawn by J. Johnson, engraved by J. Basire. 1788. fol.  
 Description of Woodbridge Church. fol.  
 Mathematical Essays, by the Rev. J. Hellins. London, 1788. 4°  
 The Generation of animal Heat investigated, by E. Peart. Gainsborough, 1788. 8°  
 The Duties of a Regimental Surgeon considered, by R. Hamilton. 2 Vols. London. 8°  
 Extrait des Observations Astronomiques et Physiques faites à l'Observatoire Royal, en l'Année 1787. 4°  
 Memoire sur les Hopitaux de Paris, par M. Tenon. Paris, 1788. 4°  
 Essai sur le Phlogistique et sur la Constitution des Acides, traduit de l'Anglois de M. Kirwan, avec des Notes de M. M. de Morveau, Lavoisier, De la Place, Monge, Berthollet, et de Fourcroy. Paris, 1788. 8°  
 Examen Physico-chimique des Principes de l'Air et du Feu, par M. Le Semelier. 2 Vols. Paris, 1788. 8°  
 Observations generales sur les Hopitaux, par M. Iberti. Londres, 1788. 8°  
 Memoire sur l'Epidemie qui a regné dans la Province du Poitou pendant 1784 et 1785, par M. J. G. Gallot. Poitiers, 1787. 4°  
 Ephemerides Astronomicæ annorum 1788 et 1789 ad meridianum Mediolanensem supputatæ ab Ang. de Cefaris. 2 Vol. Mediolani, 1787. 8°  
 Occursus Medici de Vaga ægritudine infirmitatis Nervorum, Andreae Comparetti. Venetiis, 1780. 8°  
 A. Comparetti Observationes opticae de Luce inflexa et coloribus. Patavii, 1787. 4°  
 13. A Map of the Peninsula of India, from the Khrishnah River to Cape Comorin, by J. Rennell, in Two Sheets.  
 A new Grammar, to teach French to Englishmen, by Dom. Blondin. London, 1788. 8°

M. R. Loder and Mr. J. Nichols.

---

The Rev. John Hellins.

E. Peart, M. D.

R. Hamilton, M. D.

Comte de Caffini, F. R. S.

M. Tenon, of the Royal Academy of Sciences of Paris.

The Translator.

M. Le Semelier.

M. Iberti.

M. Gallot.

The Astronomers of Milan.

Professor Comparetti of Padoua.

---

James Rennell, Esq. F. R. S.

Dom. Blondin.

Nov.

Presents.	Donors.
1788.	
Nov. 13. Discours de l'Academie Royale de l'Histoire au Roi, à l'Occasion de la Naissance de l'Infant, en Espagnol et en Francois. Madrid, 1788. 8°	Dom. Blondin.
20. Linguarum totius Orbis vocabularia comparativa, Augustissimæ cura collecta. Sectionis primæ pars prior. Petropol. 1786. 4°	The Empress of Russia.
Kongl. Vetenskaps Academiens Nya Handlingar. Tom. VIII. för år 1787, 3d and 4th Quarter; and Tom. IX. för år 1788, 1st and 2d Quarter. Stockholm. 8°	The Royal Academy of Sciences of Stockholm.
Meteorological Journal kept at Fort William, Bengal, from Nov. 20, 1773, to June 28, 1787. 3 Vols. MS. fol.	Col. Pearce.
The Natural History and Antiquities of Selborne, in the County of Southampton, by Gil. White. London, 1789. 4°	Messrs. B. White and Son.
27. A Brass Model of a Machine to turn a Mill by the Steam.	Col. Pearce.
A Model in Ivory of a vertical Axis, supported by three friction Wheels.	_____
A Model in Wood of the same, with four friction Wheels, when the Weight is great.	_____
A Model of a Marine Barometer Tube.	_____
Meteorological Diary, kept at Madras, from Feb. 18, to Sept. 30, 1787. MS. 4°	William Petrie, Esq.
Perspective View of the Font in the Church of Worlingworth, in the County of Suffolk, drawn by N. Revett, engraved by G. Vertue. 1753.	Mr. J. Nichols and Mr. R. Loder.
Rede bey der Gedächtnisfeyer Hevelii den 28 Jan. 1787, gehalten von E. P. Blech. Dantzig. 4°	Dr. E. P. Blech, of Dantzig.
Dec. 11. Floræ Cantabrigiensi Supplementum alterum, Auctore R. Relhan. Cantabrigiæ, 1788. 8°	The Rev. Richard Relhan, M. A. F. R. S.
18. Supplement à l'Edition in folio des Oeuvres de M. Perronet. Paris, 1789. fol.	M. Perronet, F. R. S.
Commentarii de Rebus in Scientia Naturali et Medicina gestis. Vol. XXIX. Lipsiæ, 1787. 8°	Mr. Hurlock, F. R. S.
1789.	
Jan. 8. Tabula exhibens crania Rhinocerotis Africae, et Rhinocerotis Asiae, 1787.	Petrus Camper, M. D. F. R. S.
An Account of the Pelew Islands, composed from the Journals of Capt. Henry Wilson	George Keate, Esq. F. R. S.

Presents.

Donors.

1789.

and some of his Officers, by G. Keate.  
London, 1788. 4°

Jan. 8. The London Medical Journal. Vol. IX.  
London, 1788. 8°

Description des Gîtes de Minerai et des  
Bouches à Feu de la France. 3<sup>me</sup> et 4<sup>me</sup>  
Partie. Paris, 1789. 4°

An Essay on the Recovery of the apparently  
dead, by C. Kite. London, 1788. 8°

15. An Essay on the Fracture of the Patella or  
Knee-pan, by John Sheldon. London,  
1789. 8°

22. Method to discover the Difference of the  
Earth's Diameters, by T. Williams. Lon-  
don, 1788. 8°

29. Decouverte d'une Methode peu couteuse,  
efficace et assurée de traiter tous les  
Hommes decedés, afin de rappeler à la  
Vie ceux qui ne sont morts qu'en appa-  
rence, par J. G. C. A. Baron de Hupfch.  
Cologne, 1789. 8°

Enquiry in the First and general Principles of  
Animal and vegetable Life, by R. Harring-  
ton. London, 1781. 8°

Thoughts on the Properties and Formation of  
the different Kinds of Air. London, 1785,  
8°

A Letter endeavouring to prove, that the  
newly adopted Opinions of inflammable and  
dephlogisticated Airs forming Water, and  
the Acids being compounded of the different  
Kinds of Air, are fallacious, by R. Har-  
rington. London, 1788. 8°

Feb. 5. Abstracts from a Meteorological Register, kept  
in Hudson's Bay, from October, 1784, to  
July, 1788, MS. 4°

Astronomisches Jahrbuch für das Jahr 1791,  
von J. E. Bode. Berlin, 1788. 8°

12. A short Appendix to Dr. Monro's Treatise on  
Medical and Pharmaceutical Chemistry.  
London, 1789. 8°

A Meteorological Journal of the Year 1788,  
kept in Pater-noster-Row. London, by  
W. Bent. 4°

Samuel Foart Simmons,  
M. D. F. R. S.

Baron de Dietrich.

Mr. Charles Kite.

Mr. Sheldon, F. R. S.

Mr. Thomas Williams.

Baron de Hupfch.

Mr. John Sherwen.

—————

—————

John M<sup>c</sup> Nab, Esq.

Mr. J. E. Bode, F.R.S.

Donald Monro, M. D.  
F. R. S.

Mr. Willam Bent.

## Presents.

## Donors.

- 1789.
- Feb. 19. *Il Mercurio Italico*. Gennajo, 1789. London. 8°  
 A Register of the Barometer, Thermometer, Hygrometer, Rain, Dew, Evaporation, Wind, and Appearance of the Sky, at Gravesend, in Kent, for the Years 1787 and 1788, MS. fol.; with a Chart of Meteorology for the Year 1788. Mr. Francis Saftres.  
 Mr. Charles Kite.
- March 5. The Preface to a Specimen of a general Astronomical Catalogue, arranged in Zones of North Polar Distance, by F. Wollaston. London, 1789. 8°  
 The Rev. Francis Wollaston, LL.B. F. R. S.  
 Chronological Tables of the High Sheriffs of the County of Lincoln. London, 1779. 4°  
 Edmund Turnor, Esq. F. R. S.
12. A System of Hydrostatics, by T. Parkinson. Cambridge, 1789. 4°  
 The Rev. Thomas Parkinson, B. D. F. R. S.
- April 2. A Treatise of the *Materia Medica*. 2 vols. Edinburgh, 1789. 4°  
 William Cullen, M. D. F. R. S.
30. *Memoires de l'Academie Imperiale et Royale des Sciences et Belles Lettres de Bruxelles*: Tome V. Bruxelles, 1788. 4°  
 The Imperial and Royal Academy of Sciences of Brussels.  
*Memoirs of the Medical Society of London*. Vol. II. London, 1789. 8°  
 The Medical Society of London.  
 A General History of Music, by C. Burney. Vol. III. and IV. London, 1789. 4°  
 Charles Burney, Mus. D. F. R. S.  
*Reliquiæ Rudbeckianæ*, cura J. E. Smith. Londini, 1789. fol.  
 James Edward Smith, M. D. F. R. S.  
*Il Mercurio Italico*. Febraio e Marzo, 1789. London. 8°  
 Mr. Francis Saftres.
- May 7. *Histoire de l'Academie Royale des Sciences*, année 1786. Paris, 1788. 4°  
 The Royal Academy of Sciences of Paris.  
 Meteorological Account of the Weather at Madras, from June 1, 1787, to May 31, 1788. Madras, 1788. 4°  
 Sir Joseph Banks, Bart. Pr. R. S.  
*Codex Diplomaticus Siciliæ ab a. 827 ad 1072*, cura Alph. Airoidi. Tom. I. Panormi, 1788. fol.  
 Sir William Hamilton, K. B. F. R. S.  
 The History and Antiquities of Newcastle upon Tyne, by J. Brand. 2 Vols. London, 1789. 4°  
 The Rev. John Brand, M. A. Sec. A. S.  
*Histoire Naturelle du Jorat, et de ses Environs*, par M. le C. G. de Razoumowsky. Tomes II. Lausanne, 1789. 8°  
 Comte G. de Razoumowsky.

Presents.

Donors.

1789.

- |  |   |
|--|---|
| <p>May 7. Lettre contenant la Description des frottoirs Electriques d'une nouvelle Construction. Harlem, 1789. 4°</p> <p>14. Original Letters written during the Reigns of Henry VI. Edward IV. and Richard III. published by Sir John Fenn. Vol. III. and IV. London, 1789. 4°</p> <p>Lettre au Docteur Maty, par l'Abbé Coyer. Bruxelles, 1767. 12°</p> <p>21. Experiences sur les Vegetaux, par J. Ingen-houfz. Tome II. Paris, 1789. 8°</p> <p>Traité des Principales Maladies externes et internes, par J. F. de Herrenschwand. Berne, 1788. 4°</p> <p>Memoires sur l'Agriculture du Boulonnois. Boulogne, 1784. 8°</p> <p>A Portrait of the late Sir William Watfon, Knt. M. D. V. P. R. S. painted by Abbot.</p> <p>28. A new English Translation of the Pentateuch, by Ifaac Dalgado. London, 1789. 4°</p> <p>De la Liberté de Commerce. Beauvais, 1789. 8°</p> <p>June 11. Britannia, by W. Camden, translated and enlarged, by R. Gough. London, 1789. 3 Vols. fol.</p> <p>Essai sur la Montagne salifere du Gouvernement d'Aigle dans le Canton de Berne, par F. S. Wild. Geneve, 1788. 8°</p> <p>The Coach of Safety. Two coloured Plates, with a printed Account of it.</p> <p>18. J. Gærtner de fructibus et feminibus Plantarum. Stutgardia, 1788. 4°</p> <p>July 2. A View of the Reign of Frederick II. of Prussia, by J. Gillies. London, 1789. 8°</p> <p>Letters from Barbary, France, Spain, Portugal, &amp;c. by an English Officer. 2 Vols. London, 1788. 8°</p> <p>Vitæ antiquæ Sanctorum qui habitaverunt in ea parte Britannniæ nunc vocata Scotia vel in ejus insulis; edidit J. Pinkerton. Londini, 1789. 8°</p> <p>The History and Antiquities of Canonbury-house, at Iflington, by J. Nichols. London, 1789. 4°</p> | <p>Martin Van Marum, M. D. of Harlem.</p> <p>Sir John Fenn, Knt.</p> <p>William Seward, Esq. F. R. S.</p> <p>John Ingen-houfz, M. D. F. R. S.</p> <p>Jonathan Watfon, Esq. F. R. S.</p> <p>Le Baron de Courfel.</p> <p>William Watfon, Esq. F. R. S.</p> <p>Mr. William Blizard, F. R. S.</p> <p>M. Briffon.</p> <p>Richard Gough, Esq. F. R. S.</p> <p>M. Wild.</p> <p>Mr. John Hatchett.</p> <p>Joseph Gærtner, M. D. F. R. S.</p> <p>John Gillies, LL.D. F. R. S.</p> <p>Major Jardine.</p> <p>John Pinkerton, Esq.</p> <p>Mr. John Nichols.</p> |
|--|---|

Presents.

Donors.


1789.

July 2. Voyage à la Nitriere naturelle qui se trouve à Molfetta en Pouille, par M. Zimmerman. Paris, 1789. 8°

Professur Zimmerman, of Brunswick.

J. Churchman's Address in Support of the Principles of the Magnetic Variation, and their applications in determining the Longitude at Sea. fol.

Mr. John Churchman.





---

A N  
I N D E X  
T O T H E  
S E V E N T Y - N I N T H V O L U M E  
O F T H E  
P H I L O S O P H I C A L T R A N S A C T I O N S .

A.

*ACCOUNT* of a bituminous lake or plain, in the island of Trinidad, p. 65.  
Account of a particular change of structure in the human ovarium, p. 71. Account of a monster of the human species, p. 157. Account of a sinking-in of the ground near Ketton, in Rutland, p. 164.

*Acid, nitrous.* A red heat not necessary to the conversion of nitrous acid into pure air, p. 147. Volatile vitriolic acid contains the proper element of dephlogisticated air, p. 292. Prussian blue supposed to have acquired something that is of the nature of an acid, p. 298. Experiments on the production of nitrous acid and nitrous air, p. 300. A relation subsists between nitrous acid and volatile alkali, *ibid.* Nitrous acid produced by a mixture of nitrous air and dephlogisticated air, p. 311. Vitriolic acid supposed to have a stronger affinity to the inflammable principle than it has to phlogisticated air, p. 312.

*Action* of points in electricity, experiments on, pp. 278, 281.

*Air, inflammable or dephlogisticated,* when either of them is extracted from any substance in contact with the other kind of air, so that one of them is made to unite with the other in what may be called its *nascent state*, the result will be *fixed air*; but if both

of them be completely formed before their union, the result will be *nitrous acid*, p. 12. All inflammable air, according to the antiphlogistic hypothesis, comes from water only, *ibid.* A small quantity of inflammable air is procured, by sending a steam over melted sulphur, p. 15. Inflammable air produced by a dissolution of iron in diluted vitriolic acid, *ibid.* Vitriolic acid air contains the same inflammable principle with inflammable air, p. 16. Nitrous air, when mixed with dephlogisticated air, has no tendency to produce phlogisticated air, p. 149. Nitrous air has no effect on phlogisticated air, *ibid.* Nitrous air does not contain so much phlogiston as an equal bulk of inflammable air, p. 150. Alkaline air converted into inflammable air, p. 295. Nitrous air and dephlogisticated air by mixture produce nitrous acid; and nitrous air, by mere heat, is converted into a mixture of phlogisticated and dephlogisticated airs, p. 309. Nitrous air changed into phlogisticated air, by abstracting from the nitrous air a quantity of dephlogisticated air, *ibid.* The residuum of volatile alkaline air, after the calces of lead have been revived in it, is phlogisticated air, p. 311. The vitriolic acid supposed to have a stronger affinity to the inflammable principle than it has to phlogisticated air, p. 312.

*Alkali*, volatile, changed into a mixture of phlogisticated and inflammable air by mere heat, p. 311.

*Almanack*, nautical, the most perfect work of the kind, p. 60.

*Altitude*. See *Quadrant of Altitude*.

*Amphibia*, a class of animals so called by Linnæus, p. 21. Observations on this class, *ibid.* Linnæus particularly unfortunate in the construction of this class, *ibid.* *Amphibia Nantes* not furnished with lungs, *ibid.*

*Anderfon*, Alexander, on a bituminous lake in the Island of Trinidad, p. 65.

*Affam*, a country bordering on, and much connected with, Thibet, p. 107.

*Attraction*, law of. See *Problems*.

*Azimuth*. See *Quadrant of Altitude*.

## B.

*Babar*, the country adjacent, highly injurious to European constitutions, p. 80.

*Baillie*, Matthew, on a particular change of structure in the human ovarium, p. 71. See *Ovarium*, also *Ovaria*.

*Balls* in electricity, a variety of, unnecessary, p. 280.

*Barker*, Thomas, Esq. Abstract of a register for the year 1788, of the barometer, thermometer, and rain, at Lyndon in Rutland. Also of the rain in Hampshire and Surrey, p. 162.

*Barometer*, state of, at London during the year 1788, p. 114—138.

*Borer*, destructive to snips, p. 68.

*Boutan* and Thibet, some account of the vegetable and mineral productions of, p. 79.

*Bray* (La). See *Tar-lake*.

*Buxaduar*,

*Buxaduar*, many of the plants peculiar to Bengal require nursing at, p. 80. *Buxaduar* unhealthy to strangers from May till September, p. 81. State of the thermometer there, *ibid.* Several excellent springs of water in its neighbourhood, *ibid.*

C.

*Chalu*, three springs discovered near this place, forcing their way through the ground with violence, and giving rise to a lake many miles in extent, well-stored with water, fowl and excellent fish, p. 92. The water of the lake impregnated with alum and selenetic earth, p. 93. The productions of *Chalu*, *ibid.*

*Chepta*, its situation and productions, p. 84. Here are several springs, and one slightly impregnated with iron, *ibid.* Between *Chepta* and *Pagha*, a mineral well, strongly impregnated with iron, discovered by Mr. Saunders, *ibid.* State of the thermometer at *Chepta*, *ibid.*

*Chooka*, two mineral wells slightly impregnated with iron, discovered by Mr. Saunders near, p. 83. Natives of *Chooka* not unacquainted with the method of extracting iron from the stones, but despise its use in building, *ibid.*

*Cold*, artificial, experiments relating to the production of, p. 199.

*Coluber lauticaudatus*, a species of venomous serpents, the fangs of which are as small as common teeth, p. 30.

*Comet*, observations on, p. 151. Its situation, p. 152.

D.

*Datura ferox*, or Thorn-apple, used medicinally in China and some parts of Thibet, p. 89. A powerful narcotic, *ibid.*

*Dog*. See *Letter*.

*Duina*, productions of, p. 92. State of the thermometer at, *ibid.*

*Dukaigua*, state of the thermometer at, soil, and productions of, p. 90. Here are many springs slightly impregnated with a selenetic earth, *ibid.*

E.

*Earth*, and other planets of the Copernican system, are bodies not luminous in themselves, p. 213.

*Eclipse*. See *Sun*.

*Electricity*, experiments and observations on, p. 265. On the excitation of electricity, *ibid.* Plate machines do not collect more electricity than cylinders do with half the rubbed surface, p. 269. The line of light on a cylinder departing from a simple cushion consists of returning electricity, p. 273. The projecting part of the cushion compensates the electricity upon the cylinder, and by diminishing its intensity prevents its striking back in such large quantities as it otherwise would do, *ibid.* The method of producing considerable intensity, *ibid.* The action of the cylinder, by a simple cushion

cushion or the hand, was first improved by the addition of a leathern flap; then by moistening the rubber; afterwards by applying the amalgam; and, lastly, by the addition of a silk flap, p. 278. Experiments on the luminous appearance of electricity and the action of points, *ibid.* The escape of negative electricity from a ball attended with the appearance of strait sharp sparks with a hoarse or chirping noise; pp. 278, 279. A variety of balls in electricity unnecessary, p. 280. Volta's opinion, that a point is the coating to an infinitely small plate of air, erroneous, p. 281. In any electrified conductor the transition or escape of electricity will be made chiefly from that part of the surface which is the most remote from the natural state, p. 282. Thicker glasses require much less electricity to produce an intensity which breaks them than thinner do, p. 284. The perforation of glass by the long spark, or by the spark through oil or cement, supposed to depend on the intensity of the electricity, which has not time to diffuse itself, but charges a minute part of the surface very high, p. 285. Muscovy talc, a very perfect non-conductor; capable of being divided into plates of less thickness than a two-hundredth part of an inch; in consequence of its great capacity gives very strong electric shocks; and its laminæ, contrary to the assertion of Beccaria, are naturally in strong opposite states of electricity, and flash to each other when torn asunder in the dark, *ibid.*

*Euler's* hypothesis of refraction of the differently refrangible rays of light proved to be false, p. 257. His hypothetical principle neither fit for rendering a telescope achromatic, nor to account for the distinctness of the human vision, *ibid.*

*Excitation* of electricity, experiments on, p. 265. Strength of the excitation ascertained, *ibid.* Silk the chief agent in excitation of electricity, p. 271.

*Experiments* and observations on the principle of acidity, the composition of water, and phlogiston, p. 7. Experiments on the phlogistication of spirit of nitre, p. 139. Experiment proving that *heat* and not *light* gives colour to spirit of nitre, p. 140. Experiments on the congelation of quicksilver in England, p. 199. Experiments relating to the production of artificial cold, *ibid.* Experiment proving that mercury may be frozen not only in England in summer, but even in the hottest climate, at any season of the year, p. 202. Experiments illustrating the subject of producing cold, pp. 209, 210. Experiment illustrating the theory of vision, p. 261. Experiments and observations on electricity, p. 265. Experiments shewing that the office of silk is not merely to prevent the return of electricity from the cylinder to the cushion, but that it is the chief agent in the excitation; while the cushion serves only to supply the electricity, and perhaps increase the pressure at the entering part, pp. 265, 266. Experiments on the luminous appearances of electricity and the action of points, p. 278. Experiments on the action of points in electricity, pp. 281, 282. Experiments on the transmission of the vapour of acids through an hot earthen tube, and further observations relating to phlogiston, p. 289. Experiment on oil of vitriol, *ibid.* Experiment on spirit of nitre, p. 291. Experiment in which alkaline air is converted into inflammable air, p. 295. Experiments relating to the doctrine of  
phlogiston,

phlogiston, *ibid.* Experiments shewing that manganese, *per se*, gives airs of different kinds, but chiefly fixed and dephlogisticated airs, as soon as it is subjected to a considerable heat, p. 306.

*Eyes.* Fifteen inches the distance at which the generality of eyes see with the most distinctness, p. 258.

## F.

*Fangs*, venomous, in serpents, merely offensive weapons, p. 33. Situation of venomous fangs always in the anterior and exterior parts of the upper jaw, p. 34.

*Fruetification* chiefly depends on rain falling at the latter end of the season of flowering, p. 39.

## G.

*Gentoo Boy*, an astonishing living subject, having his brother adhering to his breast, p. 157. Portrait of, p. 159.

*Glass*, the perforation of, by the long electric spark, or by the spark through oil or cement, supposed to depend on the intensity of the electricity, which has not time to diffuse itself, but charges a minute part of the surface very high, p. 285.

*Globe*, terrestrial and coelestial, few instruments that better fulfil their design in general, p. 1.

*Gray*, Edward Whitaker, on the class of animals called, by Linnæus, Amphibia, and particularly on the means of distinguishing those serpents which are venomous from those which are not so, p. 21.

*Ground*, account of a sinking in of the, near Ketton in Rutland, p. 164.

## H.

*Heavens*, remarks on the construction of, p. 212.

*Herschel*, Dr. William, on a comet, p. 151.

———— Catalogue of a second thousand of new nebulae and clusters of stars; with introductory remarks on the construction of the heavens, p. 212.

*Hunter*, John, on the identity of the species of the Dog, Wolf, and Jackal, p. 160.

*Hutchinson*, Rev. Mr. B. on the dryness of the year 1788, p. 37.

## I.

*Jackal.* See *Letter*.

*Iron*, dissolved in concentrated acid of vitriol, produces vitriolic acid air; but being dissolved in diluted vitriolic acid, produces inflammable air, p. 15. The decomposition of water, by means of iron, a fallacy, p. 17. Iron imbibes nothing but water when it parts with its phlogiston, *ibid.*

## K.

*Kite* (electrical) raised by Loammi Baldwin during the approach of a severe thunder-storm, p. 275. Effect of the electricity upon him, p. 276.

## L.

*Lac*, the produce of, and a staple article of commerce in, Assam, p. 107. Lac neither a gummy nor a resinous substance, though it has some properties which are common to both, *ibid.* Known in Europe by the different appellations of stick-lac, seed-lac, and shell-lac, *ibid.* Used as a dye by the natives of Assam, p. 109. The method of purifying it, *ibid.*

*Lake*, bituminous. See *Trinidad*, also *Tar-lake*.

— Tincal found in a lake in Thibet, which is fifteen days journey northward from Tissoolumboo, p. 96. Great quantities of rock salt found in this lake, p. 97.

*Letter*, supplementary, on the identity of the species of the Dog, Wolf, and Jackal, p. 160.

*Letters*, two, concerning a monster of the human species, p. 157, 158.

*Light*, capable of giving colour to spirit of nitre, p. 140.

— an attempt to explain a difficulty in the theory of vision, depending on the different refrangibility of light, p. 256. Euler's hypothesis of refraction of the differently refrangible rays of light proved to be false, p. 257.

*Lime-stone*, rock of, discovered at Pুনukha; very advantageously situated for being worked, and exceedingly pure, p. 83.

*Linnaeus*, many of his descriptions given in a very careless manner, p. 21. Particularly unfortunate in the construction of the class called amphibia, *ibid.*

*Liquor*, green acid, procured by the explosion of dephlogisticated and inflammable airs in close vessels, pp. 8, 9, 10.

## M.

*Manganese, per se*, gives airs of different kinds, but chiefly fixed and dephlogisticated airs, as soon as it is subject to a considerable heat, p. 306.

*Mangrove* swamps in the island of Trinidad, p. 65.

*Marsham*, Robert, indications of spring, p. 154—156.

*Maskelyne*, Dr. Nevil, on a difficulty in the theory of vision, depending on the refrangibility of light, p. 256.

*Mercury,*

- Mercury*, experiments on the congelation of, p. 199. Vide *Experiments. Quicksilver.*
- Meteorological journal* for the year 1788, kept at the apartments of the Royal Society, p. 114—137. See *Tables.*
- Milner*, Rev. Isaac, on the production of nitrous acid and nitrous air, p. 300.
- Monster* of the human species, account of, p. 157. See *Gentoo Boy.*
- Morgan*, William, on the method of determining, from the real probabilities of life, the value of a contingent reversion in which three lives are involved in the survivorship, p. 40.
- Murishong*, a pleasant and healthy situation, p. 82. The soil, rich and fertile, produces good crops, *ibid.*
- Muscovy talc*, a very perfect non-conductor; capable of being divided into plates of less thickness than a two-hundredth part of an inch; in consequence of its great capacity gives very strong electric shocks; and its laminæ, contrary to the assertion of Becaria, are naturally in strong opposite states of electricity, and flash to each other when torn asunder in the dark, p. 285. One solid inch of Muscovy talc contains electricity enough to charge a conductor of 7 inches diameter, and 135 feet long, so high as to give a nine-inch spark at least, p. 286.

## N.

- Naja*, a species of serpent, very venomous, p. 25.
- Natron*, phosphorated, produces more cold by solution in the diluted nitrous acid, than the vitriolic natron, p. 208. Vitriolated natron, added to marine acid, undiluted, produces very nearly as great a degree of cold as when mixed with the diluted nitrous acid, p. 210.
- Nebulae* and clusters of stars, catalogue of a thousand, p. 236—254. See *Tables.*
- Nicholson*, Mr. William, on electricity, p. 265.
- Nitre*, spirit of, experiments on the phlogistication of, p. 139. Light only capable of giving colour to spirit of nitre, contained in phials with ground stoppers, p. 140. Experiment proving that *heat* and not *light* gives colour to spirit of nitre, *ibid.* Spirit of nitre, in its most dephlogisticated state, consists of a proper saturation of the acids with phlogiston, p. 289. Experiment on spirit of nitre, p. 291.
- Nitrous air.* See *Air.*

## O.

- Objections* to the experiments and observations relating to the principle of acidity, the composition of water, and phlogiston, considered; with further experiments and observations on the same subject, p. 7.
- Observations* and experiments on the principle of acidity, the composition of water, and phlogiston, p. 7. Observations on the class of animals called, by Linnæus, Amphibia;

phibia; particularly on the means of distinguishing those serpents which are venomous from those which are not so, p. 21. Observations on the dryness of the year 1788, p. 37. Observations on a comet, p. 151. Observations and experiments on electricity, p. 265. Further observations relating to phlogiston, p. 289.

*Ovaria* in women subject to a great variety of changes, p. 71. Many of these changes similar to those which take place in other parts of the body, *ibid.* Ovaria in women have some power within themselves of taking on a process which is imitative of generation, without any previous connection with a male, p. 72.

*Ovarium*, account of a particular change of structure in the human, p. 71. Natural substance of an ovary changed into a fatty mass intermixed with hair and teeth, *ibid.* The growth of hair and teeth in the ovary supposed by Dr. Tyson to be a *lusus naturæ*, *ibid. note.*

## P.

*Pagba*, productions of, p. 84.

*Paragbon*, its soil rich, and abounding with pasture, p. 88. Its productions, *ibid.* Colder here at all seasons than at Tassésudon, p. 89. Iron stones found near Paragbon, and one spring highly impregnated with this mineral, *ibid.*

*Phlogiston*, experiments relating to the doctrine of, p. 295.

*Phosphorus*, after the ascension of it in dephlogisticated air, there is a considerable quantity of fixed air in the residuum, p. 16.

*Piazzi*, Rev. Joseph, result of calculations of the observations made at various places of the eclipse of the sun, which happened June 3, 1788, p. 55.

*Point*. A force acting at a given point may be resolved by an infinite number of ways into two, three, or more forces acting at the same point, either in the same or different planes with the given force and each other; and, *vice versa*, any number of such forces acting in the same or different planes may be reduced into one, p. 185.

Example proving this assertion, *ibid.* See *Problem*.

*Points* in electricity, experiments on the action of, pp. 281, 282. The effects of points depend on the remoteness of their extremities from the other parts of the conductor, p. 282.

*Portrait* of a Gentoo boy, p. 156.

*Precipitate, per se*, yields no fixed air by heat, p. 12.

*Presents*, list of, made to the Royal Society from November 1788 to July 1789, p. 314.

*Priestley*, Rev. Dr. Joseph, objections to the experiments and observations relating to the principle of acidity, the composition of water, and phlogiston, considered; with further experiments and observations on the same subject, p. 7. On the phlogistication.



tion of spirit of nitre, p. 139. On the transmission of acids through an hot earthen tube, and further observations relating to phlogiston, p. 289.

PROBLEMS. If the ages of three persons be given, to determine, from any table of observations, the value of the sum payable on the contingency of the last person's surviving the second, provided the life of the first shall be then extinct, p. 41. This problem solved and demonstrated, *ibid.*

I. Given the law of attraction of each of the parts of a given line in terms of their distance from a given point; to find the attraction of the whole line on the point, p. 186.

II. III. Given the attraction of each of the parts of a given surface in terms of their distance from a given point, and an equation expressing the relation between an absciss and its correspondent ordinates of the surface; to find the attraction of the surface on the given point, p. 187. Also, to find the attraction of a given solid on a given point, p. 189.

IV. Given an equation expressing the relation between two abscissæ and their correspondent ordinate of a solid; to find its solid contents contained between two values of its first abscissa, p. 194.

V. Given an equation expressing the relation between the two abscissæ of a solid, and the correspondent ordinates; to transform the first abscissa into any other, p. 196.

*Prussian* blue generally supposed to be a calx of iron super-saturated with phlogiston, though supposed by some to have acquired something that is of the nature of an acid, p. 298. The quantity of fixed air that *Prussian* blue would yield by mere heat determined, p. 299.

*Punukba*, a root of pure lime-stone discovered here by Mr. Saunders, p. 83. Abundance of fire-wood in this part of the country, p. 83. The houses of this country are lofty, and the timbers substantial, but the inhabitants are unacquainted with the use of lime, p. 83, 84. State of the thermometer here, p. 84.

## Q.

*Quadrant* of altitude, description of an improvement in the application of, to a celestial globe, for the resolution of problems dependent on azimuth and altitude, p. 1.

*Quicksilver*, experiments on the congelation of, p. 199. Experiment proving that quicksilver may be frozen not only in England in summer, but even in the hottest climate, at any season of the year, p. 202. Another experiment illustrating the congelation of quicksilver, *ibid.* Another experiment on the congelation of quicksilver, p. 203.

*Quicksilver*, being frozen and broken, afforded a beautiful appearance of flat plates converging towards a center, p. 204. Congealed quicksilver bore an exact resemblance, both in colour and plated structure, to sulphurated antimony, p. 205.

*Rain,*

- Rain*, a considerable defect of, in the year 1788, p. 37. The quantity of rain fallen at Kimbolton compared with that of the seven preceding years, *ibid.* Monthly state of rain for 1788, p. 38. Quantities of rain during the year 1788, p. 114—138.
- Register*. Abstract of a register, of the barometer, thermometer, and rain, at Lyndon in Rutland: also of the rain at South Lambeth in Surrey, and at Selbourn and Fyfield in Hampshire, p. 162.
- Reichel*, Baron, on a monster of the human species, p. 157.
- Remarks* on the construction of the heavens, p. 212.
- Rock salt*, univervally used for all domestic purposes in Thibet, Boutan, and Naphaul, p. 97.

## S.

- Salts*, neutral, which produce any remarkable degree of cold by solution in mineral acids, lose their freezing property, when deprived of their water of crystallization, p. 209.
- Sarba*, productions of, p. 90. State of the thermometer at, *ibid.*
- Satellite*. See *Saturn*.
- Saturn* (the planet) has a sixth satellite revolving round it, p. 255.
- Saunders*, Robert, on the vegetable and mineral productions of Boutan and Thibet, p. 79.
- Selub*, here is a hot well much frequented by people with venereal complaints, rheumatism, and all cutaneous diseases, p. 94. State of the thermometer when immersed in the water of this well, *ibid.* Its water contains a portion of hepar sulphuris, *ibid.*
- Serpents*, venomous fangs in, merely offensive weapons, p. 33. Situation of venomous fangs always in the anterior and exterior parts of the upper jaw, p. 34. All venomous serpents have only two rows of teeth, but all others have four, p. 35. Proportion of venomous serpents to others, as 1 to 10, p. 36.
- Simadar*, its soil very barren and unpromising, p. 93. Its productions, *ibid.*
- Small-pox*, not treated properly by the physicians of Thibet, p. 105.
- Smeaton*, Mr. John, description of an improvement in the application of the quadrant of altitude to a celestial globe, for the resolution of problems dependent on azimuth and altitude, p. 1.
- Spring*, indications of, p. 154—156.
- Stars*, catalogue of a thousand new nebulae and clusters of, p. 226—254. Every star supposed to be a sun, shining by its own native brightness, pp. 213, 214.
- Sulphur*. By sending a steam over melted sulphur, a small quantity of inflammable air is procured, p. 15. Turbith mineral mixed with sulphur makes it yield vitriolic acid

acid air, *ibid.* Sulphur not that simple substance which the anti-phlogistians suppose it to be, but contains phlogiston, *ibid.*

*Sun*, result of calculations of the observations made at various places of the eclipse of the sun, which happened on June 3, 1788, p. 55.

*Survivorship*. On the method of determining, from the real probabilities of life, the value of a contingent reversion in which three lives are involved in the survivorship, p. 40.

T.

TABLES.

Table of the observations made at Greenwich, Loampit-hill, Oxford, Dublin, Mittau, Berlin, Vienna, Viviers, Perinaldo, Rouen, Milan, Bologna, and Padua, on the eclipse of the sun, which happened June 3, 1788, and of results deduced from the same, p. 62.

Table representing observations made at Warsaw, Prague, Marseilles, Presmunster, and Bagdad, on the eclipse of the sun, which happened June 3, 1788, and results deduced from the same, p. 64.

Meteorological journal for the year 1788, kept at the apartments of the Royal Society, for January, p. 114, 115. February, 116, 117. March, 118, 119. April, 120, 121. May, 122, 123. June, 124, 125. July, 126, 127. August, 128, 129. September, 130, 131. October, 132, 133. November, 134, 135. December, 136, 137.

Table of the greatest, least, and mean heights of the thermometer without and within, and of the barometer; also observations of the quantities of rain, p. 138.

Table containing indications of spring, p. 154—156.

Table of a second thousand of new nebulæ and clusters of stars, p. 226—254. First class, bright nebulæ, p. 286. Second class, faint nebulæ, p. 229. Third class, very faint nebulæ, p. 238. Fourth class, planetary nebulæ, p. 246. Fifth class, very large nebulæ, p. 248. Sixth class, very compressed and rich clusters of stars, p. 250. Seventh class, pretty much compressed clusters of large or small stars, p. 251. Eighth class, coarsely scattered clusters of stars, p. 253.

*Takui*, its productions, p. 94.

*Talc.* See *Muscovy Talc.*

*Tar-lake*, a bituminous lake in the island of Trinidad, p. 65. Called by the French *La Bray*, from the resemblance to, and answering the intention of, ship pitch, *ibid.* Situate in the leeward-side of the island, on a point of land which extends into the sea about two miles, *ibid.* Situation similar to a Savannah, p. 66. Its colour, and even surface, present at first the aspect of a lake of water, *ibid.* Of a circular form, and about three miles in circumference, *ibid.* Its common consistence is that of  
pit-coal,

pit-coal, the colour rather greyer, very friable, and, when liquid, of a jet black colour, p. 67. Some parts of the surface covered with a thin and brittle scoria, a little elevated, *ibid.* Calcined earth mixed with some parts of its common substance, *ibid.* The bituminous substance of this lake supposed to be the *bitumen asphaltum Linnæi*, p. 68. Its substance rendered ductile by heat; and, mixed with a little grease or common pitch, is much used for the bottoms of ships, *ibid.* Supposed to be a preservative against the Boer, *ibid.*

*Taffesudon*, productions of, p. 85. *Taffesudon* lower than the level of Paraghon, p. 89. The capital of Boutan, p. 111.

*Thermometer*, state of, between Bahar and Buxaduar, p. 80. State of it at Buxaduar, p. 81. Its state at Punakha, p. 84. Its state at Chepta, *ibid.* Its state at Sanha, p. 90. Its state at Duina, p. 92. Its state at Tiffolumboo, p. 97, 98. State of the thermometer at London during the year 1788, p. 114—138.

*Thibet*, the hills in this country have, from their general appearance, strong marks of containing those fossils which are inimical to vegetation, p. 95. Tincal, which is afterwards refined into Borax, is found in inexhaustible quantities in Thibet, p. 96. Rock salt found in great abundance in this country, *ibid.* Diseases of, p. 98—106. Method of preparing mercury in Thibet, p. 100. The physicians of this country do not treat the small-pox properly, their whole attention being to preserve themselves from the disease rather than to assist those who are infected with it, p. 105. All communication with those who have the small-pox strictly forbidden, *ibid.* Hot baths used by the inhabitants of Thibet in many disorders, *ibid.*

*Tincal*, found in inexhaustible quantities in Thibet, p. 96. The lake where tincal is collected about fifteen days journey from Tiffolumboo, *ibid.* Tincal deposited or formed in the bed of a lake; dug up in large masses, p. 97. Tincal is found only in shallow water, on the borders of the lake, *ibid.* Used in Thibet for solder, and to promote the fusion of gold and silver, *ibid.*

*Tiffolumboo*, productions of, p. 94. Soil, *ibid.* State of the thermometer at, pp. 97, 98. The capital of Thibet, p. 111.

*Trinidad*, account of a bituminous lake or plain in the island of, p. 65. In several parts of the woods in this island are hot springs, p. 69.

*Turkish mineral*, being mixed with sulphur, produces vitriolic acid air, p. 15. See *Sulphur*.

## V.

*Values*, &c. on the method of correspondent, p. 166. The method of correspondent values easily affords a resolution of the problems contained in Mr. Brigg's or Sir Isaac Newton's method of differences, p. 177.

*Veneral disease*, frequent in Thibet, p. 100. Method of treating that disorder in Thibet, p. 101.

*Vision*.

*Vision*, an attempt to explain a difficulty in the theory of, p. 256. Experiment illustrating the theory of vision, p. 261.

*Vitriol*, oil of, in its most dephlogisticated state, consists of a proper saturation of the acid with phlogiston, p. 289.

*Volatile vitriolic acid* contains the proper element of dephlogisticated air, p. 292.

*Volta's* opinion, that a point in electricity is the coating to an infinitely small plate of air, erroneous, p. 281.

## W.

*Walker*, Richard, on the congelation of quicksilver in England, p. 199.

*Waring*, Dr. on the method of correspondent values, &c, p. 166. On the resolution of attractive powers, p. 185.

*Water*, the decomposition of, by means of iron, a fallacy, p. 17. A great want of water on the close of the year 1788 universally felt, p. 37.

*Wolf*. See *Letter*.

---

FROM THE PRESS OF JOHN NICHOLS.

---















